














# NSW Childhood Immunisation Schedule

## Primary and Community Health



<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>6 Weeks and 4 Months</b></p>	<p>Diphtheria-tetanus-pertussis-hepatitis B-polio-<i>Haemophilus influenzae</i> type b</p>  <p>Infanrix Hexa or Vaxelis</p>	<p>Pneumococcal</p>  <p>Prevenar 13</p>	<p>Rotavirus</p>  <p>Rotarix</p>	<p>Meningococcal B (Aboriginal# children only)</p>  <p>Bexsero</p>
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>6 Months</b></p>	<p>Diphtheria-tetanus-pertussis-hepatitis B-polio-<i>Haemophilus influenzae</i> type b</p>  <p>Infanrix Hexa or Vaxelis</p>	<ul style="list-style-type: none"> <li>At risk children <math>\geq 6</math> months require an additional dose of Prevenar 13 (Pneumococcal). Refer to the online Australian Immunisation Handbook for the list of conditions</li> <li>Aboriginal# children <math>\geq 6</math> months with certain medical conditions may require an additional dose of Meningococcal B. Refer to the online Australian Immunisation Handbook for the list of conditions</li> </ul>		
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>12 Months</b></p>	<p>Meningococcal ACWY</p>  <p>Nimenrix</p>		<p>Pneumococcal</p>  <p>Prevenar 13</p>	<p>Measles-mumps-rubella</p>  <p>Priorix or MMR II</p>
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>18 Months</b></p>	<p>Measles-mumps-rubella-varicella</p>  <p>Priorix-Tetra</p>	<p><i>Haemophilus influenzae</i> type b</p>  <p>Act-HIB</p>	<p>Diphtheria-tetanus-pertussis</p>  <p>Infanrix or Tripacel</p>	
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>4 Years</b></p>	<p>Diphtheria-tetanus-pertussis-polio</p>  <p>Quadracel or Infanrix IPV</p> <div data-bbox="1291 2136 1743 2315" style="background-color: #800000; color: white; padding: 5px;"> <p>At risk children require a dose of Pneumovax 23 (pneumococcal) at 2-12 months after Prevenar 13 or at age 4 years <u>whichever is later</u>.</p> </div>			
<p style="writing-mode: vertical-rl; transform: rotate(180deg);"><b>*Influenza vaccines 6 months to less than 5 years</b></p>	<p>All children aged 6 months to less than 5 years are recommended to receive the annual influenza vaccine (NIP funded)*. Children aged 6 months to less than 9 years of age require 2 doses at least 4 weeks apart in their first year of influenza vaccination.</p>  <p>Fluarix Tetra or Vaxigrip Tetra</p>			

# The term Aboriginal is inclusive of Aboriginal and Torres Strait Islander people. \*NIP funded influenza vaccines for all children 6 months <5 years. October 2023 © NSW Health. SHPN (HP NSW) 230818.