


FEVER OR SUSPECTED INFECTION IN PAEDIATRIC ONCOLOGY & STEM CELL TRANSPLANTATION PATIENTS

Minimum Triage Category 2 for patients presenting to the ED

For the following patients with fever or reported fever $\geq 38.0^{\circ}\text{C}$ or who are unwell

- Patients on treatment for cancer
- Patients who ceased treatment for cancer within the last 3 months
- Recipients of Stem Cell Transplantation (SCT) within the last 12 months or on immunosuppressive therapy
- Oncology or SCT patients with Central Venous Access Device (CVAD) in situ

Do Not Wait for local anaesthetic to take effect. (e.g. to access port or insert peripheral line)
Access CVAD or establish peripheral IV and collect Blood cultures, FBC, VBG (Lactate & glucose) EUC, LFT
DO NOT WAIT FOR BLOOD RESULTS TO START ANTIBIOTICS



Inform Paediatrician as per local CERS and Oncologist on call as soon as possible
Discuss the management plan with the patient and family

[^]Indications for Vancomycin: Obviously infected intravascular devices (erythema/tenderness along subcutaneous track or purulent exit site discharge), MRSA carriers with clinical instability, high dose cytarabine recipients (>2gm/m²) with clinical instability #Patients with penicillin allergy: refer to table 1 for initial antibiotic choice
October 2015 Page 1

Table 1 First Dose Empiric Antibiotics

TREATMENT		Clinically Stable	Clinically Unstable	Severe Sepsis/Shock
EMPIRICAL ANTIBIOTIC REGIMEN*		Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV (max. dose 320 mg) Single dose only then Piperacillin + Tazobactam 100 mg/kg/dose IV 6 hourly (max. dose 4 g Piperacillin component) ADD Vancomycin [^] if clinically indicated	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg) then Piperacillin + Tazobactam 100 mg/kg/dose IV 6 hourly (max. dose 4 g Piperacillin component) ADD Vancomycin [^] if clinically indicated	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg) then Piperacillin + Tazobactam 100 mg/kg/dose IV 6 hourly (max. dose 4 g Piperacillin component) AND Vancomycin 15 mg/kg/dose IV 6 hourly (max. dose 750 mg)
ALLERGY	Non-life threatening penicillin hypersensitivity	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV (max. dose 320 mg) Single dose only AND Cefepime 50 mg/kg/dose IV 8 hourly (max. dose 2 g)	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg). AND Cefepime 50 mg/kg/dose IV 8 hourly (max. dose 2 g) OR Meropenem 40 mg/kg/dose IV 8 hourly (max. dose 2 g)	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg). AND Cefepime 50 mg/kg/dose IV 8 hourly (max. dose 2 g) OR Meropenem 40 mg/kg/dose IV 8 hourly (max. dose 2 g) AND Vancomycin 15 mg/kg/dose IV 6 hourly (max. dose 750 mg)
	Life-threatening Penicillin Hypersensitivity not known to tolerate Cephalosporins /Meropenem safely	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV (max. dose 320 mg) Single dose only AND Ciprofloxacin 10 mg/kg/dose IV 8 hourly (max. dose 400 mg) AND Vancomycin 15 mg/kg/dose IV 6 hourly (max. dose 750 mg)	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg) AND Ciprofloxacin 10 mg/kg/dose IV 8 hourly (max. dose 400 mg) AND Vancomycin 15 mg/kg/dose IV 6 hourly (max. dose 750 mg)	Gentamicin (dose based on lean body weight for obese patients) 7.5 mg/kg/dose IV 24 hourly (max. dose 320 mg) AND Ciprofloxacin 10 mg/kg/dose IV 8 hourly (max. dose 400 mg) AND Vancomycin 15 mg/kg/dose IV 6 hourly (max. dose 750 mg)
MODIFICATIONS		Add Metronidazole 12.5 mg/kg/dose IV 8 hourly (max. dose 500 mg) if Cefepime or Ciprofloxacin used AND evidence of abdominal/perineal infection	Use Meropenem rather than Cefepime or Piperacillin/Tazobactam if colonised with a multi-resistant GNR (e.g. ESBL) Add Metronidazole 12.5 mg/kg/dose IV 8 hourly (max. dose 500 mg) if Cefepime or Ciprofloxacin used and evidence of abdominal/ perineal infection	Use Meropenem rather than Cefepime or Piperacillin/Tazobactam if colonised with a multi-resistant GNR (e.g. ESBL) Add Metronidazole 12.5 mg/kg/dose IV 8 hourly (max. dose 500 mg) if Cefepime or Ciprofloxacin used AND evidence of abdominal/perineal infection

All Antibiotic doses are based on actual body weight except Gentamicin. Gentamicin: Dose based on lean body weight for obese patients – see *appendix 3* for method for calculating lean body weight in obese children. **Administer over 5 minutes.** Ensure that line is flushed with 10-20 mL following Gentamicin and prior to any further doses of antibiotics. **Piperacillin-Tazobactam: Administer over 20- 30 mins. Vancomycin: Administer over at least 60 mins.** If patient has previously experienced ‘red man syndrome’ administer over 2 hours. **^ Indications for Vancomycin:** Obviously infected vascular devices (erythema/tenderness along subcutaneous track or purulent exit site discharge), MRSA carriers with clinical instability, High dose Cytarabine (>2gm/m²/day) recipients with clinical instability. For clinically stable patients the decision to continue Gentamicin beyond the first dose must be made after discussing with treating oncologist, but is not recommended by the Australian Therapeutic Guidelines – Antibiotic 2015 due to the lack of proven benefit and potential for toxicity. Subsequent antibiotic choice/dose (i.e. after first dose) may need modification based on patient’s renal function, clinical stability and history of colonisation with multi-drug resistant organisms. These decisions must be made after discussing with treating oncologist. For patients continuing Gentamicin, drug level must be monitored just prior to second dose. For patients continuing Vancomycin, drug level must be monitored just prior to 5th dose.