

## Children and Infants with Gastroenteritis - Acute Management

**Document Number** PD2010\_009

**Publication date** 03-Feb-2010

**Functional Sub group** Clinical/ Patient Services - Baby and child

**Summary** Clinical Practice Guidelines for the treatment of infants and children with gastroenteritis.

**Replaces Doc. No.** Children and Infants with Gastroenteritis - Acute Management [PD2009\_064]

**Author Branch** Statewide Services Development

**Branch contact** Trish Boss 9424 5706

**Applies to** Area Health Services/Chief Executive Governed Statutory Health Corporation, Board Governed Statutory Health Corporations, Affiliated Health Organisations - Non Declared, Affiliated Health Organisations - Declared, Public Health System Support Division, Community Health Centres, NSW Ambulance Service, Public Hospitals

**Audience** Emergency Departments, Paediatric Units

**Distributed to** Public Health System, Divisions of General Practice, NSW Ambulance Service, Private Hospitals and Day Procedure Centres, Tertiary Education Institutes

**Review date** 03-Feb-2013

**Policy Manual** Patient Matters

**File No.** 06/3557

**Status** Active

### Director-General

This Policy Directive may be varied, withdrawn or replaced at any time. Compliance with this directive is **mandatory** for NSW Health and is a condition of subsidy for public health organisations.

## INFANTS AND CHILDREN: ACUTE MANAGEMENT OF GASTROENTERITIS

### PURPOSE

The *infants and children: acute management of gastroenteritis* clinical practice guideline (attached) has been developed to provide direction to clinicians and is aimed at achieving the best possible paediatric care in all parts of the state.

The clinical practice guideline was prepared for the NSW Department of Health by an expert clinical reference group under the auspice of the state wide Paediatric Clinical Practice Guideline Steering Group.

### MANDATORY REQUIREMENTS

This policy applies to all facilities where paediatric patients are managed. It requires all Health Services to have local guidelines/protocols based on the attached clinical practice guideline in place in all hospitals and facilities likely to be required to assess or manage children with gastroenteritis.

The clinical practice guideline reflects what is currently regarded as a safe and appropriate approach to the acute management of gastroenteritis in infants and children. However, as in any clinical situation there may be factors which cannot be covered by a single set of guidelines. This document should be used as a guide, rather than as a complete authoritative statement of procedures to be followed in respect of each individual presentation. **It does not replace the need for the application of clinical judgement to each individual presentation.**

### IMPLEMENTATION

Chief Executives must ensure:

- Local protocols are developed based on the *infants and children: acute management of gastroenteritis* clinical practice guideline.
- Local protocols are in place in all hospitals and facilities likely to be required to assess or manage paediatric patients with gastroenteritis.
- Ensure that all staff treating paediatric patients are educated in the use of the locally developed paediatric protocols.

Directors of Clinical Governance are required to inform relevant clinical staff treating paediatric patients of the revised protocols.

### REVISION HISTORY

Version	Approved by	Amendment notes
December 2004 (PD2005_238)	Director-General	New policy
October 2009 (PD2009_064)	Deputy Director-General Population Health	Second edition
February 2010 (PD2010_009)	Deputy Director-General Population Health	Third edition. Corrects table on page 8.

### ATTACHMENT

1. Infants and Children: Acute Management of Gastroenteritis – Clinical Practice Guideline.

# Infants and children: Acute Management of Gastroenteritis

third edition

**CLINICAL PRACTICE GUIDELINES**



**NSW HEALTH**

NSW DEPARTMENT OF HEALTH

73 Miller Street

North Sydney NSW 2060

Tel. (02) 9391 9000

Fax. (02) 9391 9101

[www.health.nsw.gov.au](http://www.health.nsw.gov.au)

This work is copyright. It may be reproduced in whole or part for study or training purposes subject to the inclusion of an acknowledgement of the source. It may not be reproduced for commercial usage or sale. Reproduction for purposes other than those indicated above requires written permission from the NSW Department of Health.

NSW Department of Health 2009

SHPN: (SSD) 090178

ISBN: 978-1-74187-453-2

For further copies of this document please contact:

Better Health Centre – Publications Warehouse

PO Box 672

North Ryde BC, NSW 2113

Tel. (02) 9887 5450

Fax. (02) 9887 5452

Information Production and Distribution

Tel. (02) 9391 9186

Fax. (02) 9391 9580

E-mail: [bhc@nscchahs.nsw.gov.au](mailto:bhc@nscchahs.nsw.gov.au)

Further copies of this document can be downloaded from the

NSW Health website: [www.health.nsw.gov.au](http://www.health.nsw.gov.au)

A revision of this document is due in 2011.

November 2009 - third edition

# Contents

<b>Introduction</b> .....	<b>3</b>
<b>Summary</b> .....	<b>4</b>
Significant Changes from 2002 CPG Version .....	4
<b>Gastroenteritis in Infancy and Childhood</b> .....	<b>5</b>
Principles of Fluid Management .....	6
Medications.....	6
Differential Diagnoses.....	7
Table 1: Clinical Assessment of Dehydration and Initial Treatment.....	8
Management Algorithm .....	9
<b>Enteral Rehydration Therapy</b> .....	<b>10</b>
Oral Rehydration Solutions (ORS) .....	10
Method of Giving Oral Fluids .....	12
Discharge Criteria .....	12
<b>Nasogastric Rehydration Therapy</b> .....	<b>13</b>
<b>Intravenous Fluid Therapy</b> .....	<b>15</b>
Introductory Notes .....	15
Resuscitation .....	15
Rapid IV Rehydration .....	16
Standard IV Rehydration .....	17
Hypernatraemia .....	19
Hyponatraemia .....	19
<b>Investigations and Observations</b> .....	<b>20</b>
Reintroduction of Diet .....	21

References.....22

Bibliography .....24

Appendices .....29

    Appendix One – Glossary..... 29

    Appendix Two – IVT Composition.....30

    Appendix Three – Parent Oral Rehydration Documentation Form ..... 31

    Appendix Four – Parent Information ..... 33

    Appendix Five – Resources..... 34

    Appendix Six – Significant Changes From 2002 CPG Version ..... 35

    Appendix Seven – Alternative Calculation for Maintenance Fluids ..... 35

    Appendix Eight – Working Party Members..... 36

# Introduction

These Guidelines are aimed at achieving the best possible paediatric care in all parts of the State. The document should not be seen as a stringent set of rules to be applied without the clinical input and discretion of the managing professionals. Each patient should be individually evaluated and a decision made as to appropriate management in order to achieve the best clinical outcome.

The formal definition of clinical practice guidelines comes from the National Health and Medical Research Council:

'Systematically developed statements to **assist** practitioner and patient decisions about appropriate health care for specific clinical circumstances.'

*(National Health and Medical Research Council 'A Guide to the Development, implementation and evaluation of Clinical Practice Guidelines', Endorsed 16 November 1998, available from [www.nhmrc.gov.au/publications/synopses/cp30syn.htm](http://www.nhmrc.gov.au/publications/synopses/cp30syn.htm))*

It should be noted that this document reflects what is currently regarded as a safe and appropriate approach to care. However, as in any clinical situation there may be factors which cannot be covered by a single set of guidelines. This document

should be used as a guide, rather than as a complete authoritative statement of procedures to be followed in respect of each individual presentation. It does not replace the need for the application of clinical judgment to each individual presentation.

This document represents basic clinical practice guidelines for the acute management of gastroenteritis in children and infants. Further information may be required in practice; suitable widely available resources are included as Appendix Five.

Each Area Health Service is responsible for ensuring that local protocols based on these guidelines are developed. Area Health Services are also responsible for ensuring that all staff treating paediatric patients are educated in the use of the locally developed paediatric guidelines and protocols.

In the interests of patient care it is critical that contemporaneous, accurate and complete documentation is maintained during the course of patient management from arrival to discharge.

**Parental anxiety should not be discounted: it is often of significance even if the child does not appear especially unwell.**

# Summary

Gastroenteritis is a common paediatric condition. Appropriate management attempts to avoid dehydration, but when this occurs appropriate fluid management is essential. **For rehydration the enteral route is preferred**, but if intravenous fluids are used then low sodium containing fluids must be avoided. For intravenous rehydration 0.9% Normal Saline (NaCl) + 2.5% Glucose is preferred. 0.9% Normal Saline (NaCl) + 5% Glucose may also be used. If rehydrating over 24 hours then 0.45% NaCl + 2.5% Glucose is an acceptable alternative. Oral rehydration solutions may be offered orally or administered nasogastrically.

In cases of severe dehydration or clinical deterioration after admission or despite treatment, the Admitting Medical Officer in charge or consulting paediatrician should be notified and should personally review the patient as soon as possible. Where other medical staff act as a delegate for the Admitting Medical Officer, the hospital must have clear written protocols defining this arrangement.

For hospitals employing junior medical staff: **The Admitting Medical Officer MUST be notified within an hour of the decision to admit the child.** Details of the physical findings and proposed fluid therapy should

be discussed. Hospitals should have an internal policy that defines roles if senior registrars act as a delegate for the AMO.

These Clinical Practice Guidelines should be read in conjunction with other relevant Clinical Practice Guidelines (e.g. the *Recognition of a Sick Child in Emergency Departments*, and *Acute Management of Infants and Children with Acute Abdominal Pain*). When dealing with children suspected of having gastroenteritis, **it is essential that infection control measures be implemented** to prevent cross-contamination and spread.

# Gastroenteritis in Infancy and Childhood

- n This common acute intestinal communicable infection causes vomiting, diarrhoea and fever. It is usually viral, but sometimes bacterial or parasitic. Community outbreaks are sporadic and seasonal.
- n A small proportion of those affected will suffer severe dehydration and electrolyte disturbance. Untreated or poorly treated dehydration may progress to shock and death. There are also risks from over-hydration and/or inappropriate electrolyte replacement, including death from cerebral oedema.
- n Some other serious illnesses are sometimes incorrectly diagnosed as gastroenteritis. **Warning signs of other diagnoses must be recognised and investigated** (see page 7).
- n Availability of standard resuscitation intravenous fluids, including 0.9% NaCl (without added glucose) or Hartmann's solution (without added glucose).
- n Availability of rehydration intravenous fluids, including 0.9% NaCl + 2.5% Glucose, 0.45% NaCl + 2.5% Glucose.
- n Intravenous paediatric giving sets with burettes, appropriate infusion pumps.
- n Appropriate Oral Rehydration Solutions such as Gastrolyte®, Gastrolyte-R®, Repalyte®, Hydralyte®
- n Appropriate giving sets and enteral infusion pumps (e.g. Kangaroo® pump).

## Suggested hospital requirements for management of children with gastroenteritis

- n 24-hour availability of nurses and medical practitioners experienced in the management of sick children.
- n Access to 24-hour standard biochemistry for inpatient management. This may include point of care testing.

## Availability of assistance when treating severely ill children

- n The treatment of children with severe dehydration should be discussed with a paediatrician and consideration be given to transfer to a facility with a paediatric intensive care unit.
- n For advice regarding the management of seriously ill children or to arrange their transfer to any of the children's hospitals contact NSW Newborn and Paediatric Emergency Transport Service **(NETS) Hotline number: 1300 36 2500.**

Calls to NETS are voice recorded and form part of the NETS medical record for the patient.

## Principles of Fluid Management

- n Infants and children with gastro-enteritis require additional fluids to prevent dehydration, or for rehydration.
- n The **enteral route is preferred** for rehydration of children with mild or moderate dehydration. This is with an Oral Rehydration Solution (ORS) either by mouth or via nasogastric tube.
- n Suitable fluids should be offered, for oral rehydration
  - **Babies** who are breastfed should receive small frequent breastfeeds to ensure normal urine output. This may be supplemented with an ORS.
  - **For all other children**, offer an ORS. Those requiring mixing must follow the manufacturer's instructions. *Do not add flavouring or sweet drinks to an ORS.*
  - If an ORS is unavailable, or refused, dilute juice/lemonade (mixed as 1 part juice/lemonade with 4 parts water) can be used **only if a child is not dehydrated**. *These are less desirable fluid options.*
  - Do not use 'sports drinks' as they are not an appropriate rehydration fluid for children with gastroenteritis.
  - Do not use low-calorie or diet drinks.
- n Suitable volumes should be offered: try to give about 0.5mL/kg every 5 minutes.
- n Achieving successful oral rehydration demands constant attention and persistence, usually by parents.
- n The principles and practice of oral replacement therapy are described on page 10.
- n Intravenous rehydration is often a reasonable alternative for moderate dehydration (see Table 1 on page 8) and is essential where severe dehydration and/or shock are present.
- n Children receiving fluid rehydration require regular timely reassessment.
- n The principles and practice of intravenous replacement therapy are described on page 15.
- n **NB: Careful calculations of fluid volume and rate are required regardless of route of administration.**

## Medications

There are no indications for using anti-motility or anti-diarrhoeal agents in the management of acute gastroenteritis in infants or children.

Many antiemetic medications have a risk of significant side effects, like dystonic reactions and sedation, and should be avoided [e.g. promethazine, prochlorperazine]. Medications such as 5HT-3 receptor antagonists, such as ondansetron, may have some clinical benefit, however the evidence is not conclusive. Experienced clinicians choosing

to use that medication generally should limit the use to a single dose.

Pro-biotics and Zinc may have some clinical benefits in the management of gastroenteritis, and may be available in some commercially available products such as yoghurts. These can be given to children when a normal diet is reintroduced.

Antibiotics are rarely required in gastroenteritis, even when bacterial in aetiology. If unsure, consult a paediatrician or paediatric infectious disease specialist.

Rotavirus vaccines are available and have a significant benefit in the prevention of gastroenteritis in young infants. Additional information is available at the National Centre for Immunisation Research and Surveillance of Vaccine Preventable Diseases website

[http://www.ncirs.usyd.edu.au/facts/rotavirus\\_vaccine\\_for\\_children\\_june\\_2007.pdf](http://www.ncirs.usyd.edu.au/facts/rotavirus_vaccine_for_children_june_2007.pdf)

## Differential Diagnoses

Always keep in mind the possibility that the diagnosis of gastroenteritis could be incorrect. *Gastroenteritis consists of the triad of vomiting, diarrhoea and fever.* Be cautious of evaluating the child with vomiting alone. The following list is not exclusive. Consider also:

- n Acute appendicitis
- n Strangulated hernia
- n Intussusception or other causes of bowel obstruction

- n Urinary tract infection
- n Meningitis and other types of sepsis
- n Any cause of raised intracranial pressure
- n Diabetic ketoacidosis
- n Inborn errors of metabolism
- n Inflammatory bowel disease
- n Haemolytic uraemic syndrome

Always consider another diagnosis if there is:

- n Abdominal distension
- n Bile-stained vomiting
- n Fever >39°C
- n Blood in vomitus or stool
- n Severe abdominal pain
- n Vomiting in the absence of diarrhoea
- n Headache

**Beware! The very young infant and the malnourished child are more likely to suffer severe disease, or to have another diagnosis.**


Table 1 on page 8 gives an overview of dehydration definition, signs and symptoms, along with initial enteral or parenteral fluid therapy. Wherever 0.9% NaCl + 2.5% Glucose is advised, 0.9% NaCl + 5% Glucose would be an acceptable alternative. The flowchart on page 9 outlines a treatment overview and highlights decision points in regard to the initial management of an infant or child with gastroenteritis.

**Table 1: Clinical Assessment of Dehydration and Initial Treatment**

No single symptom or clinical sign reliably predicts the degree of dehydration

Dehydration (% of Body Weight)		Signs and Symptoms	Replacement Fluid Route	Replacement Fluid Type
No Clinical Signs of Dehydration		Reduced urine output	Oral	In order of preference 1. Frequent breastfeeds where appropriate/possible 2. Oral Rehydration Solution (see page 10) 3. 1/5 strength clear fluids i.e.: 4 parts water and 1 part juice/lemonade (if an ORS refused)
		Thirst		
Mild	3%	No physical signs	Oral	In order of preference 1. Frequent breastfeeds where possible/appropriate may be supplemented with an ORS 2. Oral Rehydration Solution (see page 10)
		Reduced urine output		
Moderate		Thirst	Nasogastric	Oral Rehydration Solution e.g. Gastrolyte® (see p. 10)
		Dry mucous membranes	Intravenous	0.9% NaCl + 2.5% Glucose
		Tachycardia	- Rapid	0.9% NaCl + 2.5% Glucose <b>or</b> 0.45% NaCl + 2.5% Glucose
Severe	5%	Abnormal respiratory pattern	Nasogastric	1. Oral Rehydration Solution e.g. Gastrolyte® (see page 10)
		Lethargy	Intravenous	0.9% NaCl + 2.5% Glucose
		Reduced skin turgor	-Rapid	0.9% NaCl + 2.5% Glucose <b>or</b> 0.45% NaCl + 2.5% Glucose
Severe	10%	Sunken eyes	Intravenous or intraosseous	Use either: 0.9% NaCl or Hartmann's solution for resuscitation and then reassess the child. Ongoing fluid replacement should be with either: 0.9% NaCl + 2.5% Glucose <b>or</b> 0.45% NaCl + 2.5% Glucose
		Above signs	20mL/kg stat and reassess fluid needs	
		Poor Perfusion: Mottled, cool limbs/Slow capillary refill/Altered consciousness		
		Shock: thready peripheral pulses with marked tachycardia and other signs of poor perfusion stated above		

# Management Algorithm



# Enteral Rehydration Therapy

Most children with gastroenteritis and mild- moderate dehydration can be successfully rehydrated with oral rehydration solutions either by mouth or nasogastric tube.

## Oral Rehydration Solutions

Oral rehydration solutions (ORS) are specifically designed fluids that contain an appropriate amount of sodium, glucose and other electrolytes and are of the appropriate osmolality, to maximise water absorption from the gut. They use the principle of glucose-facilitated sodium transport whereby glucose enhances sodium and secondarily water transport across the mucosa of the upper intestine. The sodium and glucose concentrations and the osmolality are of vital importance.

The World Health Organisation (WHO) recommends an ORS that has a sodium concentration of 90mmol/L. In developed countries with non-cholera diarrhoea, it is generally thought that 90mmol/L is a little high, as non-cholera gastroenteritis does not result in the same sodium losses that are seen in cholera.

Many different ORS with varying sodium concentrations have been developed. It has been shown that water absorption across the lumen of the human intestine is

maximal using solutions with a sodium concentration of 60mmol/L<sup>1</sup> (such as Gastrolyte®) and this is the concentration recommended by the European Society of Paediatric Gastro-enterology and Nutrition.<sup>2</sup> However some children who are not particularly dehydrated will refuse to drink such an ORS because of its salty taste. ORS with slightly less sodium such as Hydralyte® may be more palatable, particularly as this comes in an iceblock form.

ORS with similar compositions to Hydralyte® are safe and effective. These hypo-osmolar solutions (such as Gastrolyte® and Hydralyte®) are more effective at promoting water absorption than isotonic or hypertonic solutions.<sup>3,4,5</sup>

The composition of various ORS and other fluids is shown in Tables 2 and 3. Fruit juices and soft drinks are inappropriate because of the minimal sodium content and the excessive glucose content and hence excessive osmolality, which will worsen diarrhoea. Although diluting juices and soft drinks reduces glucose concentration, the fluid has insufficient sodium to act as a rehydration fluid.

Sports drinks have varying sodium and carbohydrate levels, and are considered inappropriate as rehydration solutions.

**Table 2: Composition of Oral Rehydration Solutions**

Comparisons of ORS					
	Na	Carbohydrate			Osmolality
	(mmol/L)		(mmol/L)	(%)	(mOsm/L)
WHO	90	G	111	(2%)	331
Gastrolyte®	60	G	90	(2%)	240
Gastrolyte-R®	60	RSS	6g/L	(2.5%)	226
Repalyte®	60	G	90	(2%)	240
Terry White/Chem-mart®	60	G	90	(2%)	240
Hydralyte®	45	G	90	(2.5%)	240
G = glucose, RSS = rice syrup solids					

**Table 3: Composition of Oral Fluids**

Comparisons of Oral Fluids			
	Na <sup>+</sup>	Carbohydrate	Osmolality
	(mmol/L)	(mmol/L)	(mOsm/L)
Apple Juice	3	690	730
Soft drinks	~2	~700	~750
Sports drinks	~20	~255	~330

## Method of Giving Oral Fluids

It is important to give small amounts of fluid frequently, for example 0.5mL/kg every five minutes. The fluid can be measured in a syringe and given to the child either by syringe, teaspoon or cup. The child is far more likely to tolerate these small amounts of fluid than if he/she drinks a large amount at once.

Obviously if the child tolerates this fluid the parent can gradually increase the volume and decrease the frequency of the fluid offered. Success can be optimised in the Emergency Department setting by giving the parents a documentation chart (see Appendix 3) to record the fluid given and any vomits, diarrhoea or urine passed.

It is important to educate the parents that seeing a doctor will not cure their child of his/her vomiting and diarrhoea. Small, frequent amounts of fluid will hopefully minimise the vomiting, but will not reduce the diarrhoea. The aim is for the input to exceed the output by enough to rehydrate and then maintain hydration.

**Occasional vomiting alone should not be considered as failure of oral rehydration therapy.**

## Discharge Criteria

Children with gastroenteritis can be discharged, even if they still have some vomiting, if the following discharge criteria are met:

1. Diagnosis of gastroenteritis
2. Child is rehydrated or only mildly dehydrated
3. Gastrointestinal losses not profuse
4. Child has passed urine in ED or within the last 4 hours
5. Parent has demonstrated the ability to give an ORS appropriately
6. Clinical staff confident parent will take child to GP for review within 48hrs and represent for medical review if child's condition deteriorates.
7. If a child with gastroenteritis and dehydration does not fulfil the discharge criteria, they will need to be admitted for ongoing management.

# Nasogastric Rehydration Therapy

Increasing numbers of hospitals in developed countries are using an ORS via continuous nasogastric (NG) infusion.<sup>8,9</sup> This has been shown to be as effective as intravenous rehydration,<sup>8-15</sup> less expensive<sup>8,15</sup> and reduces lengths of hospital stay<sup>8</sup> when compared with standard intravenous rehydration. It is usually unnecessary to perform EUC for children being rehydrated with nasogastric ORS.

Nasogastric rehydration is where an ORS is infused continuously via a nasogastric tube with a pump such as a Kangaroo® pump. **Choose an ORS with a sodium concentration of 60mmol/L such as Gastrolyte®, as this is the optimal concentration and taste is not an issue when using an NG tube.** Hydralyte® has only 45 mmol/L of sodium, and is not the preferred nasogastric solution (but when given orally often has better compliance due to taste). Nasogastric rehydration is often successful even in children with frequent vomiting. Staff need to be competent in placing nasogastric tubes in children and babies and follow local protocols and training/accreditation procedures. Facilities need to be equipped to deliver NG rehydration in regards to equipment and education before this form of rehydration should be introduced.

## **Do not use nasogastric rehydration if the child has:**

- n an ileus (check for bowel sounds)
- n significantly reduced level of consciousness

## **Do not use 'rapid' nasogastric rehydration if child:**

- n is younger than 6 months old
- n has a medical condition which increases the risk of fluid overload

Different regimens are used for continuous nasogastric rehydration. One simple method is described below:

- n Perform observations – temperature (T), pulse (P), respiratory rate (R), blood pressure (BP) and mental state – before commencing, then repeat T, P, and R at least hourly.
- n Establish NG access
- n Give 10mL/kg/hour of Gastrolyte® over four hours for all mild-moderately dehydrated children, **after which the infusion is ceased.**
- n Do not take blood for EUC and BGL as a routine.
- n After completion, the child is then re-examined by the medical officer and a re-trial of oral fluids is commenced.

After 1–2 hours of completing rehydration the child is reassessed and if he/she fulfils the discharge criteria (page 12), may be discharged with appropriate advice and follow-up.

If the child does not tolerate NG rehydration, (NB: 1 or 2 small vomits does not necessarily mean NG rehydration has failed), IV rehydration will probably need to be commenced. **This should be over 24 hours i.e. do not give rapid IV rehydration after the child has already received 'rapid' NG rehydration** (see page 16 for calculation).

Some clinicians may choose to use a lower NG infusion rate, either initially or over a longer period of time; similar to the standard IV rehydration rate (see page 17).

If the child tolerated the NG rehydration but fails the subsequent trial of oral fluids, the child will usually need to stay in hospital. If further fluid in addition to that taken orally is required after reassessment, this can be given via the nasogastric tube. **A second administration of 'rapid' NG rehydration should not be given** (see page 18 for calculation). At this point, EUC and BGL should be checked to ensure that an electrolyte abnormality is not the cause for failure of rehydration.

NG rehydration is most suitable for infants and young children. From a practical viewpoint, older children would be more suitable to be rehydrated orally or intravenously.

# Intravenous Fluid Therapy

## Introductory notes

- n When IV fluid therapy is commenced, the first decision is whether the child is severely dehydrated and needs resuscitation (see below).
- n When resuscitation fluid (a bolus) has been given for shock, and the signs of shock corrected, then the next phase of treatment is to provide standard IV rehydration.
- n If a bolus has not been needed, the next phase of treatment is to provide either rapid IV rehydration or standard IV rehydration.
- n In this Guideline, 0.9% NaCl + 2.5% Glucose is considered to be an isotonic solution, as the glucose is rapidly metabolised after infusion.
- n Whenever an IV cannula is inserted for the provision of IV fluids, blood should be withdrawn and sent for EUC and BGL.
- n Children with gastroenteritis and dehydration are at risk of hypoglycaemia; any fluid used for rehydration should contain some glucose. For hypoglycaemic children (i.e. BGL <2.6 mmol/L), treat with **2–5mL/kg** of 10% Glucose and recheck BGL within 20–30 minutes.

Check urinalysis for ketones. Most healthy children who have ketonuria and hypoglycaemia in this situation will be due to starvation. Consult a paediatrician and discuss any need for further investigations.

- n For standard IV rehydration therapy (**not** rapid IV rehydration – see below) it is acceptable to use **either** 0.9% NaCl + 2.5% Glucose **or** 0.45% NaCl + 2.5% Glucose. **DO NOT USE ANY FLUID CONTAINING LESS SODIUM THAN 0.45% NaCl.**

When ordering IV fluids it is preferable to write:

'0.9% NaCl + 2.5% Glucose' rather than 'N/S + 2.5% Glucose',

or '0.45% NaCl + 2.5% Glucose', rather than 'N/2 Saline + 2.5% Glucose'

to avoid confusion between 'N/S' and 'N/2' and subsequent error.

## Resuscitation

- n Where severe dehydration is accompanied by shock or imminent shock, bolus administration of intravenous fluids for resuscitation is required. Commence with intraosseous infusion if IV access cannot be

established. If neither is possible commence nasogastric rehydration while awaiting assistance. These administration routes do not negate the need for an IV line.

- n 0.9% NaCl or Hartmann's solution, 20 mL/kg, should be given IV/IO initially over 10–20 minutes. These fluids should NOT contain Glucose.
- n For hypoglycaemic children (i.e. BGL <2.6 mmol/L), treat with 2–5mL/kg of 10% Glucose and recheck BGL within 20–30 minutes.
- n Check serum electrolytes.
- n Repeat boluses of 10–20 mL/kg until signs of shock are reversed.
- n After signs of shock are reversed, proceed to **standard IV rehydration** (see page 16).

## Rapid IV Rehydration

- n Rapid rehydration refers to the correction of dehydration over a relatively short time e.g. 4 hours, with the expectation that the child may subsequently be able to be discharged from hospital if tolerating oral fluids. This method has been shown to be safe and effective.<sup>16</sup>

**THE IV FLUIDS MUST NOT BE CONTINUED AT THE RAPID RATE BEYOND 4 HOURS.**

- n All children should be given a trial of oral fluids during or after rapid IV rehydration has finished. IV fluids should only be recommenced after careful reassessment of the child and if the child still has signs of dehydration,

or there are significant ongoing losses. These should be given at standard rehydration rates only (i.e. rapid IV rehydration should NOT be repeated).

- n The advice on rapid IV rehydration in this guideline applies only to those children with gastroenteritis.
- n **Do not use rapid IV rehydration if:**
  - *The patient is less than 6 months old*
  - *The patient is severely dehydrated (10%) or shocked*
  - *The patient has an altered level of consciousness*
  - *The serum sodium, if known, is <130 mmol/L or >149 mmol/L.*

- n Establish IV access. Send blood for EUC and BGL. Also checking the BGL by bedside meter is good clinical practice. Results should be available and checked within 1 hour.

**IF LABORATORY ACCESS TO THIS TURN AROUND IS NOT AVAILABLE THEN DO NOT USE RAPID IV REHYDRATION.**

- n Perform observations – T,P,R, BP and assessment of mental state – before commencing rapid IV rehydration , then repeat T,P,R at least hourly.
- n Commence 0.9% NaCl + 2.5% Glucose at 10 mL/kg/hr and continue for 4 hours. 0.9% NaCl + 5% Glucose is an acceptable alternative. **0.45% NaCl + 2.5% Glucose must not be used for rapid IV rehydration.**
- n After 4 hours, cap IV cannula. The treating doctor should re-examine the child, and oral fluids should be offered.

Body Weight	Fluid requirement mL/ day	Fluid requirement mL/ hour
First 10 kg	100 mL/kg	4 mL/kg/hr
Second 10 kg	+ 50 mL/kg	+ 2 mL/kg/hr
Subsequent kg	+ 20 mL/kg	+ 1 mL/kg/hr

Small amounts of oral fluids may be commenced earlier, during IV rehydration, if appropriate. Intravenous fluids should only be recommenced if on review the child still has signs of dehydration, or if there are significant ongoing losses.

- n If the rapid rehydration has been completed late at night, and the child has no signs of dehydration, the child may safely be allowed to sleep with the IV capped, with a trial of oral fluids to commence when the child wakes.
- n It is expected that the clinical status (e.g. heart rate, perfusion and mental state) of the child receiving rapid IV rehydration for gastroenteritis should gradually improve. Failure to improve, any deterioration, or the development of unexpected signs or symptoms should lead to a reconsideration of the diagnosis and management, and discussion with a Paediatrician.

## Standard IV Rehydration

- n Standard rehydration refers to the provision of maintenance fluids, and the correction of dehydration, usually over 24 hours.
- n When the IV cannula is inserted, send blood for EUC and BGL.
- n It is acceptable to commence fluid therapy with either 0.9% NaCl + 2.5% Glucose or 0.45% NaCl + 2.5% Glucose for standard IV rehydration, pending EUC results. 0.9% NaCl + 5% Glucose and 0.45% NaCl + 5% Glucose are acceptable alternatives.
- n Calculate the total volume of IV fluids likely to be needed for the next 24 hours, being maintenance and deficit. Do not include fluids already given for resuscitation. If resuscitation has been required, reassess dehydration state now and calculate fluid requirement from now. Note that this calculation is a 'starting point' which will be reviewed according to progress, including assessment of general appearance, heart rate, urine output, ongoing losses (vomiting, diarrhoea), or fever, at intervals of not more than 6 hours.

- n The majority of children will not require rehydration for more than a 5% deficit in the first 24 hours.
- n The volumes for rehydration and maintenance are calculated separately, as the basis for calculation of each is different.
- n For **rehydration**:  $\text{Weight (in kg)} \times \% \text{ dehydration} \times 10 = \text{mL deficit needed for rehydration}$ . For example, a 9 kg child estimated to be 5% dehydrated, rehydration volume is  $9 \times 5 \times 10 = 450 \text{ mL}$ . (see Table 1 on page 8 for a guide to assessing dehydration)
- n For **maintenance**: Calculate volume according to the child's weight, as in the table (page 17):  
For example: A child weighing 25 kg has a maintenance fluid requirement for 24 hours of:  
 $(10 \times 100) + (10 \times 50) + (5 \times 20) = 1600 \text{ mL per 24 hours}$ .  
Note:
  - Maintenance fluids for infants less than 6–9 months is 120mL/kg
  - There are alternative methods for calculating maintenance fluid requirements (see Appendix 7 for an example).
- n Add the two volumes (rehydration + maintenance) together. Calculate the rate to give the total volume over 24 hours. Start by giving 0.9% NaCl + 2.5% Glucose or 0.45% NaCl + 2.5% Glucose over the first 6-8 hours, then review the child's progress.

**DO NOT USE LOW SODIUM-CONTAINING FLUIDS** (any fluid with less sodium than 0.45% NaCl) **FOR ONGOING TREATMENT.**

- n Formal review by a Medical Officer after 6–8 hours is generally required. Check hydration. Check patient physically, including mental state.
- n Take note of parental observations or concerns.
- n There is some emerging evidence that 0.9% NaCl + 2.5% Glucose solution may be a more appropriate IV solution for rehydration in gastroenteritis<sup>16</sup>.
- n Rapid improvement over 2–4 hours is the 'norm'. Onset of any new symptoms (e.g. drowsiness, headache, abdominal pain) demand urgent review. Atypical behaviour of the patient should raise the question of an alternate diagnosis.
- n Repeat EUC if the child still appears unwell, if the electrolytes were markedly abnormal initially, or if the child was seriously unwell initially. Repeat EUC should also be planned for the child who continues on IV fluids.
- n Add potassium ~3 mmol/kg/24hrs when urine is passed, if initial serum potassium was normal (up to 5 mmol/kg/24hrs if marked hypokalaemia is present). Generally this is achieved by adding 10 mmol of potassium as KCl to each 500 mL bag of IV fluid.

## **Hypernatraemia** (serum sodium >149 mmol/L)

**Do not follow the Acute Gastroenteritis Guideline for fluid administration.**

**Early consultation with a Paediatrician is essential.**

Hypernatraemic dehydration is uncommon, but potentially more dangerous than when serum sodium is initially normal or slightly low. There is a greater likelihood of cerebral oedema, seizures and brain damage.

Clinically, the degree of dehydration may be underestimated.

If shock is present, resuscitate with a fluid bolus of 20 mL/kg, using 0.9% NaCl or Hartmann's solution.

Continuing rehydration should proceed slowly (usually over at least 48 hours), initially using 0.9% NaCl + 2.5% Glucose.

## **Hyponatraemia** (serum sodium <135 mmol/L)

There is some evidence that isotonic solutions, such as 0.9% NaCl (with added glucose) protect against hyponatraemia, both in the setting of Rapid IV and Standard IV Rehydration.

**We caution against the use of 0.45% NaCl + 2.5% Glucose in the setting of hyponatraemia.**

If there is severe hyponatraemia (serum sodium <130 mmol/L), ensure that the IV fluid being given is 0.9% NaCl + 2.5% Glucose, and discuss urgently with a Paediatrician.

# Investigations and Observations

- n Generally, children being enterally rehydrated do not require blood tests.
- n If nasogastric rehydration is required beyond 4 hours of rapid nasogastric rehydration, check EUC and BGL. Medical reassessment of the patient, including hydration status, is required.
- n All children with severe dehydration or with intravenous therapy, need EUC, BGL.
- n Consider blood culture and FBC if the child has a temperature  $>39^{\circ}\text{C}$ .
- n Generally urine culture is not required but urinalysis is helpful.
- n It is generally unnecessary to send stool for MC & S or viral studies. In some circumstances (e.g. bloody diarrhoea, history of travel, and community outbreak of gastroenteritis) it may be appropriate to undertake these tests.
- n Infants and children who are severely dehydrated require constant observation and monitoring, including, where possible, cardiac monitoring, pulse oximetry, frequent blood pressure measurement and urine output measurement.
- n Every child being treated in hospital for gastroenteritis, whether or not having intravenous therapy, requires observation of, and recording of, standard observations (e.g. pulse, respiration, temperature etc.) on a regular basis (not less than 4-hourly).
- n Children needing IV fluid therapy require EUC and BGL check at initial assessment. If initial EUC was markedly abnormal, or if the child's condition has not started to improve, or if the child was severely dehydrated recheck EUC at 6–8 hours. Results should be checked within two hours.
- n If there is failure to improve, deterioration or development of new signs, there should be discussion with the Admitting Medical Officer.
- n A daily lightly clothed weight can be a useful clinical parameter in the assessment of progress after admission, as well as a retrospective guide to the accuracy of the initial assessment of dehydration.

## Reintroduction of Diet

Children who are not dehydrated should continue to be fed an age-appropriate diet. Children who require rehydration should recommence age appropriate diets as soon as vomiting settles. This should be within the first 12–24 hours. Formula-fed infants should recommence full strength formula.

Refer to Gastroenteritis fact sheet jointly developed by the Children's Hospital Westmead, the Sydney Children's Hospital and the John Hunter Children's Hospital. The fact sheet is available at:

[www.chw.edu.au/parents/factsheets](http://www.chw.edu.au/parents/factsheets)

[www.sch.edu.au/health/factsheets](http://www.sch.edu.au/health/factsheets)

[www.kaleidoscope.org.au/parents/factsheets.htm](http://www.kaleidoscope.org.au/parents/factsheets.htm)

# References

1. Hunt JB, Elliott EJ, Fairclough PD et al. Water and solute absorption from hypotonic glucose-electrolyte solutions in human jejunum. *Gut*. 1992;33:479–483.
2. Booth I, Ferreira R, Desjeux JF et al. Recommendation for composition of oral rehydration solutions for the children of Europe. Report of an ESPGAN working group. *J Pediatric Gastroenterology and Nutrition*. 1992;14:113–115.
3. Ferreira RMCC, Elliott EJ, Watson AJM et al. Dominant role for osmolality in the efficacy of glucose and glycine-containing oral rehydration solutions: studies in a rat model of secretory diarrhoea. *Acta Paediatrica*. 1991; 81:46–50.
4. Hunt JB, Thillainayagam AV, Salim AFM et al. Water and solute absorption from a new hypotonic oral rehydration solution: evaluation in human and animal perfusion models. *Gut*. 1992;33:1652–1659.
5. International Study Group on Reduced-osmolality ORS solutions. Multicentre evaluation of reduced-osmolality oral rehydration salts solution. *Lancet*. 1995;345:282–285.
6. Walker-Smith JA, Sandhu BK, Isolauri E et al. Recommendations for feeding in childhood gastroenteritis. *J Pediatric Gastroenterology and Nutrition*. 1997;24:619–620.
7. Mackenzie A, Barnes G. Randomised controlled trial comparing oral and intravenous rehydration therapy in children with diarrhoea. *BMJ*. 1991;303:393–6.
8. Sharifi J, Ghavami F, Nowrouzi Z et al. Oral versus intravenous rehydration therapy in severe gastroenteritis. *Archives of Diseases in Children*. 1985; 60:856–860.
9. Issenman RM, Leung AK. Oral and intravenous rehydration of children. *Can Fam Phy*. 1993;39:2129–2136.
10. Vesikari T, Isolauri E, Baer M. A comparative trial of rapid oral and intravenous rehydration in acute diarrhoea. *Acta Paediatric Scand*. 1987;76:300–305.
11. Santosham M, Daum RS, Dillman L et al. Oral rehydration therapy of infantile diarrhoea. *New Eng J Med*. 1982;306:1070–1076.

12. Tamer AM, Friedman LB, Maxwell SRW et al. Oral rehydration of infants in a large US urban medical center. *Journal of Pediatrics*. 1985;107:14–19.
13. Listerick R, Zieserl E, Davis AT. Outpatient oral rehydration in the United States. *Am J Dis Child*. 1986;140:211–215.
14. Nager AL, Wang VJ. Comparison of nasogastric and intravenous methods of rehydration in pediatric patients with acute dehydration. *Pediatrics*. 2002;109(4):566–572.
15. Gremse DA. Effectiveness of nasogastric rehydration in hospitalised children with acute diarrhoea. *J Paediatric Gastroenterology and Nutrition*. 1995;21:145–148.
16. Neville KA et al. Isotonic is better than hypotonic saline for intravenous rehydration of children with gastroenteritis: a prospective randomised study. *Arch Dis Child* 2006 Mar; 91 (3): 226–232

# Bibliography

- Aperia A et al. Salt and water homeostasis during oral rehydration therapy. *J Paediatr* 1983;103:364–69
- Armon K, Stephenson T, MacFaul R et al. An evidence and consensus based guideline for acute diarrhoea management. *Arch Dis Child* 2001;85:132–142
- Bhargava SK et al. Oral therapy of neonates and young infants with World Health Organisation Rehydration Packets: a controlled trial of two sets of instructions *J Paediatr Gastroenterol Nutr* 1986;5:416–22
- Blum D et al. Safe oral rehydration of hypertonic dehydration. *J Paediatr Gastroenterol Nutr* 1986;5:232–35
- Borowitz SM. Are antiemetics helpful in young children suffering from acute viral gastroenteritis? *Arch Dis Child* 2005; 35:646–648
- Brill SA, Stewart TR, Brundage SI, Schreiber MA. Base deficit does not predict mortality when secondary to hyperchloremic acidosis. *Shock*.17(6):459–62, 2002.
- Bullivant EM, Wilcox CS, Welch WJ. Intrarenal vasoconstriction during hyperchloremia: role of thromboxane. *American Journal of Physiology*. 256(1 Pt 2):F152–7, 1989.
- Carey MJ, Aitken ME. Diverse effects of antiemetics in children. *NZ Med J*. 1994;107(989):452–3
- Centers for Disease Control and Prevention. Managing acute gastroenteritis among children: oral rehydration, maintenance, and nutritional therapy. *Pediatrics*. 2004;114:507
- CHOICE Study Group. Multicenter, randomized, double-blind clinical trial to evaluate the efficacy and safety of a reduced osmolarity oral rehydration salts solution in children with acute watery diarrhoea. *Pediatrics*. 2001 107(4):613–8
- Christakis DA, Wright JA, Rivara F. Promethazine therapy for gastroenteritis: towards a better understanding of use, risks and benefits...including commentary by Saur P. *Ambulatory Child Health*. 1998; 4(2):181–7
- Chubeddu LX, Trujillo LM, Talmaciu I, Gonzales V, Guariguata J, Seijas J, Miller IA, Paska W. Antiemetic activity of ondansetron in acute gastroenteritis. *Aliment Pharmacol Ther*. 1997 Feb;11(1):185–91

- Cutting WAM et al. Safety and efficacy of three oral rehydration solutions for children with diarrhoea (Edinburgh 1985–5). *Acta Paediatr Scand* 1989 Mar; 78(2):253–8.
- Duggan C, Fontaine O, Pierce NF, et al. Scientific rationale for a change in the composition of oral rehydration Solution. *JAMA*. 2004; 291:2628–31
- Durward A, Skellett S, Mayer A, Taylor D, Tibby SM, Murdoch IA. The value of the chloride: sodium ratio in differentiating the aetiology of metabolic acidosis. *Intensive Care Medicine*. 27(5):828–35, 2001.
- Durward A, Tibby SM, Skellett S, Austin C, Anderson D, Murdoch IA. The strong ion gap predicts mortality in children following cardiopulmonary bypass surgery. *Pediatric Critical Care Medicine*. 6(3):281–5, 2005.
- ESPGAN Working Group. Recommendations for composition of oral rehydration solutions for the children of Europe. *J Pediatr Gastroenterol Nutr*. 1992; (1):113–5
- Farthing MJG. Oral rehydration: an evolving solution. *J Pediatr Gastroenterol Nutr*. 2002 May–Jun;34 Suppl 1:S64–7
- Freedman SB, Adler M, Seshadri R, et al. Oral ondansetron for gastroenteritis in a paediatric emergency department. *N Engl J Med*. 2006 354(16):1698–705
- Hahn S, Kim Y, Garner P. Reduced osmolarity oral rehydration solution for treating dehydration due to diarrhoea in children: systematic review. *BMJ*. 2001; 323:81–85
- Hahn S, Kim Y, Garner P. Reduced osmolarity oral rehydration solution for treating dehydration caused by acute diarrhoea in children (Cochrane Review). *Cochrane Database Sys Rev*. 2002;(1):CD002847
- Ho AM, Karmakar MK, Contardi LH, Ng SS, Hewson JR. Excessive use of normal saline in managing traumatized patients in shock: a preventable contributor to acidosis. *Journal of Trauma–Injury Infection & Critical Care*. 51(1):173–7, 2001.
- Hunt JB, Elliott EJ, Fairclough PD, Clark ML, Farthing MJG. Water and solute absorption from hypotonic glucose–electrolyte solutions in human jejunum. *Gut*. 1992 Apr;33(4):479–83
- Isolauri E. Evaluation of an oral rehydration solution with Na 60mmol/L in infants hospitalised for acute diarrhoea or treated as outpatients. *Acta Paediatr Scand*; 74:643–49
- Kellum JA. Fluid resuscitation and hyperchloremic acidosis in experimental sepsis: improved short-term survival and acid-base balance with Hextend compared with saline. *Critical Care Medicine*. 30(2):300–5, 2002.

- King CK, Glass R, Bresee JS, Duggan C, Centers for Disease Control and Prevention. Managing acute gastroenteritis among children: oral rehydration, maintenance, and nutritional therapy. *MMWR Recomm Rep*. 2003 Nov 21;52(RR-16):1-16
- Kwon KT, Rudkin SE, Langdorf MI. Antiemetic use in pediatric gastroenteritis: a rational survey of emergency physicians, pediatricians, and pediatric emergency physicians. *Clin Pediatr*. 2002 Nov-Dec 41(9):641-52
- Leman P. Utility of ondansetron in children with vomiting. *Ann Emerg Med*. 2002 Sep;40(3):366-7
- Li ST, DiGiuseppe DL, Christakis DA. Antiemetic use for acute gastroenteritis in children. *Arch Pediatr Adolesc Med*. 2003 May;157(5):475-9
- Listernick R, Zieseri E et al. Outpatient oral rehydration in the United States. *Am J Dis Child*. 1986;140:211-215
- Mackenzie A, Barnes G. Randomised controlled trial comparing oral and intravenous rehydration therapy in children with diarrhoea. *Br Med J*. 1991;303:393-396
- Murphy MS. Guidelines for managing acute gastroenteritis based on a systematic review of published research. *Arch Dis Child*. 1998 Sep;79(3):279-84
- Murphy C, Hahn S, Volmink J. Reduced osmolarity oral rehydration solution for treating cholera. *Cochrane Database Sys Rev*. 2004, Issue 4. Art. No:CD003754
- Nager AL, Wang VJ. Comparison of nasogastric and intravenous methods of rehydration in paediatric patients with acute dehydration. *Paediatrics* 2002;109:566-72
- Nalin DR, Hirschhorn N, Greenough W 3rd, Fuchs GJ, Cash RA. Clinical concerns about reduced-osmolarity oral rehydration solution. *JAMA*. 2004 Jun 2;291(21):2632-5
- Neville KA, Verge CF, Rosenberg AR, O'Meara MW, Walker JL. Isotonic is better than hypotonic saline for intravenous rehydration of children with gastroenteritis: a prospective randomised study. *Archives of Disease in Childhood*. 91(3):226-32, 2006.
- Oral rehydration solutions for the children of Europe. Proceedings of a workshop held at XXI annual meeting of ESPGAN, Copenhagen 1988. *Acta Paediatr Scand Suppl*. 1989;364:1-50
- O'Malley CM, Frumento RJ, Hardy MA, Benvenisty AI, Brentjens TE, Mercer JS et al. A randomized, double-blind comparison of lactated Ringer's solution and 0.9% NaCl during renal transplantation. *Anesthesia & Analgesia*. 100(5):1518-24, table of contents, 2005.

- Ozuah PO, Avner JR, Stein RE. Oral rehydration, emergency physicians, and practice parameters: a national survey. *Pediatrics*. 2002 Feb;109(2):259–61
- Prough DS, Bidani A. Hyperchloremic metabolic acidosis is a predictable consequence of intraoperative infusion of 0.9% saline. *Anesthesiology*. 90(5):1247–9, 1999.
- Powell CVE, Heine RG, Priestley SJ. Randomised-controlled trial of rapid vs 24-hour nasogastric rehydration in children with acute gastroenteritis and moderate dehydration: PG3-13. *J Pediatr Gastroenterol Nutr*. 2005 May;40(5):651
- Ramsook C, Sahagun-Carreón I, et al. A randomized clinical trial comparing oral ondansetron with placebo in children with vomiting from acute gastroenteritis. *Ann Emerg Med* 2002 Apr;39(4):397–403
- Reid F, Lobo DN, Williams RN, Rowlands BJ, Allison SP. (Ab)normal saline and physiological Hartmann's solution: a randomized double-blind crossover study. *Clinical Science*. 104(1):17–24, 2003.
- Reeves JJ, Shannon MW, Fleisher GR. Ondansetron decreases vomiting associated with acute gastroenteritis: a randomized, controlled trial. *Pediatrics*. 2002 Apr;109(4):e62
- Spirko BA. Antiemetic use for gastroenteritis in children. *Ann Emerg Med*. 2003 Apr;41(4):585–6; author reply 586–7
- Loo DM, van der Graaf F, Ten WT. The effect of flavouring oral rehydration solution on its composition and palatability. *J Pediatr Gastroenterol Nutr*. 2004 ;39(5):545–8
- Saberi MS et al. Oral rehydration of diarrhoeal dehydration. Comparison of high and low sodium concentration in rehydration solutions. *Acta Paediatr Scand* 1983;72:167–70
- Santosham M et al. Oral rehydration therapy of infantile diarrhoea; a controlled study of well-nourished children hospitalised in the United States and Panama. *N Engl J Med*. 1982;306:1070–1076
- Santosham M et al. Oral rehydration therapy for acute diarrhoea in ambulatory children in the United States: a double-blind comparison of four different solutions. *Paediatrics*.1985;76:159–166
- Scheingraber S, Rehm M, Sehmisch C, Finsterer U. Rapid saline infusion produces hyperchloremic acidosis in patients undergoing gynecologic surgery. *Anesthesiology*. 90(5):1265–70, 1999.
- Skellett S, Mayer A, Durward A, Tibby SM, Murdoch IA. Chasing the base deficit: hyperchloraemic acidosis following 0.9% saline fluid resuscitation. *Archives of Disease in Childhood*. 83(6):514–6, 2000.
- Sokucu S et al. Oral rehydration therapy in infectious diarrhoea. Comparison of rehydration solutions with 60 and 90 mmol sodium per litre. *Acta Paediatr Scand* 1985;74:489–94

- Steele A, Gowrishankar M, Abrahamson S, Mazer CD, Feldman RD, Halperin ML. Postoperative hyponatremia despite near-isotonic saline infusion: a phenomenon of desalination. *Annals of Internal Medicine*. 126(1):20–5, 1997.
- Vesikari T et al. A comparative trial of rapid and intravenous rehydration in acute diarrhoea. *Acta Paediatr Scand* 1987;76:300–305
- Walker-Smith JA, Sandhu BK, Isolauri E et al. Guidelines prepared by the ESPGAN working group on acute diarrhoea: recommendations for feeding in childhood gastroenteritis. *J Pediatr Gastroenterol Nutr*. 1997 24(5):619–20
- Waters JH, Gottlieb A, Schoenwald P, Popovich MJ, Sprung J, Nelson DR. Normal saline versus lactated Ringer's solution for intraoperative fluid management in patients undergoing abdominal aortic aneurysm repair: an outcome study. *Anesthesia & Analgesia*. 93(4):817–22, 2001.
- Wilcox CS. Regulation of renal blood flow by plasma chloride. *Journal of Clinical Investigation*. 71(3):726–35, 1983.
- Wilkes NJ, Woolf R, Mutch M, Mallett SV, Peachey T, Stephens R et al. The effects of balanced versus saline-based hetastarch and crystalloid solutions on acid-base and electrolyte status and gastric mucosal perfusion in elderly surgical patients. *Anesthesia & Analgesia*. 93(4):811–6, 2001.
- Williams EL, Hildebrand KL, McCormick SA, Bedel MJ. The effect of intravenous lactated Ringer's solution versus 0.9% sodium chloride solution on serum osmolality in human volunteers. *Anesthesia & Analgesia*. 88(5):999–1003, 1999.

# Appendices

## Appendix One – Glossary

Word/ Abbreviation	Definition
Admitting Medical Officer	Most senior medical officer under whom the child is admitted to hospital
BGL	Blood Glucose (Sugar) Level
FBC	Full Blood Count
Hartmann's solution	Isotonic intravenous solution (see 'Composition' table Appendix Two)
ORS	Oral Rehydration Solution
EUC	Electrolytes, Urea and Creatinine. Ideally this should include measurement of serum sodium, potassium, chloride, bicarbonate, urea and creatinine. It is recognised that not all local laboratories offer all of these parameters 24 hours. It is essential that the serum sodium be measured on any child who is receiving intravenous rehydration therapy.

# Appendix Two – IVT Composition

	Osmolality mOsm/L	Na <sup>+</sup> mmol/L	Cl <sup>-</sup> mmol/L	Glucose g/L	K <sup>+</sup> mmol/L
0.9% NaCl	300	150	150	-	-
Hartmann's Solution	274	129	109	-	5
0.45% NaCl & 2.5% Glucose	292	76	76	25	-
0.9% NaCl & 2.5% Glucose	448	150	150	25	-

# Appendix Three – Parent Oral Rehydration Documentation Form

## Oral Fluids for your Child with Gastroenteritis

Please give your child:

1. Frequent breastfeeds if you are breastfeeding or
2. An Oral Rehydration Solution

Dilute juice (e.g. 1 part apple juice to 4 parts water) is not as effective but sometimes may be used if your child is not dehydrated.

Your child's weight is \_\_\_\_\_ kg.

Your child should drink about \_\_\_\_\_ mL every 5 minutes ( $\frac{1}{2}$  mL/kg) or 1 Hydralyte® iceblock (62.5mL) every \_\_\_\_\_ Minutes.

Use the 10 mL syringe to measure the fluid unless using Hydralyte® iceblock. Give the fluid to your child in a syringe, teaspoon, bottle or cup.

(One Hydralyte® iceblock = 62.5 mL)

Please record every time you give your child fluid and every time your child vomits, passes urine or has diarrhoea:

	TIME	FLUID TYPE	VOLUME	VOMIT	DIARRHOEA	URINE
1						
2						
3						
4						
5						
6						
7						
8						
9						
10						
11						
12						
13						
14						
15						
16						



## **ORAL FLUIDS for YOUR CHILD with GASTROENTERITIS**

The nurse who assessed your sick child has placed you into an appropriate category for urgency to see the doctor. It is most likely that your child has gastroenteritis and needs fluid treatment. Here in hospital we use oral fluids while you are waiting to see a doctor.

If you have been giving your child fluids at home, you are probably here because you feel this has been unsuccessful. The way we give oral fluids here may be slightly different and is often successful. The other side of this sheet explains exactly how much fluid and how often we want you to give it to your child.

When your child sees the doctor a decision will be made as to whether you can go home, or whether your child needs a small tube through the nose into the stomach or a drip to provide extra fluid for a few hours. Sometimes this is all it takes to make your child feel a lot better and you will then be able to go home. If this doesn't improve your child, he or she may need to be admitted to hospital for further treatment.

## Appendix Four – Parent Information

A Gastroenteritis Fact Sheet jointly developed by John Hunter Children's Hospital, Sydney Children's Hospital and Children's Hospital Westmead is available at:

[www.kaleidoscope.org.au/parents/factsheets.htm](http://www.kaleidoscope.org.au/parents/factsheets.htm)

[www.sch.edu.au/health/factsheets](http://www.sch.edu.au/health/factsheets)

[www.chw.edu.au/parents/factsheets](http://www.chw.edu.au/parents/factsheets)

*Disclaimer: The fact sheet is for educational purposes only. Please consult with your doctor or other health professional to ensure this information is right for your child.*

# Appendix Five – Resources

Fuller details may be necessary in practice, especially for the management of children with moderate or severe dehydration. Possible sources include:

NSW Health Department CIAP website, Managing Young Children and Infants with Gastroenteritis in Hospitals at: [www.ciap.health.nsw.gov.au](http://www.ciap.health.nsw.gov.au) also

The Children's Hospital Westmead Handbook, 2004 (Sections 7 — Fluid Therapy, and Section 16 — Gastroenterology), available as a book from the Children's Hospital at Westmead, or at: [www.chw.edu.au/parents/factsheets](http://www.chw.edu.au/parents/factsheets)

Gastroenteritis Fact Sheet jointly developed by the John Hunter Children's Hospital, Sydney Children's Hospital and Children's Hospital Westmead at:

[www.kaleidoscope.org.au/parents/factsheets.htm](http://www.kaleidoscope.org.au/parents/factsheets.htm)

[www.sch.edu.au/health/factsheets](http://www.sch.edu.au/health/factsheets)

[www.chw.edu.au/parents/factsheets](http://www.chw.edu.au/parents/factsheets)

## Appendix Six – Significant Changes From 2002 CPG Version

- n Levels of dehydration modified
- n Rapid and standard rehydration techniques included and volume calculations amended
- n IV Fluid Therapy section modified
- n Enteral Rehydration Therapy section included
- n Medications section revised and expanded
- n Indication for blood chemistry revised
- n Reintroduction of diet modified
- n Hyponatremia section added
- n Parent fluid documentation form included

## Appendix Seven – Alternative Calculation for Maintenance Fluids

Calculate the maintenance fluid requirement, for 24 hours, by age:

- Infants up to 9 months: 120–140mL/kg/24hrs
- Children 9–24 months: 90–100mL/kg/24hrs
- Children 2–4 years: 70–90mL/kg/24hrs
- Children 4–8 years: 60–70mL/kg/24hrs
- Older children: 50–60mL/kg/24hrs

# Appendix Eight – Working Party Members

**Dr Christopher Webber (Chair)**  
Paediatric Emergency Physician and  
Consultant Paediatrician  
Emergency Department  
Sydney Children's Hospital

**Dr Matthew Chu**  
Director of Emergency Medicine  
Canterbury Hospital

**Dr Steven Doherty (to March 2007)**  
Emergency Physician  
Emergency Department  
Tamworth Hospital

**Dr Patrick Moore**  
Staff Specialist Paediatrician  
Fairfield Hospital

**Dr Kristen Neville**  
Paediatric Endocrinologist  
Sydney Children's Hospital

**Dr Susan Phin**  
Paediatric Emergency Physician  
Emergency Department  
Children's Hospital Westmead

**Mr Phillip Way**  
Clinical Nurse Consultant  
Rural Critical Care  
Hunter New England Area Health Service

**Ms Rhonda Winskill**  
Clinical Nurse Consultant, Paediatrics  
Hunter New England Area Health Service

**Ms Leanne Crittenden**  
Coordinator  
Northern Child Health Network

**Ms Judy Lissing**  
Coordinator  
Greater Eastern and Southern  
Child Health Network

**Ms Halina Nagiello**  
Coordinator  
Western Child Health Network

**Ms Mary Crum**  
Senior Analyst  
Clinical Policy Branch (Secretariat)  
NSW Health

**Mr Bart Cavalletto**  
Manager, Statewide Paediatric Services  
NSW Health

