

Influenza Monthly Epidemiology Report, NSW

January 2012

Produced by: Population Health Division, NSW Health.

Please note influenza reports will now only be produced on a monthly basis until May 2012, unless unusual influenza activity becomes apparent prior to this time.

This report describes the surveillance for influenza and other respiratory pathogens, undertaken by NSW Health to date. This includes data from a range of surveillance systems.

For weekly communicable disease surveillance updates refer to the Communicable Disease Weekly Report at <http://www.health.nsw.gov.au/publichealth/infectious/index.asp>.

1. Summary

In January 2012:

- the rate of influenza-like illness (ILI) presentations to selected emergency departments was low and was below the normal range expected for January
- 14 cases with laboratory-confirmed influenza A – predominantly H3N2 – and seven cases of influenza B were identified by sentinel laboratories
- influenza A (H3) infection was confirmed in five month old child who died
- Rhinovirus was the most common respiratory virus identified by sentinel laboratories.

2. Emergency Department (ED) presentations


Data from 56 NSW emergency departments are included. Comparisons are made with data for the preceding six years. Recent counts are subject to change.

Presentations for influenza-like illness

- In January 2012 there were 67 presentations with influenza-like illness (rate 0.4 per 1,000 presentations) (Figure 1). This is similar to the previous month (December -70 presentations, rate 0.5 per 1,000 presentations), lower than the count of 107 (rate 0.7 per 1,000 presentations) for the month of January in 2011 and lower than the historical average for January.
- Total admissions from ED to critical care units for influenza-like illness and pneumonia were elevated in the early weeks of January but subsequently decreased, and overall were within the usual range for this time of year (Figure 2).
- Total ED presentations for bronchiolitis were within the usual range (Figure 3) for this time of year.

Figure 1: Comparison of weekly influenza-like illness presentations to NSW EDs, 2007-2012.*


Category: All visits with the above inclusions
Total 1-week counts



Note: Excludes data from 2009 to enable easier comparison of 2011 data with data from previous non-pandemic years. Includes data from 56 emergency departments. Source: NSW Health Public Health Real-time Emergency Department Surveillance System (PHREDSS) and the Centre for Epidemiology and Research, NSW Health Department.

Figure 2: Comparison of weekly admissions to hospital critical care units for ILI and pneumonia, 2007-2012..


Category: All visits with the above inclusions
Total 1-week counts



Note: As for Figure 1, although includes 2009

Figure 3: Comparison of weekly bronchiolitis presentations to NSW EDs, 2007-2012.

Category: All visits with the above inclusions
Total 1-week counts



Note: As for Figure 1, although includes 2009

3. Laboratory testing summary for influenza

In January 2012:

- 1509 tests for respiratory viruses were performed at sentinel NSW laboratories (Table 1).

- 14 specimens tested positive for influenza A – four of these have tested positive for A(H3N2), four tested positive for influenza A (pH1N1). The remainder tested negative to influenza A (pH1N1) and are assumed to be A (H3N2) (Table 1, Figure 4).
- seven cases of influenza B were reported (Table 1, Figure 4).
- the number of positive influenza tests in January was lower than the previous month and lower than that for the same month in 2011.

Laboratory testing suggests influenza has occurred at low levels. In addition, two of the influenza isolates were from people known to have acquired their infections overseas. Rhinoviruses were the most common respiratory viruses identified by laboratories.


Table 1: Summary of testing for respiratory viruses and influenza at NSW public hospital laboratories, 1 January to 27 January 2012.

Four week period ending	Virology specimens tested	Influenza A (total pos) (%)	H1N1** influenza 09 (total pos) (%)	Influenza B (total pos) (%)	Adenovirus	Parainfluenza 1, 2 & 3	RSV	Rhinovirus	Enterovirus	HMPV***
27/01/2012	1509	14 (0.9%)	4 (29%)	7 (0.5%)	37	59	37	112	64	34
Week ending										
06/01/2012	442	4 (0.9%)	0	2 (0.5%)	11	22	13	44	15	15
13/01/2012	454	4 (0.9%)	1 (25%)	1 (0.9%)	14	18	7	30	16	8
20/01/2012	349	4 (1.1%)	3 (75%)	2 (0.6%)	7	12	11	19	13	8
27/01/2012	264	2 (0.8%)	0	2 (0.8%)	5	7	6	19	20	3

** Subset of influenza A cases *** HMPV = Human metapneumovirus

Note: Data is provided by laboratories on a weekly basis. Excludes point of care tests. Influenza laboratory diagnoses using virology are reported by South Eastern Area Laboratory Services (SEALS), Institute of Clinical Pathology and Medical Research (ICPMR), The Children's Hospital at Westmead (CHW), South West Area Pathology Services (SWAPS), Pacific Laboratory Medicine Services (PaLMS), Royal Prince Alfred Hospital (RPAH), Hunter Area Pathology Service (HAPS).

Figure 4: Percent of laboratory tests positive for influenza A and influenza B, 1 January 2007 – 27 January 2012, New South Wales.



Note: Data is provided by laboratories on a weekly basis. Excludes point of care tests. Influenza laboratory diagnoses using virology are reported by South Eastern Area Laboratory Services (SEALS), Institute of Clinical Pathology and Medical Research (ICPMR), The Children's Hospital at Westmead (CHW), South West Area Pathology Services (SWAPS), Pacific Laboratory Medicine Services (PaLMS), Royal Prince Alfred Hospital (RPAH), Hunter Area Pathology Services (HAPS), St Vincents (SydPath), Nepean (no data between Oct 2010 to June 2011), Douglas Hanley Moir (DHM), VDRLab from 5 March 2010, Laverty (data from 1 April 2010 to February 2011) and St Vincent's (data since November 2010).

4. Deaths with pneumonia or influenza reported on the death certificate


Deaths registration data is routinely reviewed for deaths attributed to pneumonia or influenza. While pneumonia has many causes, a well-known indicator of seasonal and pandemic influenza activity is an increase in the number of death certificates that mention pneumonia or influenza as a cause of death.

The predicted seasonal baseline estimates the predicted rate of influenza or pneumonia deaths in the absence of influenza epidemics. If deaths exceed the epidemic threshold, then it may be an indication that influenza is beginning to circulate widely.

For the week ending 13 January:

- There were 1.0 pneumonia or influenza deaths per 100,000 NSW population, below the seasonal threshold of 1.2 per 100,000 population (Figure 5).*
- There was a significant increase in the number of deaths reported for the last week of December 2011. This was consistent with an increased number of presentations to ED's with pneumonia.

Figure 5: Rate of deaths classified as influenza and pneumonia (by NSW Registered Death Certificates) per 100,000 NSW population, 2007-2012



Source: NSW Registry of Births, Deaths and Marriages.

* Notes on interpreting death data:

- (1) The number of deaths mentioning "Pneumonia or influenza" is reported as a rate per 100,000 NSW population. Using the NSW population provides a more stable and reliable denominator than deaths from all causes. This is because pneumonia and influenza are known to contribute to increases in deaths from non-respiratory illnesses, such as deaths due to ischaemic heart disease. As the number of these deaths will increase with rises in influenza activity, the actual effect of influenza on mortality rates will be obscured if all-cause mortality is used as the denominator. This limitation is avoided by using the NSW population, which is relatively constant throughout the year, as the denominator.
- (2) Deaths referred to a coroner during the reporting period may not be available for analysis. Deaths in younger people may be more likely to require a coronial inquest. Therefore influenza-related deaths in younger people may be under-represented in these data.
- (3) The interval between death and death data availability is usually at least 7 days, and so these data are one week behind reports from emergency departments and laboratories. In addition, previous weekly rates may also change due to longer delays in reporting some deaths.

5. National and International Influenza Surveillance Links

For the latest information on national influenza activity please see the Australian Influenza Surveillance Reports at the following website:

<http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-ozflu-2011.htm>

For the latest information on international influenza activity please see the World Health Organization Influenza Updates at the following website:

<http://www.who.int/csr/disease/influenza/en/index.html>

For the information on current strains covered in this years influenza vaccine see WHO Collaborating Centre for Reference and Research on Influenza at the following website:

http://www.influenzacentre.org/centre_vaccines.htm