

# New South Wales Population Health Survey

2004

## REPORT ON ADULT HEALTH



NORTH WEST SLOPES  
DIVISION OF GENERAL PRACTICE

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice

---

**Acknowledgements:**

The development and preparation of this edition of the General Practice Division Reports was a team effort involving staff from the NSW Health Survey Program, the Centre for Epidemiology and Research, and the Alliance of NSW Divisions of General Practice. The development of these reports was part funded by the Program for Enhanced Population Health Infrastructure.

**Source:** New South Wales Population Health Survey 2002, 2003 and 2004 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

---

**Produced by:** Centre for Epidemiology and Research, Population Health Division, NSW Department of Health.

**Last updated on:** 26 September 2005

---

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Introduction

---

This report is the first of a series of triennial reports that will be produced for each division of general practice from the New South Wales Population Health Survey. The North West Slopes Division of General Practice Report aims to assist divisions in monitoring the health status of their population, and to support needs assessment, monitoring and improving quality of care in general practice, and policy planning and service development.

The NSW Department of Health in conjunction with the Alliance of NSW Divisions of General Practice has identified a number of indicators for inclusion in the reports. The indicators are:

### **Health behaviours**

Alcohol risk drinking behaviour  
Fruit intake  
Vegetable intake  
Adequate physical activity  
Smoking status  
Cancer screening  
Influenza immunisation last 12 months 65 years and over  
Pneumococcal immunisation last 5 years 65 years and over  
Homes with a smoke alarm

### **Health services**

Difficulties getting health care when needing it  
Emergency department care rating  
Hospital care rating

### **Social Capital**

Attended a community event in the last 6 months  
Most people can be trusted  
Visit neighbours

### **Health Status**


Excellent, very good or good self-rated health status  
Current asthma  
Diabetes or high blood sugar  
All natural teeth missing  
Overweight and obesity  
High and very high psychological distress

These indicators are presented in graphical and tabular form. For each indicator, the report includes a bar chart of the indicator by age group (2002, 2003 and 2004 North West Slopes data), compared with the whole of New South Wales. The tables below the charts present the indicator estimates for males, females and persons and includes the 95% confidence intervals (CI). The 95% confidence interval provides a range of values that should contain the actual value 95% of the time. In general, a wider confidence interval reflects less certainty in the indicator estimate. If confidence intervals do not overlap then the observed estimates are significantly different. **Source:** New South Wales Population Health Survey 2002, 2003 and 2004 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

---

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Risk alcohol drinking by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	38.8 (31.4-46.2)	9,100	27.6 (21.6-33.7)	6,200	33.3 (28.6-38.1)	15,300
	16-24	31.5 (12.6-50.4)	1,000	48.2 (26.5-69.9)	1,800	40.6 (25.0-56.2)	2,700
	25-34	47.4 (23.0-71.8)	1,600	38.9 (23.5-54.4)	1,400	42.9 (28.6-57.2)	3,000
	35-44	37.4 (19.4-55.3)	2,000	29.3 (16.6-42.0)	1,400	33.6 (22.6-44.6)	3,300
	45-54	55.4 (40.3-70.5)	2,600	22.9 (12.1-33.7)	800	41.8 (31.2-52.4)	3,300
	55-64	31.4 (16.7-46.2)	1,100	16.7 (7.9-25.5)	500	25.1 (15.6-34.6)	1,600
	65-74	23.1 (9.4-36.9)	600	9.8 (2.9-16.8)	200	16.6 (8.7-24.6)	800
NSW	75+	29.7 (11.6-47.8)	300	9.0 (1.3-16.8)	200	16.6 (8.1-25.1)	500
	All Ages	40.2 (39.0-41.5)	907,800	30.0 (29.0-31.0)	700,800	35.0 (34.2-35.7)	1,610,900
	16-24	49.3 (45.8-52.9)	178,300	44.8 (41.5-48.0)	157,500	46.9 (44.6-49.3)	336,500
	25-34	47.2 (43.8-50.7)	205,100	35.2 (32.7-37.7)	156,600	41.1 (38.9-43.2)	361,700
	35-44	38.0 (35.0-40.9)	169,700	32.1 (29.7-34.5)	146,500	35.0 (33.1-36.9)	317,000
	45-54	36.4 (33.6-39.1)	143,500	25.9 (23.9-28.0)	104,500	31.0 (29.3-32.7)	248,100
	55-64	36.0 (33.3-38.6)	105,500	23.4 (21.4-25.4)	66,900	29.7 (28.0-31.4)	172,600
	65-74	31.2 (28.7-33.7)	62,300	18.1 (16.1-20.1)	39,200	24.4 (22.8-26.0)	101,900
75+	34.2 (30.6-37.8)	43,400	16.6 (14.6-18.6)	29,500	23.9 (21.9-25.8)	73,100	

**Note:** Estimates are based on 683 respondents in North West Slopes and 34,980 in NSW. For this indicator 5 (0.73%) were 'not stated' (Don't know or Refused) in North West Slopes and 389 (1.11%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those people who meet Guideline 1 of the NHMRC Australian Alcohol Guidelines, as 1 or more of the following: consuming alcohol every day, consuming on average more than [4 if male/2 if female] standard drinks, consuming more than [6 if male/4 if female] on any 1 occasion or day. Questions used to define the indicator were 'How often do you usually drink alcohol?', 'On a day when you drink alcohol, how many standard drinks do you usually have?', 'In the past 4 weeks have you had more than [7-10 if male/5-6 if female] drinks in a day?', and 'In the past 4 weeks how often have you had [11 or more if male/7 or more if female] drinks in a day?'. Questions used to define the 1997 and 1998 indicator were 'How often do you have an alcoholic drink of any kind?', 'On a day when you have alcoholic drinks, how many standard drinks do you usually have?' and 'On the last occasion that you had more than [4 if male/2 if female] drinks in a day, how many drinks did you actually have?'

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Recommended fruit consumption by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	37.3 (29.7-45.0)	8,800	48.4 (42.3-54.6)	10,900	42.6 (37.7-47.6)	19,700
	16-24	38.7 (17.3-60.1)	1,200	50.7 (29.2-72.2)	1,900	45.1 (29.7-60.6)	3,100
	25-34	12.0 (0.0-24.0)	400	36.1 (20.7-51.5)	1,300	24.7 (13.9-35.6)	1,700
	35-44	45.7 (24.0-67.4)	2,400	38.8 (24.7-52.9)	1,800	42.0 (28.5-55.5)	4,200
	45-54	24.6 (12.4-36.8)	1,100	62.5 (50.0-75.0)	2,100	40.7 (30.8-50.5)	3,300
	55-64	47.6 (32.7-62.5)	1,700	44.4 (32.3-56.6)	1,300	46.2 (36.3-56.1)	3,000
	65-74	52.5 (36.3-68.7)	1,300	53.9 (40.2-67.6)	1,300	53.2 (42.6-63.8)	2,600
	75+	54.5 (33.6-75.4)	600	65.7 (51.0-80.3)	1,200	61.5 (49.3-73.6)	1,800
NSW	All Ages	40.4 (39.2-41.7)	918,800	52.6 (51.5-53.6)	1,234,800	46.6 (45.8-47.4)	2,159,200
	16-24	39.1 (35.6-42.5)	144,400	44.5 (41.3-47.8)	158,000	41.6 (39.3-44.0)	302,900
	25-34	32.9 (29.7-36.2)	142,700	44.9 (42.2-47.6)	200,200	39.0 (36.8-41.1)	343,000
	35-44	38.2 (35.3-41.2)	171,800	50.4 (47.9-53.0)	231,000	44.4 (42.4-46.4)	403,900
	45-54	41.9 (39.0-44.7)	166,600	54.7 (52.4-57.1)	222,100	48.3 (46.5-50.2)	389,200
	55-64	44.0 (41.2-46.7)	129,800	60.3 (58.0-62.6)	173,500	52.0 (50.2-53.8)	304,500
	65-74	50.0 (47.2-52.8)	99,900	63.0 (60.6-65.4)	137,100	56.8 (55.0-58.6)	237,900
	75+	50.5 (46.7-54.3)	63,600	63.4 (60.7-66.1)	112,700	58.2 (56.0-60.4)	177,900

**Note:** Estimates are based on 687 respondents in North West Slopes and 35,162 in NSW. For this indicator 1 (0.15%) were 'not stated' (Don't know or Refused) in North West Slopes and 207 (0.59%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who meet the recommended fruit consumption. The recommended consumption of fruit according to the National Campaign is 2 serves for people aged 16 years and over. One serve is equivalent to 1 medium piece or 2 small pieces of fruit. The question used to define the indicator was 'How many serves of fruit do you usually eat each day?'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Recommended vegetable consumption by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	10.7 (4.0-17.5)	2,500	15.6 (11.2-20.0)	3,500	13.1 (9.1-17.1)	6,100
	16-24	14.6 (0.0-29.4)	500	10.6 (0.0-25.6)	400	12.4 (1.9-23.0)	800
	25-34	. (-.-)	.	2.1 (0.0-6.3)	100	1.1 (0.0-3.3)	100
	35-44	19.8 (0.0-44.4)	1,000	10.7 (2.0-19.4)	500	15.5 (1.2-29.8)	1,500
	45-54	4.5 (0.0-9.1)	200	26.3 (14.6-38.0)	900	13.7 (7.5-19.9)	1,100
	55-64	15.6 (3.9-27.2)	600	20.6 (10.4-30.7)	600	17.7 (9.8-25.7)	1,100
	65-74	5.1 (0.0-10.9)	100	27.0 (12.7-41.3)	600	15.8 (7.4-24.3)	800
NSW	75+	11.3 (0.0-23.8)	100	22.4 (8.7-36.1)	400	18.3 (8.4-28.3)	600
	All Ages	7.0 (6.5-7.6)	159,600	13.6 (12.9-14.3)	320,100	10.4 (9.9-10.9)	479,700
	16-24	5.4 (4.0-6.8)	19,800	8.0 (6.2-9.7)	28,300	6.7 (5.5-7.8)	48,100
	25-34	5.2 (3.8-6.7)	22,700	9.2 (7.5-10.8)	40,900	7.2 (6.1-8.3)	63,600
	35-44	5.2 (4.0-6.5)	23,600	11.6 (10.0-13.3)	53,300	8.5 (7.4-9.5)	76,900
	45-54	7.1 (5.7-8.5)	28,200	17.8 (16.0-19.6)	72,200	12.5 (11.3-13.7)	100,400
	55-64	9.2 (7.7-10.8)	27,200	19.3 (17.5-21.1)	55,400	14.2 (13.0-15.4)	82,600
	65-74	11.9 (10.1-13.7)	23,700	20.4 (18.4-22.5)	44,300	16.3 (15.0-17.7)	68,000
75+	11.5 (9.2-13.9)	14,400	14.4 (12.4-16.4)	25,700	13.2 (11.7-14.8)	40,100	

**Note:** Estimates are based on 687 respondents in North West Slopes and 35,000 in NSW. For this indicator 1 (0.15%) were 'not stated' (Don't know or Refused) in North West Slopes and 369 (1.05%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who meet the recommended consumption of vegetables. The recommended vegetable intake according to the National Campaign is 5 serves per day for persons aged 16 years and over. One serve is equivalent to 1/2 cup of cooked vegetables or 1 cup of salad vegetables. The question used to define the indicator was: 'How many serves of vegetables do you usually eat each day?'

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Adequate physical activity by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	51.9 (44.1-59.7)	12,300	41.1 (34.9-47.3)	9,300	46.5 (41.5-51.5)	21,500
	16-24	71.1 (52.0-90.2)	2,200	56.8 (35.6-77.9)	2,100	63.4 (48.8-78.0)	4,300
	25-34	70.6 (51.3-90.0)	2,300	45.0 (29.0-61.0)	1,700	57.1 (43.4-70.8)	4,000
	35-44	39.7 (20.3-59.1)	2,100	37.6 (23.5-51.7)	1,700	38.3 (26.2-50.4)	3,800
	45-54	48.5 (33.0-63.9)	2,300	43.2 (30.5-55.8)	1,500	46.2 (35.8-56.7)	3,700
	55-64	46.3 (31.4-61.2)	1,700	32.6 (21.4-43.7)	900	40.3 (30.6-50.0)	2,600
	65-74	42.2 (26.2-58.2)	1,100	46.0 (32.4-59.7)	1,100	44.1 (33.5-54.7)	2,200
	75+	53.7 (32.8-74.7)	600	15.2 (4.9-25.6)	300	29.3 (17.9-40.8)	900
NSW	All Ages	52.5 (51.2-53.8)	1,199,100	44.0 (43.0-45.1)	1,037,600	48.1 (47.3-49.0)	2,241,000
	16-24	66.8 (63.4-70.1)	247,000	57.5 (54.4-60.7)	205,200	62.2 (59.9-64.5)	453,800
	25-34	53.7 (50.3-57.1)	234,900	46.5 (43.9-49.2)	208,200	50.0 (47.8-52.2)	443,200
	35-44	49.4 (46.4-52.5)	223,500	43.6 (41.1-46.2)	200,700	46.5 (44.5-48.5)	425,200
	45-54	49.5 (46.6-52.4)	197,600	43.2 (40.9-45.5)	175,600	46.3 (44.4-48.2)	374,200
	55-64	49.8 (47.0-52.5)	147,500	41.6 (39.3-43.9)	120,000	45.6 (43.8-47.4)	267,500
	65-74	49.6 (46.9-52.4)	99,800	37.3 (34.9-39.7)	81,500	43.2 (41.3-45.0)	181,800
	75+	38.1 (34.4-41.8)	48,600	26.0 (23.6-28.3)	46,500	30.8 (28.8-32.9)	95,200

**Note:** Estimates are based on 688 respondents in North West Slopes and 35,365 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 4 (0.01%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who did adequate physical activity. Adequate physical activity is a total of 150 minutes per week on 5 separate occasions. The total minutes were calculated by adding minutes in the last week spent walking continuously for at least 10 minutes, minutes doing moderate physical activity, plus 2 x minutes doing vigorous physical activity. The questions used to define the indicator were 'In the last week, how many times have you walked continuously for at least 10 minutes for recreation or exercise or to get to or from places?', 'What do you estimate was the total time you spent walking in this way in the last week?', 'In the last week, how many times did you do any vigorous physical activity which made you breathe harder or puff and pant?', 'What do you estimate was the total time you spent doing this vigorous physical activity in the last week?', 'In the last week, how many times did you do any other more moderate physical activity that you have not already mentioned?' and 'What do you estimate was the total time that you spent doing these activities in the last week?'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Current daily or occasional smoking by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	23.6 (17.2-30.0)	5,600	25.7 (19.7-31.6)	5,800	24.5 (20.1-28.9)	11,400
	16-24	16.6 (3.9-29.2)	500	35.6 (13.6-57.6)	1,300	26.8 (12.7-40.9)	1,800
	25-34	38.5 (15.1-61.9)	1,300	35.7 (19.7-51.7)	1,300	37.0 (23.1-50.9)	2,600
	35-44	30.8 (13.0-48.5)	1,600	32.4 (18.8-46.0)	1,500	31.2 (19.9-42.5)	3,100
	45-54	17.4 (7.2-27.5)	800	27.7 (16.0-39.4)	900	21.7 (13.9-29.6)	1,800
	55-64	25.0 (11.1-38.8)	900	20.0 (10.1-29.8)	600	22.8 (13.8-31.7)	1,500
	65-74	8.2 (0.0-19.1)	200	2.7 (0.0-5.8)	100	5.5 (0.0-11.4)	300
NSW	75+	20.6 (2.4-38.9)	200	5.0 (0.0-12.2)	100	10.7 (2.3-19.2)	300
	All Ages	23.9 (22.8-25.0)	546,100	19.4 (18.6-20.2)	456,900	21.7 (21.0-22.3)	1,007,600
	16-24	26.7 (23.7-29.8)	98,800	26.0 (23.2-28.8)	92,600	26.6 (24.5-28.6)	193,700
	25-34	32.9 (29.8-36.1)	144,000	25.8 (23.4-28.1)	115,200	29.3 (27.3-31.2)	259,500
	35-44	29.1 (26.5-31.8)	131,400	22.4 (20.5-24.4)	103,100	25.8 (24.1-27.4)	235,300
	45-54	24.3 (21.9-26.8)	97,300	19.6 (17.8-21.4)	79,700	22.0 (20.4-23.5)	177,400
	55-64	15.7 (13.8-17.6)	46,600	14.6 (13.1-16.2)	42,200	15.2 (14.0-16.4)	89,200
	65-74	10.3 (8.7-11.9)	20,700	7.9 (6.6-9.2)	17,200	9.1 (8.1-10.1)	38,300
75+	5.7 (4.0-7.4)	7,300	3.9 (3.0-4.8)	6,900	4.6 (3.7-5.5)	14,200	

**Note:** Estimates are based on 688 respondents in North West Slopes and 35,353 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 16 (0.05%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who smoked daily or occasionally. The question used to define the indicator was 'Which of the following best describes your smoking status? Smoke daily, Smoke occasionally, Do not smoke now, but I used to, I have tried it a few times but never smoked regularly, and I have never smoked'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Smoke-free households by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	83.3 (77.8-88.9)	19,700	84.8 (80.5-89.1)	19,100	84.1 (80.6-87.6)	38,900
	16-24	86.6 (75.5-97.7)	2,700	100 (100-100)	3,700	93.8 (88.7-98.9)	6,400
	25-34	81.1 (63.6-98.7)	2,700	83.7 (71.8-95.6)	3,100	82.5 (72.1-92.9)	5,800
	35-44	77.2 (61.2-93.1)	4,100	72.2 (59.3-85.1)	3,300	75.1 (64.7-85.5)	7,500
	45-54	77.0 (63.2-90.9)	3,600	82.4 (72.7-92.0)	2,800	79.3 (70.3-88.3)	6,400
	55-64	92.1 (85.8-98.4)	3,400	86.5 (78.4-94.7)	2,400	89.7 (84.6-94.8)	5,700
	65-74	92.4 (84.0-100)	2,300	85.3 (71.4-99.2)	2,000	89.0 (80.7-97.2)	4,300
NSW	75+	87.2 (72.9-100)	1,000	89.8 (79.4-100)	1,800	88.9 (80.4-97.3)	2,700
	All Ages	81.7 (80.7-82.6)	1,863,400	83.3 (82.6-84.1)	1,962,200	82.5 (81.9-83.1)	3,837,200
	16-24	76.8 (73.9-79.7)	283,100	79.3 (76.7-82.0)	282,800	78.1 (76.1-80.0)	568,700
	25-34	81.4 (79.0-83.9)	356,100	84.6 (82.7-86.5)	378,600	83.0 (81.5-84.6)	735,800
	35-44	83.4 (81.4-85.5)	376,500	84.1 (82.4-85.9)	386,500	83.8 (82.4-85.1)	764,800
	45-54	79.5 (77.4-81.7)	317,900	81.8 (80.1-83.5)	332,300	80.7 (79.3-82.0)	651,600
	55-64	82.0 (80.0-84.1)	243,200	82.0 (80.3-83.8)	236,300	82.0 (80.6-83.4)	481,300
	65-74	86.0 (84.1-87.9)	172,900	84.4 (82.5-86.4)	184,300	85.1 (83.8-86.5)	358,400
75+	89.1 (86.8-91.5)	113,700	90.3 (88.8-91.8)	161,400	89.7 (88.4-91.0)	276,500	

**Note:** Estimates are based on 686 respondents in North West Slopes and 35,328 in NSW. For this indicator 2 (0.29%) were 'not stated' (Don't know or Refused) in North West Slopes and 39 (0.11%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those households who indicated that their home was smoke-free. The question used to define the indicator was 'Which of the following best describes your home situation? My home is smoke-free, People occasionally smoke in the house, and People frequently smoke in the house'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Homes with a smoke alarm or detector by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	75.7 (69.3-82.0)	17,600	78.0 (73.3-82.7)	17,300	76.6 (72.6-80.6)	34,900
	16-24	74.3 (55.3-93.3)	2,300	82.3 (69.3-95.4)	2,800	78.5 (66.9-90.0)	5,200
	25-34	69.8 (46.5-93.0)	2,300	72.3 (58.4-86.2)	2,700	71.1 (57.9-84.3)	5,000
	35-44	84.0 (72.6-95.5)	4,300	83.1 (72.8-93.4)	3,800	82.6 (74.7-90.6)	8,100
	45-54	82.2 (71.9-92.4)	3,700	78.7 (68.3-89.1)	2,600	80.7 (73.3-88.1)	6,400
	55-64	60.1 (44.5-75.7)	2,100	69.7 (57.9-81.6)	1,900	64.3 (53.9-74.7)	4,100
	65-74	77.9 (64.7-91.1)	1,900	78.9 (68.5-89.3)	1,900	78.4 (70.0-86.8)	3,800
	75+	76.9 (59.6-94.2)	900	78.7 (66.3-91.1)	1,500	78.0 (67.9-88.1)	2,400
NSW	All Ages	71.8 (70.7-72.9)	1,599,000	72.6 (71.7-73.6)	1,676,900	72.2 (71.5-72.9)	3,286,100
	16-24	72.1 (68.9-75.2)	258,400	69.9 (67.0-72.9)	244,300	71.1 (68.9-73.2)	505,500
	25-34	71.4 (68.3-74.4)	302,700	70.1 (67.6-72.6)	307,300	70.7 (68.7-72.7)	610,700
	35-44	74.1 (71.5-76.8)	326,200	78.0 (75.9-80.1)	350,100	76.1 (74.4-77.8)	678,200
	45-54	72.8 (70.2-75.5)	284,400	71.5 (69.3-73.7)	284,500	72.1 (70.4-73.8)	569,900
	55-64	68.7 (66.1-71.3)	200,100	70.9 (68.8-73.1)	201,600	69.8 (68.2-71.5)	403,500
	65-74	71.1 (68.5-73.7)	140,200	75.6 (73.4-77.7)	162,900	73.3 (71.6-75.0)	304,000
	75+	69.5 (65.9-73.0)	87,000	72.0 (69.5-74.5)	126,200	70.8 (68.8-72.9)	214,300

**Note:** Estimates are based on 680 respondents in North West Slopes and 34,762 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 0 (0%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who have a smoke alarm or detector in their home. The question used to define the indicator was 'Do you have any of the following fire safety measures in your home? Fire alarm (hard wired), Fire alarm (battery operated only), Fire sprinkler system, Safety switch or circuit breaker, Fire extinguisher, Fire evacuation plan, External water supply, External sprinkler'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Screening mammogram within the last 2 years by age




Division	Age (years)	Females (95% CI)	Females (est. no.)
North West Slopes	All Ages	85.9 (75.5-96.3)	1,500
	50-54	86.4 (66.9-100)	500
	55-59	76.5 (54.6-98.3)	400
	60-64	100 (100-100)	300
	65-69	86.7 (68.3-100)	200
NSW	All Ages	75.6 (73.0-78.2)	130,300
	50-54	70.6 (65.5-75.7)	39,300
	55-59	77.3 (72.6-82.1)	34,400
	60-64	82.5 (78.5-86.6)	32,200
	65-69	73.4 (66.6-80.2)	24,500

**Note:** Estimates are based on 49 respondents in North West Slopes and 2,202 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 5 (0.23%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who have had a screening mammogram in the last 2 years. This excludes women who had a mammogram for a breast problem or had had breast cancer in the past. The questions used to define the indicator were "Have you ever had a mammogram?" and "When did you last have a mammogram?" and "Can you tell me all the reasons why you had your last mammogram?".

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Pap test within the last 2 years by age




Division	Age (years)	Females (95% CI)	Females (est. no.)
North West Slopes	All Ages	82.7 (73.5-91.9)	4,500
	20-29	93.4 (83.0-100)	1,300
	30-39	79.6 (59.8-99.4)	1,300
	40-49	74.0 (51.8-96.3)	900
	50-59	84.5 (71.2-97.8)	700
	60-69	79.9 (55.7-100)	300
NSW	All Ages	74.6 (72.8-76.4)	435,300
	20-29	65.1 (60.7-69.6)	95,100
	30-39	78.1 (74.6-81.5)	123,500
	40-49	81.1 (77.9-84.4)	111,000
	50-59	77.8 (74.2-81.4)	71,300
	60-69	67.4 (63.0-71.9)	34,400

**Note:** Estimates are based on 86 respondents in North West Slopes and 4,499 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 23 (0.51%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who have had a Pap test in the last 2 years and have not had a hysterectomy. The questions used to define the indicator were "A Pap test, is a routine test carried out by a doctor. It is recommended for all women for early detection of cancer of the cervix. Have you ever had a Pap test?", "When did you last have a Pap test?" and "Have you ever had a hysterectomy?".

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Vaccinated against influenza in the last 12 months by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	83.9 (73.5-94.3)	3,000	77.6 (69.3-86.0)	3,400	80.5 (73.9-87.0)	6,400
	65-69	78.6 (60.4-96.8)	1,100	59.6 (43.7-75.6)	800	69.7 (57.2-82.3)	1,900
	70-75	89.1 (69.2-100)	900	91.0 (81.7-100)	1,000	90.1 (79.3-100)	2,000
	75+	85.9 (71.0-100)	1,000	81.6 (68.3-94.8)	1,600	83.2 (73.1-93.2)	2,600
NSW	All Ages	75.5 (73.5-77.4)	247,700	76.3 (74.8-77.8)	302,900	75.9 (74.7-77.1)	553,300
	65-69	65.2 (61.6-68.7)	69,900	68.8 (65.7-72.0)	77,000	66.9 (64.5-69.3)	147,100
	70-75	77.2 (73.8-80.6)	72,400	77.3 (74.4-80.2)	82,200	77.2 (75.0-79.4)	155,300
	75+	82.9 (79.9-85.8)	105,500	80.4 (78.3-82.6)	143,700	81.5 (79.7-83.2)	250,900


**Note:** Estimates are based on 196 respondents in North West Slopes and 9,684 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 12 (0.12%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those respondents that are aged 65 years and over who have been vaccinated against flu in the past 12 months.

The question used was 'Were you vaccinated or immunised against flu in the past 12 months?'

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Vaccinated against pneumococcal disease in the last 5 years by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	43.4 (30.2-56.6)	1,500	47.2 (36.9-57.5)	2,000	45.5 (37.3-53.7)	3,500
	65-69	33.3 (12.2-54.4)	500	25.9 (12.2-39.7)	300	29.8 (16.9-42.7)	800
	70-75	56.4 (31.8-81.1)	500	47.5 (24.7-70.3)	500	51.7 (34.6-68.9)	1,000
	75+	40.1 (16.7-63.6)	300	58.6 (42.9-74.3)	1,100	52.9 (39.4-66.4)	1,400
NSW	All Ages	42.3 (40.0-44.5)	134,100	47.1 (45.2-48.9)	182,800	44.9 (43.5-46.3)	318,300
	65-69	27.5 (24.1-30.9)	28,400	32.2 (28.8-35.5)	35,200	29.9 (27.5-32.3)	63,700
	70-75	42.0 (37.8-46.1)	38,000	49.2 (45.8-52.7)	51,500	45.8 (43.1-48.5)	89,700
	75+	56.7 (52.6-60.7)	60,200	54.8 (51.9-57.7)	84,800	55.6 (53.2-57.9)	146,000

**Note:** Estimates are based on 190 respondents in North West Slopes and 9,430 in NSW. For this indicator 6 (3.16%) were 'not stated' (Don't know or Refused) in North West Slopes and 266 (2.82%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those respondents that are aged 65 years and over who have been vaccinated against pneumococcal disease in the last 5 years. The question used was "When were you last vaccinated or immunised against pneumonia?".

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Excellent, very good, or good self-rated health status by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	84.1 (79.5-88.8)	19,900	79.2 (74.3-84.1)	17,900	81.8 (78.4-85.1)	37,900
	16-24	86.7 (72.4-100)	2,700	86.2 (73.8-98.7)	3,200	86.5 (77.0-95.9)	5,900
	25-34	96.8 (90.6-100)	3,200	78.1 (63.5-92.7)	2,900	86.9 (78.0-95.9)	6,100
	35-44	90.5 (81.4-99.7)	4,800	72.8 (59.5-86.1)	3,400	82.4 (74.1-90.8)	8,300
	45-54	87.3 (78.4-96.3)	4,100	80.2 (69.9-90.6)	2,700	84.3 (77.5-91.2)	6,800
	55-64	74.5 (62.4-86.6)	2,700	81.9 (73.0-90.8)	2,300	77.7 (69.9-85.6)	5,000
	65-74	65.5 (49.6-81.4)	1,600	78.0 (68.1-87.9)	1,900	71.6 (61.9-81.3)	3,500
	75+	68.7 (49.2-88.2)	800	78.8 (67.1-90.5)	1,500	75.1 (64.7-85.6)	2,300
NSW	All Ages	81.2 (80.2-82.2)	1,852,000	79.8 (79.0-80.6)	1,878,300	80.5 (79.8-81.1)	3,741,600
	16-24	85.1 (82.4-87.8)	314,200	83.2 (80.8-85.5)	296,600	84.2 (82.4-86.0)	613,600
	25-34	84.6 (82.1-87.2)	370,200	85.4 (83.5-87.2)	381,700	85.0 (83.4-86.6)	753,200
	35-44	83.2 (81.1-85.3)	376,100	82.0 (80.1-83.8)	376,800	82.6 (81.2-84.0)	754,800
	45-54	81.6 (79.4-83.7)	325,300	80.0 (78.2-81.8)	324,900	80.7 (79.3-82.1)	651,700
	55-64	76.1 (73.7-78.5)	225,600	75.5 (73.5-77.5)	217,500	75.7 (74.1-77.3)	444,000
	65-74	77.1 (74.7-79.4)	154,600	73.2 (71.0-75.5)	159,500	75.0 (73.4-76.7)	315,300
	75+	68.0 (64.5-71.5)	86,100	68.0 (65.5-70.6)	121,400	68.1 (66.0-70.2)	209,000

**Note:** Estimates are based on 688 respondents in North West Slopes and 35,324 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 45 (0.13%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those responding excellent, very good, or good to a global self-rated health status question. The question used to define the indicator was 'Overall, how would you rate your health during the past 4 weeks? Was it excellent, very good, good, fair, poor, or very poor'. Question used to define the indicator in 1997 and 1998 was 'In general would you say your health is excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Current asthma by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	9.6 (5.7-13.5)	2,300	17.5 (12.8-22.2)	4,000	13.4 (10.3-16.5)	6,200
	16-24	22.8 (5.2-40.4)	700	31.6 (13.0-50.3)	1,200	27.5 (14.6-40.4)	1,900
	25-34	4.9 (0.0-10.9)	200	13.8 (3.1-24.4)	500	9.6 (3.1-16.1)	700
	35-44	10.9 (1.3-20.4)	600	22.5 (10.2-34.8)	1,000	16.1 (8.2-24.0)	1,600
	45-54	5.8 (0.0-12.0)	300	11.2 (3.7-18.8)	400	8.1 (3.2-12.9)	600
	55-64	11.1 (2.3-19.9)	400	14.0 (6.4-21.5)	400	12.4 (6.4-18.3)	800
	65-74	3.4 (0.0-8.7)	100	12.9 (5.2-20.6)	300	8.1 (3.4-12.8)	400
	75+	4.4 (0.0-12.8)	0	8.2 (1.2-15.3)	200	6.8 (1.4-12.3)	200
NSW	All Ages	9.1 (8.4-9.9)	208,300	12.3 (11.6-12.9)	288,700	10.7 (10.2-11.2)	498,500
	16-24	11.8 (9.4-14.2)	43,600	15.0 (12.9-17.1)	53,400	13.3 (11.7-14.9)	97,100
	25-34	12.0 (9.5-14.5)	52,400	10.7 (9.1-12.3)	47,900	11.3 (9.9-12.8)	100,300
	35-44	6.8 (5.4-8.2)	30,600	10.3 (8.9-11.7)	47,300	8.6 (7.6-9.6)	78,300
	45-54	8.1 (6.5-9.6)	32,300	13.8 (12.2-15.4)	56,000	11.0 (9.8-12.1)	88,500
	55-64	7.4 (6.0-8.8)	22,000	12.9 (11.4-14.4)	37,100	10.2 (9.1-11.2)	59,700
	65-74	8.3 (6.8-9.7)	16,600	13.0 (11.3-14.6)	28,300	10.7 (9.6-11.8)	45,100
	75+	8.4 (6.4-10.4)	10,800	10.5 (8.7-12.2)	18,700	9.6 (8.3-10.9)	29,600

**Note:** Estimates are based on 686 respondents in North West Slopes and 35,333 in NSW. For this indicator 2 (0.29%) were 'not stated' (Don't know or Refused) in North West Slopes and 36 (0.1%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those respondents who had symptoms of asthma or had taken treatment for asthma in the last 12 months. The questions used to define the indicator were 'Have you ever been told by a doctor or at a hospital that you have asthma?' and 'Have you had symptoms of asthma or taken treatment for asthma in the last 12 months?'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Diabetes or high blood sugar by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	8.3 (4.5-12.2)	2,000	5.9 (3.1-8.6)	1,300	7.1 (4.8-9.5)	3,300
	16-24	5.5 (0.0-15.9)	200	. (-.-)	.	2.5 (0.0-7.4)	200
	25-34	. (-.-)	.	7.9 (0.0-18.6)	300	4.2 (0.0-10.0)	300
	35-44	0.7 (0.0-2.2)	0	4.8 (0.0-11.2)	200	2.6 (0.0-5.7)	300
	45-54	13.0 (0.0-26.3)	600	3.2 (0.0-7.3)	100	8.9 (0.8-16.9)	700
	55-64	14.1 (5.0-23.2)	500	9.1 (1.8-16.4)	300	11.9 (5.9-18.0)	800
	65-74	19.9 (6.8-33.0)	500	9.2 (2.8-15.5)	200	14.6 (7.1-22.1)	700
	75+	13.5 (0.4-26.6)	200	11.7 (1.7-21.7)	200	12.4 (4.4-20.3)	400
NSW	All Ages	7.2 (6.6-7.7)	163,400	5.5 (5.1-5.9)	128,900	6.3 (6.0-6.7)	294,200
	16-24	0.9 (0.3-1.5)	3,300	1.8 (0.9-2.7)	6,400	1.4 (0.8-1.9)	9,900
	25-34	2.7 (1.5-3.8)	11,600	1.8 (1.2-2.5)	8,100	2.2 (1.6-2.9)	19,700
	35-44	3.5 (2.5-4.6)	16,000	2.6 (1.9-3.3)	12,000	3.1 (2.5-3.7)	28,000
	45-54	8.0 (6.6-9.5)	32,000	5.3 (4.3-6.3)	21,400	6.7 (5.8-7.6)	54,100
	55-64	16.0 (13.9-18.0)	47,200	9.9 (8.5-11.3)	28,500	13.0 (11.7-14.3)	76,200
	65-74	17.6 (15.4-19.7)	35,200	14.2 (12.4-16.0)	30,800	15.8 (14.4-17.2)	66,300
	75+	14.3 (11.9-16.8)	18,200	12.1 (10.3-13.9)	21,700	13.0 (11.6-14.5)	40,100

**Note:** Estimates are based on 686 respondents in North West Slopes and 35,248 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 8 (0.02%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who either had diabetes or high blood sugar levels but did not have gestational diabetes. The questions used to define the indicator were 'Have you ever been told by a doctor or at a hospital that you have diabetes?', 'Have you ever been told by a doctor or at a hospital that you have high sugar levels in your blood or urine?' and if female 'Were you pregnant when you were first told that you had diabetes or high blood sugar?' and 'Have you ever had diabetes or high blood sugar apart from when you were pregnant?'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice All natural teeth missing by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	5.1 (2.6-7.5)	1,200	9.3 (6.6-12.0)	2,100	7.1 (5.3-8.9)	3,300
	16-24	. (-.)	.	. (-.)	.	. (-.)	.
	25-34	. (-.)	.	. (-.)	.	. (-.)	.
	35-44	. (-.)	.	4.0 (0.0-9.4)	200	1.8 (0.0-4.4)	200
	45-54	1.7 (0.0-4.9)	100	3.3 (0.0-7.4)	100	2.4 (0.0-4.9)	200
	55-64	9.2 (0.6-17.8)	300	14.3 (6.8-21.8)	400	11.4 (5.6-17.3)	700
	65-74	16.9 (3.9-29.9)	400	27.8 (16.6-38.9)	700	22.2 (13.7-30.8)	1,100
	75+	32.1 (12.3-51.9)	400	37.3 (21.9-52.6)	700	35.4 (23.2-47.5)	1,100
NSW	All Ages	4.5 (4.1-4.9)	103,700	7.8 (7.3-8.2)	182,900	6.2 (5.9-6.5)	288,800
	16-24	0.3 (0.0-0.7)	1,200	1.0 (0.1-1.9)	3,600	0.7 (0.2-1.1)	4,800
	25-34	0.2 (0.0-0.5)	1,100	0.4 (0.0-0.9)	2,000	0.3 (0.1-0.6)	3,100
	35-44	0.6 (0.2-1.1)	2,900	1.3 (0.7-1.8)	5,800	1.0 (0.6-1.3)	8,700
	45-54	3.5 (2.5-4.5)	14,100	3.9 (3.0-4.8)	15,900	3.7 (3.0-4.4)	30,000
	55-64	6.8 (5.5-8.1)	20,200	11.8 (10.4-13.2)	33,900	9.4 (8.4-10.4)	55,100
	65-74	15.0 (13.1-16.9)	30,200	25.3 (23.1-27.5)	55,300	20.4 (18.9-21.9)	85,900
	75+	26.7 (23.3-30.2)	34,100	37.1 (34.5-39.7)	66,400	32.8 (30.7-34.8)	101,100

**Note:** Estimates are based on 688 respondents in North West Slopes and 35,355 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 14 (0.04%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those respondents who had all of their natural teeth missing. The question used to define the indicator was 'Are any of your natural teeth missing? (Natural teeth does not include dentures but includes wisdom teeth)'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Overweight and obesity by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	61.6 (54.2-69.1)	14,200	47.7 (41.3-54.0)	10,200	55.0 (50.0-60.1)	24,500
	16-24	20.3 (4.9-35.7)	600	26.3 (5.7-46.9)	900	23.5 (10.2-36.7)	1,600
	25-34	75.5 (56.2-94.7)	2,300	47.7 (31.3-64.2)	1,700	60.3 (46.9-73.8)	4,000
	35-44	69.9 (52.2-87.6)	3,700	54.9 (39.6-70.3)	2,300	63.7 (51.4-76.0)	6,100
	45-54	79.2 (66.6-91.7)	3,500	51.7 (38.6-64.7)	1,700	67.4 (57.8-77.0)	5,300
	55-64	60.2 (45.0-75.3)	2,200	60.7 (48.0-73.3)	1,600	60.4 (50.2-70.6)	3,800
	65-74	61.0 (45.2-76.8)	1,500	45.9 (31.7-60.1)	1,000	53.9 (43.2-64.6)	2,500
	75+	37.4 (17.0-57.8)	400	48.2 (32.2-64.3)	900	44.1 (31.3-57.0)	1,300
NSW	All Ages	54.5 (53.3-55.8)	1,223,800	40.0 (39.0-41.0)	891,000	47.3 (46.5-48.1)	2,121,200
	16-24	30.3 (26.9-33.6)	108,600	18.3 (15.9-20.7)	61,200	24.6 (22.5-26.8)	171,400
	25-34	50.6 (47.2-54.1)	217,500	33.4 (30.9-36.0)	140,400	42.2 (40.0-44.3)	359,000
	35-44	61.1 (58.1-64.1)	272,300	39.0 (36.5-41.6)	170,100	50.1 (48.1-52.2)	442,900
	45-54	64.5 (61.6-67.3)	254,200	48.6 (46.2-51.0)	188,600	56.6 (54.7-58.5)	443,400
	55-64	66.2 (63.5-68.9)	193,300	53.0 (50.6-55.3)	146,800	59.8 (58.0-61.6)	341,500
	65-74	61.1 (58.4-63.8)	121,000	55.3 (52.8-57.8)	114,400	58.1 (56.2-59.9)	236,200
	75+	45.5 (41.7-49.2)	56,900	42.2 (39.3-45.0)	69,500	43.5 (41.2-45.8)	126,800

**Note:** Estimates are based on 659 respondents in North West Slopes and 33,809 in NSW. For this indicator 29 (4.4%) were 'not stated' (Don't know or Refused) in North West Slopes and 1557 (4.61%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those with a Body Mass Index (BMI) of 25 or higher. The questions used to define the indicator were 'How tall are you without shoes?' and 'How much do you weigh without clothes or shoes?'. The BMI is calculated as follows: BMI = weight(kg)/height<sup>2</sup>(m). Categories for this indicator include overweight (BMI between 25 and 29.9) and obese (BMI of 30 and over).

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice High and very high psychological distress by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	9.4 (4.6-14.2)	2,200	11.5 (7.6-15.5)	2,600	10.4 (7.3-13.6)	4,800
	16-24	3.6 (0.0-10.7)	100	8.8 (0.0-18.2)	300	6.4 (0.4-12.4)	400
	25-34	24.7 (1.9-47.5)	800	17.6 (3.9-31.4)	700	21.0 (7.9-34.0)	1,500
	35-44	9.8 (0.0-20.8)	500	18.5 (7.2-29.9)	900	13.7 (5.8-21.7)	1,400
	45-54	4.9 (0.2-9.6)	200	6.3 (0.7-11.8)	200	5.5 (1.9-9.1)	400
	55-64	9.2 (1.9-16.5)	300	7.3 (1.2-13.4)	200	8.4 (3.5-13.3)	500
	65-74	7.9 (0.0-18.5)	200	8.1 (2.0-14.2)	200	8.0 (1.8-14.1)	400
	75+	1.4 (0.0-4.1)	0	8.1 (0.0-16.8)	200	5.5 (0.0-11.1)	200
NSW	All Ages	10.4 (9.7-11.2)	235,900	13.9 (13.2-14.7)	324,900	12.2 (11.7-12.7)	562,000
	16-24	11.8 (9.5-14.0)	43,500	19.6 (17.0-22.2)	69,700	15.6 (13.9-17.3)	113,600
	25-34	12.2 (10.0-14.4)	52,600	13.9 (12.1-15.7)	61,800	13.0 (11.6-14.5)	114,500
	35-44	10.8 (9.1-12.6)	48,200	15.0 (13.3-16.8)	68,600	12.9 (11.7-14.1)	116,800
	45-54	10.7 (9.1-12.3)	42,300	14.2 (12.6-15.8)	57,000	12.5 (11.3-13.6)	99,700
	55-64	9.7 (7.9-11.4)	28,400	11.8 (10.2-13.4)	33,700	10.7 (9.5-11.8)	62,100
	65-74	6.1 (4.8-7.5)	12,200	8.4 (7.1-9.6)	18,100	7.3 (6.4-8.2)	30,500
	75+	7.0 (5.0-9.0)	8,700	9.2 (7.6-10.9)	16,100	8.2 (7.0-9.5)	24,800

**Note:** Estimates are based on 684 respondents in North West Slopes and 34,985 in NSW. For this indicator 4 (0.58%) were 'not stated' (Don't know or Refused) in North West Slopes and 384 (1.1%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those with a Kessler 10 (K10) score of 22 or above. The K10 is a 10-item questionnaire about the level of anxiety and depressive symptoms in the most recent 4-week period. K10 scores for respondents aged 65 years and over were derived using 6 questions from the K10 questionnaire.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Difficulties getting health care when needing it by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	24.0 (17.7-30.3)	5,400	30.3 (24.7-35.9)	6,600	27.1 (22.9-31.4)	12,100
	16-24	23.3 (2.9-43.8)	700	3.5 (0.0-8.4)	100	12.2 (2.1-22.4)	800
	25-34	11.8 (0.0-23.9)	400	48.8 (32.3-65.4)	1,700	31.2 (18.8-43.6)	2,100
	35-44	36.6 (17.9-55.4)	1,900	42.7 (28.2-57.3)	2,000	39.5 (27.3-51.8)	3,900
	45-54	32.4 (18.4-46.4)	1,400	36.8 (24.1-49.5)	1,200	34.3 (24.6-44.0)	2,700
	55-64	15.7 (5.2-26.3)	500	25.3 (15.1-35.6)	700	19.9 (12.4-27.4)	1,200
	65-74	14.3 (3.8-24.9)	300	20.6 (10.6-30.5)	500	17.4 (10.1-24.7)	800
NSW	75+	15.8 (0.0-33.2)	200	23.9 (9.6-38.3)	400	20.8 (9.7-31.9)	600
	All Ages	11.6 (10.8-12.3)	253,200	14.6 (13.9-15.3)	333,000	13.1 (12.6-13.6)	587,100
	16-24	5.2 (3.8-6.7)	18,800	9.9 (8.1-11.7)	34,200	7.5 (6.3-8.6)	53,000
	25-34	12.8 (10.6-15.0)	53,300	16.0 (14.2-17.8)	70,000	14.4 (13.0-15.8)	123,300
	35-44	13.6 (11.7-15.5)	59,500	19.5 (17.6-21.4)	87,700	16.6 (15.2-17.9)	147,200
	45-54	14.3 (12.4-16.1)	53,900	16.9 (15.2-18.6)	66,700	15.6 (14.4-16.9)	121,000
	55-64	11.8 (10.2-13.5)	33,400	14.1 (12.6-15.6)	38,600	12.9 (11.8-14.0)	72,100
	65-74	13.2 (11.4-15.0)	25,300	11.3 (9.8-12.7)	23,300	12.2 (11.0-13.3)	48,800
75+	7.2 (5.5-9.0)	8,900	7.5 (6.1-9.0)	12,600	7.4 (6.3-8.5)	21,700	

**Note:** Estimates are based on 661 respondents in North West Slopes and 33,854 in NSW. For this indicator 27 (4.08%) were 'not stated' (Don't know or Refused) in North West Slopes and 1515 (4.48%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who had difficulties getting health care when they needed it. It excludes those who said they do not need health care. The question used to define the indicator was 'Do you have any difficulties getting health care when you need it?'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Emergency department care rated as excellent, very good or good by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	71.6 (56.9-86.4)	3,900	76.3 (61.6-90.9)	4,100	73.9 (63.6-84.3)	8,000
	16-24	62.2 (23.2-100)	900	24.3 (0.0-55.6)	200	47.6 (18.5-76.7)	1,100
	25-34	100 (100-100)	400	61.5 (26.8-96.2)	600	72.0 (44.4-99.5)	1,000
	35-44	59.7 (32.8-86.7)	900	90.8 (79.8-100)	1,100	74.1 (57.1-91.0)	2,000
	45-54	80.7 (52.1-100)	600	91.7 (79.8-100)	600	86.0 (70.7-100)	1,200
	55-64	86.6 (68.9-100)	800	100 (100-100)	300	90.3 (77.3-100)	1,100
	65-74	76.2 (37.0-100)	300	97.8 (93.3-100)	800	90.7 (75.7-100)	1,000
	75+	65.8 (16.6-100)	200	93.2 (80.0-100)	500	84.3 (63.1-100)	600
NSW	All Ages	78.7 (76.1-81.4)	256,800	76.6 (74.2-78.9)	241,900	77.7 (75.9-79.4)	500,200
	16-24	74.1 (67.2-81.1)	48,600	67.1 (59.8-74.4)	36,100	70.8 (65.8-75.8)	85,000
	25-34	73.6 (66.6-80.7)	46,600	69.9 (63.9-75.9)	49,100	71.7 (67.1-76.3)	95,600
	35-44	77.2 (70.8-83.5)	44,800	78.0 (72.7-83.2)	43,100	77.6 (73.5-81.7)	88,100
	45-54	74.7 (67.3-82.1)	35,000	76.3 (70.6-81.9)	33,600	75.4 (70.7-80.0)	68,800
	55-64	86.0 (80.2-91.7)	32,700	81.6 (76.5-86.7)	28,000	84.0 (80.1-87.8)	61,000
	65-74	90.6 (86.7-94.4)	29,800	87.0 (82.5-91.5)	25,100	89.0 (86.0-91.9)	55,100
	75+	89.7 (84.5-94.9)	19,400	91.7 (88.2-95.2)	27,000	90.9 (87.9-93.9)	46,600

**Note:** Estimates are based on 158 respondents in North West Slopes and 5,672 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 45 (0.79%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those attending an emergency department in the last 12 months who rated the care as excellent, very good, or good for their most recent visit. The question used to define the indicator was 'In the last 12 months, have you attended a hospital emergency department or casualty for your own medical care?' and 'Overall, what do you think of the care you received at this emergency department? Was it excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Hospital care rated as excellent, very good or good by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	96.5 (91.7-100)	3,800	95.1 (89.2-100)	3,800	95.8 (92.0-99.7)	7,600
	16-24	100 (100-100)	500	89.1 (64.7-100)	500	94.4 (82.7-100)	1,000
	25-34	89.9 (69.9-100)	600	91.2 (74.1-100)	1,000	90.7 (77.7-100)	1,600
	35-44	100 (100-100)	200	100 (100-100)	700	100 (100-100)	900
	45-54	94.2 (82.6-100)	1,100	100 (100-100)	500	95.9 (87.6-100)	1,600
	55-64	100 (100-100)	600	100 (100-100)	200	100 (100-100)	800
	65-74	100 (100-100)	400	100 (100-100)	400	100 (100-100)	800
	75+	100 (100-100)	300	93.5 (80.9-100)	500	95.8 (87.6-100)	800
NSW	All Ages	92.4 (90.6-94.2)	247,800	89.9 (88.3-91.6)	329,500	91.0 (89.8-92.2)	579,300
	16-24	90.4 (83.9-96.9)	29,200	84.7 (78.1-91.3)	35,400	87.2 (82.5-91.9)	64,700
	25-34	90.5 (85.2-95.9)	33,400	89.7 (85.9-93.4)	90,000	89.9 (86.8-93.0)	123,400
	35-44	87.1 (79.3-94.9)	26,700	89.1 (84.8-93.3)	54,100	88.5 (84.6-92.3)	81,200
	45-54	90.3 (84.6-95.9)	38,700	91.1 (87.4-94.8)	42,400	90.7 (87.4-94.0)	81,400
	55-64	95.5 (93.0-98.0)	40,800	88.8 (84.7-92.9)	33,500	92.4 (90.0-94.8)	74,800
	65-74	94.8 (92.1-97.4)	39,900	92.4 (89.5-95.3)	33,700	93.5 (91.5-95.5)	73,800
	75+	96.2 (93.5-99.0)	39,100	94.4 (92.1-96.7)	40,500	95.3 (93.5-97.1)	80,000

**Note:** Estimates are based on 118 respondents in North West Slopes and 5,451 in NSW. For this indicator 1 (0.85%) were 'not stated' (Don't know or Refused) in North West Slopes and 9 (0.17%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those admitted to hospital in the last 12 months who rated the care as excellent, very good, or good for their most recent overnight stay. The questions used to define the indicator were 'In the last 12 months, have you stayed for at least 1 night in hospital?' and 'Overall, what do you think of the care you received at this hospital? Was it excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Attended a community event at least once in the last 6 months by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	58.5 (50.7-66.4)	9,100	74.5 (68.0-81.1)	12,100	66.8 (61.7-71.9)	21,300
	16-24	68.1 (45.2-90.9)	1,400	59.6 (34.0-85.3)	1,500	63.3 (45.4-81.2)	2,900
	25-34	42.0 (17.6-66.5)	900	79.1 (64.6-93.7)	2,200	62.8 (46.9-78.7)	3,100
	35-44	63.3 (45.4-81.1)	2,100	82.2 (70.2-94.2)	3,000	73.5 (63.2-83.9)	5,300
	45-54	63.4 (48.6-78.2)	2,100	78.7 (66.6-90.8)	2,100	70.2 (60.3-80.1)	4,100
	55-64	50.5 (34.7-66.4)	1,300	76.1 (64.8-87.3)	1,500	61.5 (50.6-72.5)	2,700
	65-74	65.1 (47.5-82.7)	1,000	68.5 (56.0-81.0)	1,000	66.8 (56.0-77.6)	2,100
	75+	51.7 (28.0-75.4)	400	68.4 (49.9-86.9)	800	62.2 (47.7-76.7)	1,100
NSW	All Ages	53.7 (52.3-55.1)	860,100	61.5 (60.4-62.6)	1,017,900	57.5 (56.6-58.4)	1,881,900
	16-24	50.1 (46.3-54.0)	131,100	59.0 (55.4-62.6)	152,500	54.4 (51.7-57.0)	284,300
	25-34	51.7 (47.9-55.4)	160,500	61.0 (58.1-63.9)	193,200	56.3 (53.9-58.6)	353,800
	35-44	65.1 (61.9-68.4)	208,300	72.8 (70.3-75.3)	236,700	68.9 (66.8-70.9)	445,800
	45-54	57.5 (54.4-60.7)	163,000	63.6 (61.0-66.2)	180,700	60.4 (58.4-62.5)	344,200
	55-64	48.4 (45.3-51.5)	97,900	58.7 (56.1-61.3)	116,100	53.4 (51.4-55.4)	214,900
	65-74	45.9 (42.8-49.0)	64,300	53.9 (51.1-56.8)	82,000	50.0 (47.9-52.1)	147,000
	75+	41.7 (37.4-45.9)	34,900	46.9 (43.9-50.0)	56,700	44.5 (42.0-47.0)	92,000

**Note:** Estimates are based on 529 respondents in North West Slopes and 25,602 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in North West Slopes and 44 (0.17%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who have attended at least one community event in the last six months. The question used was "In the past six months, how often have you attended a local community event such as a church or school fete, school concert, or a street fair?".

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Most people can be trusted by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	74.3 (67.3-81.4)	11,500	68.6 (61.4-75.7)	10,900	71.2 (66.1-76.2)	22,400
	16-24	72.5 (50.3-94.7)	1,400	46.4 (21.9-71.0)	1,200	57.8 (39.7-75.9)	2,600
	25-34	70.2 (46.9-93.6)	1,500	75.2 (57.3-93.0)	2,100	73.0 (58.7-87.3)	3,600
	35-44	70.5 (51.6-89.5)	2,400	64.2 (49.0-79.5)	2,300	66.2 (54.2-78.3)	4,700
	45-54	81.4 (70.1-92.8)	2,600	73.8 (60.8-86.7)	1,900	78.1 (69.5-86.7)	4,500
	55-64	67.8 (53.5-82.1)	1,700	76.3 (64.4-88.2)	1,400	71.3 (61.6-81.0)	3,100
	65-74	86.5 (74.3-98.7)	1,300	71.4 (58.8-84.0)	1,100	79.0 (70.1-88.0)	2,400
	75+	75.1 (55.9-94.3)	500	88.5 (78.3-98.8)	1,000	83.5 (73.7-93.4)	1,500
NSW	All Ages	70.2 (68.9-71.5)	1,101,100	65.3 (64.1-66.4)	1,045,900	67.6 (66.8-68.5)	2,154,500
	16-24	65.8 (62.2-69.5)	168,400	57.3 (53.7-60.9)	144,600	61.5 (58.9-64.1)	314,400
	25-34	66.6 (63.0-70.1)	202,900	60.7 (57.8-63.7)	186,100	63.6 (61.3-65.9)	389,400
	35-44	69.6 (66.4-72.7)	218,200	66.3 (63.6-69.0)	210,300	67.9 (65.8-70.0)	430,000
	45-54	74.5 (71.7-77.3)	208,300	69.3 (66.8-71.9)	192,000	71.9 (70.0-73.8)	401,100
	55-64	71.8 (69.0-74.7)	143,200	67.0 (64.5-69.6)	128,100	69.3 (67.4-71.3)	272,200
	65-74	72.6 (69.8-75.5)	98,900	68.5 (65.7-71.4)	99,900	70.5 (68.5-72.5)	200,000
	75+	76.5 (72.7-80.3)	61,200	75.1 (72.4-77.9)	84,800	75.8 (73.5-78.0)	147,400

**Note:** Estimates are based on 519 respondents in North West Slopes and 24,837 in NSW. For this indicator 10 (1.93%) were 'not stated' (Don't know or Refused) in North West Slopes and 809 (3.26%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who strongly agree, or agree that most people can be trusted. The question used was "Most people can be trusted. Do you strongly agree, agree, disagree or strongly disagree?".

# Report of the New South Wales Population Health Survey

## North West Slopes Division of General Practice Visit neighbours by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
North West Slopes	All Ages	73.4 (65.8-81.0)	11,400	69.0 (62.2-75.9)	11,200	70.9 (65.8-76.0)	22,600
	16-24	88.9 (76.2-100)	1,800	69.2 (43.7-94.6)	1,800	77.7 (61.3-94.2)	3,600
	25-34	69.0 (43.5-94.5)	1,500	61.9 (43.7-80.2)	1,700	65.1 (49.9-80.2)	3,200
	35-44	58.7 (38.0-79.3)	2,000	70.3 (56.3-84.3)	2,600	63.8 (51.2-76.3)	4,600
	45-54	81.1 (69.3-93.0)	2,600	66.8 (53.0-80.6)	1,700	74.8 (65.6-83.9)	4,400
	55-64	68.7 (52.9-84.5)	1,700	80.0 (69.6-90.5)	1,500	73.6 (63.3-83.9)	3,300
	65-74	83.1 (69.8-96.5)	1,300	68.7 (55.5-81.9)	1,000	76.0 (66.4-85.6)	2,300
	75+	73.6 (53.8-93.4)	500	68.9 (49.1-88.6)	800	70.6 (56.1-85.2)	1,300
NSW	All Ages	68.1 (66.7-69.4)	1,089,900	63.8 (62.6-64.9)	1,054,800	65.9 (65.0-66.7)	2,153,700
	16-24	74.6 (71.3-77.9)	195,200	64.0 (60.5-67.5)	165,500	69.3 (66.9-71.7)	363,100
	25-34	66.8 (63.3-70.3)	207,900	63.6 (60.7-66.4)	201,400	65.2 (62.9-67.4)	410,100
	35-44	65.9 (62.6-69.3)	210,400	66.7 (64.1-69.4)	216,600	66.4 (64.3-68.5)	428,800
	45-54	66.5 (63.4-69.6)	188,500	61.1 (58.4-63.8)	173,500	63.7 (61.7-65.8)	362,800
	55-64	68.0 (65.0-71.0)	137,200	63.5 (60.9-66.1)	125,400	65.7 (63.8-67.7)	263,900
	65-74	67.2 (64.2-70.3)	94,200	65.4 (62.7-68.2)	99,600	66.2 (64.1-68.3)	194,600
	75+	67.6 (63.5-71.6)	56,600	60.2 (57.2-63.3)	72,800	63.1 (60.7-65.6)	130,400

**Note:** Estimates are based on 528 respondents in North West Slopes and 25,605 in NSW. For this indicator 1 (0.19%) were 'not stated' (Don't know or Refused) in North West Slopes and 41 (0.16%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who visited someone in their neighbourhood at least once in the past week. The question used was "How often have you visited someone in your neighbourhood in the past week?"