

# New South Wales Population Health Survey

2004

## REPORT ON ADULT HEALTH



SOUTH EASTERN SYDNEY  
DIVISION OF GENERAL PRACTICE

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice

---

**Acknowledgements:**

The development and preparation of this edition of the General Practice Division Reports was a team effort involving staff from the NSW Health Survey Program, the Centre for Epidemiology and Research, and the Alliance of NSW Divisions of General Practice. The development of these reports was part funded by the Program for Enhanced Population Health Infrastructure.

**Source:** New South Wales Population Health Survey 2002, 2003 and 2004 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

---

**Produced by:** Centre for Epidemiology and Research, Population Health Division, NSW Department of Health.

**Last updated on:** 26 September 2005

---

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Introduction

---

This report is the first of a series of triennial reports that will be produced for each division of general practice from the New South Wales Population Health Survey. The South Eastern Sydney Division of General Practice Report aims to assist divisions in monitoring the health status of their population, and to support needs assessment, monitoring and improving quality of care in general practice, and policy planning and service development.

The NSW Department of Health in conjunction with the Alliance of NSW Divisions of General Practice has identified a number of indicators for inclusion in the reports. The indicators are:

### **Health behaviours**

- Alcohol risk drinking behaviour
- Fruit intake
- Vegetable intake
- Adequate physical activity
- Smoking status
- Cancer screening
- Influenza immunisation last 12 months 65 years and over
- Pneumococcal immunisation last 5 years 65 years and over
- Homes with a smoke alarm

### **Health services**

- Difficulties getting health care when needing it
- Emergency department care rating
- Hospital care rating

### **Social Capital**

- Attended a community event in the last 6 months
- Most people can be trusted
- Visit neighbours

### **Health Status**


- Excellent, very good or good self-rated health status
- Current asthma
- Diabetes or high blood sugar
- All natural teeth missing
- Overweight and obesity
- High and very high psychological distress

These indicators are presented in graphical and tabular form. For each indicator, the report includes a bar chart of the indicator by age group (2002, 2003 and 2004 South Eastern Sydney data), compared with the whole of New South Wales. The tables below the charts present the indicator estimates for males, females and persons and includes the 95% confidence intervals (CI). The 95% confidence interval provides a range of values that should contain the actual value 95% of the time. In general, a wider confidence interval reflects less certainty in the indicator estimate. If confidence intervals do not overlap then the observed estimates are significantly different. **Source:** New South Wales Population Health Survey 2002, 2003 and 2004 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

---

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Risk alcohol drinking by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	39.2 (30.1-48.4)	24,600	30.0 (23.2-36.8)	18,400	34.7 (28.9-40.4)	43,000
	16-24	35.0 (12.0-58.0)	3,700	29.2 (13.0-45.4)	3,700	31.8 (18.2-45.4)	7,300
	25-34	48.0 (26.6-69.3)	7,100	33.4 (16.2-50.6)	4,200	41.2 (27.3-55.2)	11,400
	35-44	50.2 (27.3-73.1)	5,900	31.1 (13.0-49.2)	3,200	41.2 (25.8-56.7)	9,100
	45-54	29.0 (9.0-49.0)	2,900	34.8 (17.9-51.7)	3,000	31.7 (18.3-45.0)	6,000
	55-64	32.5 (11.6-53.5)	2,500	24.5 (8.7-40.2)	1,700	28.7 (15.4-42.0)	4,300
	65-74	29.7 (10.4-48.9)	1,500	29.0 (8.2-49.8)	1,300	29.4 (15.3-43.5)	2,800
	75+	35.8 (5.4-66.2)	1,000	22.4 (6.2-38.5)	1,200	27.0 (12.0-42.0)	2,200
NSW	All Ages	40.2 (39.0-41.5)	907,800	30.0 (29.0-31.0)	700,800	35.0 (34.2-35.7)	1,610,900
	16-24	49.3 (45.8-52.9)	178,300	44.8 (41.5-48.0)	157,500	46.9 (44.6-49.3)	336,500
	25-34	47.2 (43.8-50.7)	205,100	35.2 (32.7-37.7)	156,600	41.1 (38.9-43.2)	361,700
	35-44	38.0 (35.0-40.9)	169,700	32.1 (29.7-34.5)	146,500	35.0 (33.1-36.9)	317,000
	45-54	36.4 (33.6-39.1)	143,500	25.9 (23.9-28.0)	104,500	31.0 (29.3-32.7)	248,100
	55-64	36.0 (33.3-38.6)	105,500	23.4 (21.4-25.4)	66,900	29.7 (28.0-31.4)	172,600
	65-74	31.2 (28.7-33.7)	62,300	18.1 (16.1-20.1)	39,200	24.4 (22.8-26.0)	101,900
	75+	34.2 (30.6-37.8)	43,400	16.6 (14.6-18.6)	29,500	23.9 (21.9-25.8)	73,100

**Note:** Estimates are based on 418 respondents in South Eastern Sydney and 34,980 in NSW. For this indicator 4 (0.96%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 389 (1.11%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those people who meet Guideline 1 of the NHMRC Australian Alcohol Guidelines, as 1 or more of the following: consuming alcohol every day, consuming on average more than [4 if male/2 if female] standard drinks, consuming more than [6 if male/4 if female] on any 1 occasion or day. Questions used to define the indicator were 'How often do you usually drink alcohol?', 'On a day when you drink alcohol, how many standard drinks do you usually have?', 'In the past 4 weeks have you had more than [7-10 if male/5-6 if female] drinks in a day?', and 'In the past 4 weeks how often have you had [11 or more if male/7 or more if female] drinks in a day?'. Questions used to define the 1997 and 1998 indicator were 'How often do you have an alcoholic drink of any kind?', 'On a day when you have alcoholic drinks, how many standard drinks do you usually have?' and 'On the last occasion that you had more than [4 if male/2 if female] drinks in a day, how many drinks did you actually have?'

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Recommended fruit consumption by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	44.9 (35.4-54.5)	28,300	51.8 (44.2-59.3)	32,300	48.3 (42.3-54.4)	60,600
	16-24	51.1 (25.1-77.1)	5,300	44.3 (25.9-62.8)	5,900	47.3 (32.0-62.7)	11,300
	25-34	62.3 (43.0-81.6)	9,000	57.0 (37.6-76.3)	7,400	59.8 (45.9-73.7)	16,400
	35-44	27.9 (6.7-49.1)	3,300	51.0 (31.1-70.8)	5,300	38.7 (23.8-53.7)	8,500
	45-54	38.7 (16.3-61.1)	4,100	49.0 (30.5-67.4)	4,300	43.3 (28.4-58.3)	8,400
	55-64	43.9 (22.0-65.8)	3,400	58.5 (41.4-75.6)	4,200	50.9 (36.8-65.0)	7,600
	65-74	26.3 (6.1-46.5)	1,400	53.7 (31.4-76.1)	2,400	38.9 (23.9-54.0)	3,800
NSW	75+	64.5 (37.7-91.2)	1,800	53.2 (35.7-70.6)	2,900	57.1 (41.8-72.3)	4,700
	All Ages	40.4 (39.2-41.7)	918,800	52.6 (51.5-53.6)	1,234,800	46.6 (45.8-47.4)	2,159,200
	16-24	39.1 (35.6-42.5)	144,400	44.5 (41.3-47.8)	158,000	41.6 (39.3-44.0)	302,900
	25-34	32.9 (29.7-36.2)	142,700	44.9 (42.2-47.6)	200,200	39.0 (36.8-41.1)	343,000
	35-44	38.2 (35.3-41.2)	171,800	50.4 (47.9-53.0)	231,000	44.4 (42.4-46.4)	403,900
	45-54	41.9 (39.0-44.7)	166,600	54.7 (52.4-57.1)	222,100	48.3 (46.5-50.2)	389,200
	55-64	44.0 (41.2-46.7)	129,800	60.3 (58.0-62.6)	173,500	52.0 (50.2-53.8)	304,500
65-74	50.0 (47.2-52.8)	99,900	63.0 (60.6-65.4)	137,100	56.8 (55.0-58.6)	237,900	
75+	50.5 (46.7-54.3)	63,600	63.4 (60.7-66.1)	112,700	58.2 (56.0-60.4)	177,900	

**Note:** Estimates are based on 420 respondents in South Eastern Sydney and 35,162 in NSW. For this indicator 2 (0.48%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 207 (0.59%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who meet the recommended fruit consumption. The recommended consumption of fruit according to the National Campaign is 2 serves for people aged 16 years and over. One serve is equivalent to 1 medium piece or 2 small pieces of fruit. The question used to define the indicator was 'How many serves of fruit do you usually eat each day?'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Recommended vegetable consumption by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	7.1 (2.5-11.7)	4,500	17.1 (11.1-23.2)	10,700	12.1 (8.2-16.0)	15,100
	16-24	1.9 (0.0-5.8)	200	21.5 (5.4-37.6)	2,800	12.8 (3.0-22.7)	3,000
	25-34	12.0 (0.0-26.5)	1,700	17.9 (3.9-31.8)	2,300	14.8 (4.7-24.9)	4,100
	35-44	2.0 (0.0-5.9)	200	26.3 (7.5-45.0)	2,700	13.4 (3.3-23.5)	2,900
	45-54	3.1 (0.0-9.2)	300	5.4 (0.0-11.7)	500	4.1 (0.0-8.6)	800
	55-64	15.0 (0.0-32.8)	1,200	22.7 (8.2-37.3)	1,600	18.7 (7.1-30.3)	2,800
	65-74	8.7 (0.0-21.9)	500	6.9 (0.0-15.0)	300	7.9 (0.0-16.0)	800
	75+	14.2 (0.0-33.8)	400	7.4 (0.0-17.4)	400	9.6 (0.4-18.7)	800
NSW	All Ages	7.0 (6.5-7.6)	159,600	13.6 (12.9-14.3)	320,100	10.4 (9.9-10.9)	479,700
	16-24	5.4 (4.0-6.8)	19,800	8.0 (6.2-9.7)	28,300	6.7 (5.5-7.8)	48,100
	25-34	5.2 (3.8-6.7)	22,700	9.2 (7.5-10.8)	40,900	7.2 (6.1-8.3)	63,600
	35-44	5.2 (4.0-6.5)	23,600	11.6 (10.0-13.3)	53,300	8.5 (7.4-9.5)	76,900
	45-54	7.1 (5.7-8.5)	28,200	17.8 (16.0-19.6)	72,200	12.5 (11.3-13.7)	100,400
	55-64	9.2 (7.7-10.8)	27,200	19.3 (17.5-21.1)	55,400	14.2 (13.0-15.4)	82,600
	65-74	11.9 (10.1-13.7)	23,700	20.4 (18.4-22.5)	44,300	16.3 (15.0-17.7)	68,000
	75+	11.5 (9.2-13.9)	14,400	14.4 (12.4-16.4)	25,700	13.2 (11.7-14.8)	40,100

**Note:** Estimates are based on 417 respondents in South Eastern Sydney and 35,000 in NSW. For this indicator 5 (1.2%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 369 (1.05%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who meet the recommended consumption of vegetables. The recommended vegetable intake according to the National Campaign is 5 serves per day for persons aged 16 years and over. One serve is equivalent to 1/2 cup of cooked vegetables or 1 cup of salad vegetables. The question used to define the indicator was: 'How many serves of vegetables do you usually eat each day?'

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Adequate physical activity by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	58.2 (48.9-67.5)	36,900	52.0 (44.5-59.5)	32,500	55.1 (49.2-61.1)	69,300
	16-24	91.2 (80.4-100)	9,500	70.3 (55.0-85.7)	9,400	79.5 (69.1-89.8)	18,900
	25-34	62.6 (41.4-83.8)	9,300	57.7 (38.7-76.7)	7,500	60.3 (45.9-74.7)	16,800
	35-44	38.5 (17.3-59.6)	4,500	53.9 (34.3-73.4)	5,600	45.7 (30.6-60.8)	10,000
	45-54	48.3 (25.2-71.3)	5,100	54.5 (36.2-72.7)	4,800	51.1 (35.9-66.2)	9,900
	55-64	52.5 (30.7-74.4)	4,100	39.7 (22.8-56.7)	2,800	46.4 (32.3-60.5)	6,900
	65-74	61.7 (40.5-82.8)	3,200	18.3 (4.2-32.3)	800	41.7 (26.5-56.8)	4,000
	75+	41.6 (10.6-72.6)	1,200	29.5 (14.6-44.3)	1,600	33.6 (18.9-48.3)	2,800
NSW	All Ages	52.5 (51.2-53.8)	1,199,100	44.0 (43.0-45.1)	1,037,600	48.1 (47.3-49.0)	2,241,000
	16-24	66.8 (63.4-70.1)	247,000	57.5 (54.4-60.7)	205,200	62.2 (59.9-64.5)	453,800
	25-34	53.7 (50.3-57.1)	234,900	46.5 (43.9-49.2)	208,200	50.0 (47.8-52.2)	443,200
	35-44	49.4 (46.4-52.5)	223,500	43.6 (41.1-46.2)	200,700	46.5 (44.5-48.5)	425,200
	45-54	49.5 (46.6-52.4)	197,600	43.2 (40.9-45.5)	175,600	46.3 (44.4-48.2)	374,200
	55-64	49.8 (47.0-52.5)	147,500	41.6 (39.3-43.9)	120,000	45.6 (43.8-47.4)	267,500
	65-74	49.6 (46.9-52.4)	99,800	37.3 (34.9-39.7)	81,500	43.2 (41.3-45.0)	181,800
	75+	38.1 (34.4-41.8)	48,600	26.0 (23.6-28.3)	46,500	30.8 (28.8-32.9)	95,200

**Note:** Estimates are based on 422 respondents in South Eastern Sydney and 35,365 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 4 (0.01%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who did adequate physical activity. Adequate physical activity is a total of 150 minutes per week on 5 separate occasions. The total minutes were calculated by adding minutes in the last week spent walking continuously for at least 10 minutes, minutes doing moderate physical activity, plus 2 x minutes doing vigorous physical activity. The questions used to define the indicator were 'In the last week, how many times have you walked continuously for at least 10 minutes for recreation or exercise or to get to or from places?', 'What do you estimate was the total time you spent walking in this way in the last week?', 'In the last week, how many times did you do any vigorous physical activity which made you breathe harder or puff and pant?', 'What do you estimate was the total time you spent doing this vigorous physical activity in the last week?', 'In the last week, how many times did you do any other more moderate physical activity that you have not already mentioned?' and 'What do you estimate was the total time that you spent doing these activities in the last week?'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Current daily or occasional smoking by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	25.5 (17.6-33.5)	16,200	13.3 (8.4-18.2)	8,300	19.5 (14.7-24.2)	24,500
	16-24	21.6 (2.4-40.8)	2,300	14.5 (3.3-25.7)	1,900	17.6 (7.1-28.1)	4,200
	25-34	21.8 (6.2-37.4)	3,200	11.2 (0.0-23.4)	1,500	16.9 (6.9-26.9)	4,700
	35-44	29.9 (9.1-50.7)	3,500	20.6 (5.5-35.7)	2,100	25.5 (12.3-38.8)	5,600
	45-54	46.9 (23.8-70.1)	5,000	19.0 (5.2-32.7)	1,700	34.3 (19.3-49.3)	6,600
	55-64	13.6 (1.0-26.2)	1,100	6.9 (0.0-14.1)	500	10.4 (3.0-17.8)	1,500
	65-74	12.9 (0.0-27.1)	700	5.8 (0.0-14.5)	300	9.6 (0.9-18.4)	900
	75+	17.3 (0.0-37.3)	500	7.2 (0.0-16.1)	400	10.7 (1.7-19.6)	900
NSW	All Ages	23.9 (22.8-25.0)	546,100	19.4 (18.6-20.2)	456,900	21.7 (21.0-22.3)	1,007,600
	16-24	26.7 (23.7-29.8)	98,800	26.0 (23.2-28.8)	92,600	26.6 (24.5-28.6)	193,700
	25-34	32.9 (29.8-36.1)	144,000	25.8 (23.4-28.1)	115,200	29.3 (27.3-31.2)	259,500
	35-44	29.1 (26.5-31.8)	131,400	22.4 (20.5-24.4)	103,100	25.8 (24.1-27.4)	235,300
	45-54	24.3 (21.9-26.8)	97,300	19.6 (17.8-21.4)	79,700	22.0 (20.4-23.5)	177,400
	55-64	15.7 (13.8-17.6)	46,600	14.6 (13.1-16.2)	42,200	15.2 (14.0-16.4)	89,200
	65-74	10.3 (8.7-11.9)	20,700	7.9 (6.6-9.2)	17,200	9.1 (8.1-10.1)	38,300
	75+	5.7 (4.0-7.4)	7,300	3.9 (3.0-4.8)	6,900	4.6 (3.7-5.5)	14,200

**Note:** Estimates are based on 422 respondents in South Eastern Sydney and 35,353 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 16 (0.05%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who smoked daily or occasionally. The question used to define the indicator was 'Which of the following best describes your smoking status? Smoke daily, Smoke occasionally, Do not smoke now, but I used to, I have tried it a few times but never smoked regularly, and I have never smoked'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Smoke-free households by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	72.5 (64.2-80.7)	45,900	88.5 (83.9-93.1)	55,200	80.4 (75.6-85.2)	101,100
	16-24	63.5 (37.0-90.1)	6,600	85.9 (71.6-100)	11,500	76.1 (61.2-91.0)	18,100
	25-34	80.9 (66.6-95.2)	12,000	98.7 (96.2-100)	12,800	89.3 (81.5-97.0)	24,800
	35-44	67.8 (47.3-88.3)	7,900	98.0 (95.3-100)	10,100	82.0 (70.5-93.6)	18,000
	45-54	67.0 (46.4-87.6)	7,100	72.2 (56.2-88.1)	6,300	69.3 (55.9-82.7)	13,400
	55-64	83.4 (69.2-97.5)	6,500	88.7 (79.4-98.0)	6,300	85.9 (77.3-94.5)	12,800
	65-74	77.0 (59.8-94.2)	4,000	79.1 (58.4-99.7)	3,500	77.9 (64.7-91.2)	7,500
NSW	75+	62.6 (31.5-93.7)	1,800	85.8 (75.6-96.0)	4,700	77.8 (64.2-91.4)	6,500
	All Ages	81.7 (80.7-82.6)	1,863,400	83.3 (82.6-84.1)	1,962,200	82.5 (81.9-83.1)	3,837,200
	16-24	76.8 (73.9-79.7)	283,100	79.3 (76.7-82.0)	282,800	78.1 (76.1-80.0)	568,700
	25-34	81.4 (79.0-83.9)	356,100	84.6 (82.7-86.5)	378,600	83.0 (81.5-84.6)	735,800
	35-44	83.4 (81.4-85.5)	376,500	84.1 (82.4-85.9)	386,500	83.8 (82.4-85.1)	764,800
	45-54	79.5 (77.4-81.7)	317,900	81.8 (80.1-83.5)	332,300	80.7 (79.3-82.0)	651,600
	55-64	82.0 (80.0-84.1)	243,200	82.0 (80.3-83.8)	236,300	82.0 (80.6-83.4)	481,300
	65-74	86.0 (84.1-87.9)	172,900	84.4 (82.5-86.4)	184,300	85.1 (83.8-86.5)	358,400
75+	89.1 (86.8-91.5)	113,700	90.3 (88.8-91.8)	161,400	89.7 (88.4-91.0)	276,500	

**Note:** Estimates are based on 422 respondents in South Eastern Sydney and 35,328 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 39 (0.11%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those households who indicated that their home was smoke-free. The question used to define the indicator was 'Which of the following best describes your home situation? My home is smoke-free, People occasionally smoke in the house, and People frequently smoke in the house'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Homes with a smoke alarm or detector by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	67.9 (59.6-76.3)	42,600	60.7 (53.3-68.2)	37,500	64.4 (58.7-70.0)	80,200
	16-24	63.9 (39.4-88.5)	6,700	68.3 (52.3-84.3)	9,100	66.4 (52.3-80.5)	15,800
	25-34	68.8 (50.6-86.9)	9,900	43.7 (25.0-62.3)	5,600	57.0 (42.6-71.3)	15,500
	35-44	73.1 (54.5-91.8)	8,500	63.5 (44.8-82.2)	6,300	68.7 (55.4-82.1)	14,800
	45-54	69.9 (48.9-90.8)	7,400	56.9 (38.4-75.4)	5,000	64.0 (49.6-78.4)	12,400
	55-64	73.1 (54.7-91.6)	5,500	58.0 (40.7-75.3)	4,100	65.8 (52.6-78.9)	9,600
	65-74	61.2 (40.8-81.7)	3,200	72.1 (54.3-89.9)	3,200	66.3 (52.4-80.1)	6,400
	75+	48.5 (17.4-79.6)	1,400	77.8 (63.5-92.0)	4,200	67.7 (53.0-82.5)	5,600
NSW	All Ages	71.8 (70.7-72.9)	1,599,000	72.6 (71.7-73.6)	1,676,900	72.2 (71.5-72.9)	3,286,100
	16-24	72.1 (68.9-75.2)	258,400	69.9 (67.0-72.9)	244,300	71.1 (68.9-73.2)	505,500
	25-34	71.4 (68.3-74.4)	302,700	70.1 (67.6-72.6)	307,300	70.7 (68.7-72.7)	610,700
	35-44	74.1 (71.5-76.8)	326,200	78.0 (75.9-80.1)	350,100	76.1 (74.4-77.8)	678,200
	45-54	72.8 (70.2-75.5)	284,400	71.5 (69.3-73.7)	284,500	72.1 (70.4-73.8)	569,900
	55-64	68.7 (66.1-71.3)	200,100	70.9 (68.8-73.1)	201,600	69.8 (68.2-71.5)	403,500
	65-74	71.1 (68.5-73.7)	140,200	75.6 (73.4-77.7)	162,900	73.3 (71.6-75.0)	304,000
	75+	69.5 (65.9-73.0)	87,000	72.0 (69.5-74.5)	126,200	70.8 (68.8-72.9)	214,300

**Note:** Estimates are based on 417 respondents in South Eastern Sydney and 34,762 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 0 (0%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who have a smoke alarm or detector in their home. The question used to define the indicator was 'Do you have any of the following fire safety measures in your home? Fire alarm (hard wired), Fire alarm (battery operated only), Fire sprinkler system, Safety switch or circuit breaker, Fire extinguisher, Fire evacuation plan, External water supply, External sprinkler'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Screening mammogram within the last 2 years by age




Division	Age (years)	Females (95% CI)	Females (est. no.)
South Eastern Sydney	All Ages	87.5 (71.9-100)	3,400
	50-54	86.7 (60.1-100)	1,500
	55-59	100 (100-100)	800
	60-64	88.9 (65.5-100)	600
	65-69	71.4 (18.1-100)	500
NSW	All Ages	75.6 (73.0-78.2)	130,300
	50-54	70.6 (65.5-75.7)	39,300
	55-59	77.3 (72.6-82.1)	34,400
	60-64	82.5 (78.5-86.6)	32,200
	65-69	73.4 (66.6-80.2)	24,500

**Note:** Estimates are based on 23 respondents in South Eastern Sydney and 2,202 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 5 (0.23%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who have had a screening mammogram in the last 2 years. This excludes women who had a mammogram for a breast problem or had had breast cancer in the past. The questions used to define the indicator were "Have you ever had a mammogram?" and "When did you last have a mammogram?" and "Can you tell me all the reasons why you had your last mammogram?".

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Pap test within the last 2 years by age




Division	Age (years)	Females (95% CI)	Females (est. no.)
South Eastern Sydney	All Ages	64.2 (50.7-77.8)	11,700
	20-29	38.9 (14.9-63.0)	2,200
	30-39	66.9 (35.6-98.3)	3,500
	40-49	81.7 (55.1-100)	1,900
	50-59	86.5 (70.0-100)	2,900
	60-69	71.7 (44.7-98.7)	1,200
NSW	All Ages	74.6 (72.8-76.4)	435,300
	20-29	65.1 (60.7-69.6)	95,100
	30-39	78.1 (74.6-81.5)	123,500
	40-49	81.1 (77.9-84.4)	111,000
	50-59	77.8 (74.2-81.4)	71,300
	60-69	67.4 (63.0-71.9)	34,400

**Note:** Estimates are based on 76 respondents in South Eastern Sydney and 4,499 in NSW. For this indicator 1 (1.32%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 23 (0.51%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who have had a Pap test in the last 2 years and have not had a hysterectomy. The questions used to define the indicator were "A Pap test, is a routine test carried out by a doctor. It is recommended for all women for early detection of cancer of the cervix. Have you ever had a Pap test?", "When did you last have a Pap test?" and "Have you ever had a hysterectomy?".

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Vaccinated against influenza in the last 12 months by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	76.5 (60.3-92.7)	6,100	75.0 (62.9-87.0)	7,400	75.7 (65.8-85.5)	13,500
	65-69	75.6 (47.7-100)	1,900	67.9 (38.9-96.9)	1,100	72.6 (52.2-93.0)	2,900
	70-75	78.5 (53.9-100)	2,100	65.5 (36.1-94.8)	1,900	71.7 (52.0-91.4)	4,000
	75+	75.5 (44.9-100)	2,100	82.0 (70.4-93.6)	4,500	79.8 (66.6-93.0)	6,600
NSW	All Ages	75.5 (73.5-77.4)	247,700	76.3 (74.8-77.8)	302,900	75.9 (74.7-77.1)	553,300
	65-69	65.2 (61.6-68.7)	69,900	68.8 (65.7-72.0)	77,000	66.9 (64.5-69.3)	147,100
	70-75	77.2 (73.8-80.6)	72,400	77.3 (74.4-80.2)	82,200	77.2 (75.0-79.4)	155,300
	75+	82.9 (79.9-85.8)	105,500	80.4 (78.3-82.6)	143,700	81.5 (79.7-83.2)	250,900


**Note:** Estimates are based on 106 respondents in South Eastern Sydney and 9,684 in NSW. For this indicator 1 (0.94%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 12 (0.12%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those respondents that are aged 65 years and over who have been vaccinated against flu in the past 12 months.

The question used was 'Were you vaccinated or immunised against flu in the past 12 months?'

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Vaccinated against pneumococcal disease in the last 5 years by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	47.7 (29.1-66.2)	3,700	39.6 (26.1-53.1)	3,800	43.2 (31.9-54.5)	7,500
	65-69	27.4 (0.0-56.1)	600	18.2 (0.0-39.6)	300	23.7 (4.3-43.2)	900
	70-75	68.6 (41.2-96.0)	1,700	29.3 (4.4-54.2)	800	47.6 (25.7-69.5)	2,600
	75+	35.0 (6.8-63.1)	800	53.5 (35.2-71.8)	2,700	47.6 (31.6-63.5)	3,500
NSW	All Ages	42.3 (40.0-44.5)	134,100	47.1 (45.2-48.9)	182,800	44.9 (43.5-46.3)	318,300
	65-69	27.5 (24.1-30.9)	28,400	32.2 (28.8-35.5)	35,200	29.9 (27.5-32.3)	63,700
	70-75	42.0 (37.8-46.1)	38,000	49.2 (45.8-52.7)	51,500	45.8 (43.1-48.5)	89,700
	75+	56.7 (52.6-60.7)	60,200	54.8 (51.9-57.7)	84,800	55.6 (53.2-57.9)	146,000

**Note:** Estimates are based on 103 respondents in South Eastern Sydney and 9,430 in NSW. For this indicator 4 (3.88%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 266 (2.82%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those respondents that are aged 65 years and over who have been vaccinated against pneumococcal disease in the last 5 years. The question used was "When were you last vaccinated or immunised against pneumonia?".

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice

### Excellent, very good, or good self-rated health status by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	81.0 (73.9-88.1)	51,300	78.9 (72.8-85.1)	49,200	80.0 (75.3-84.7)	100,400
	16-24	82.8 (58.5-100)	8,700	79.6 (62.5-96.6)	10,700	81.0 (66.7-95.3)	19,300
	25-34	94.2 (87.0-100)	13,900	95.3 (88.9-100)	12,400	94.7 (89.8-99.6)	26,300
	35-44	82.6 (65.7-99.4)	9,600	76.0 (58.4-93.6)	7,800	79.5 (67.2-91.8)	17,500
	45-54	83.8 (69.9-97.7)	8,900	70.6 (54.8-86.5)	6,200	77.8 (67.1-88.6)	15,100
	55-64	57.7 (35.5-79.9)	4,500	71.8 (56.6-87.1)	5,100	64.5 (50.4-78.5)	9,600
	65-74	66.2 (46.3-86.2)	3,400	71.3 (50.2-92.4)	3,200	68.6 (54.1-83.0)	6,600
	75+	78.8 (55.8-100)	2,200	72.5 (58.2-86.8)	3,800	74.7 (62.4-87.1)	6,100
NSW	All Ages	81.2 (80.2-82.2)	1,852,000	79.8 (79.0-80.6)	1,878,300	80.5 (79.8-81.1)	3,741,600
	16-24	85.1 (82.4-87.8)	314,200	83.2 (80.8-85.5)	296,600	84.2 (82.4-86.0)	613,600
	25-34	84.6 (82.1-87.2)	370,200	85.4 (83.5-87.2)	381,700	85.0 (83.4-86.6)	753,200
	35-44	83.2 (81.1-85.3)	376,100	82.0 (80.1-83.8)	376,800	82.6 (81.2-84.0)	754,800
	45-54	81.6 (79.4-83.7)	325,300	80.0 (78.2-81.8)	324,900	80.7 (79.3-82.1)	651,700
	55-64	76.1 (73.7-78.5)	225,600	75.5 (73.5-77.5)	217,500	75.7 (74.1-77.3)	444,000
	65-74	77.1 (74.7-79.4)	154,600	73.2 (71.0-75.5)	159,500	75.0 (73.4-76.7)	315,300
	75+	68.0 (64.5-71.5)	86,100	68.0 (65.5-70.6)	121,400	68.1 (66.0-70.2)	209,000

**Note:** Estimates are based on 421 respondents in South Eastern Sydney and 35,324 in NSW. For this indicator 1 (0.24%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 45 (0.13%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those responding excellent, very good, or good to a global self-rated health status question. The question used to define the indicator was 'Overall, how would you rate your health during the past 4 weeks? Was it excellent, very good, good, fair, poor, or very poor'. Question used to define the indicator in 1997 and 1998 was 'In general would you say your health is excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Current asthma by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	6.9 (2.5-11.2)	4,400	10.4 (5.9-14.9)	6,500	8.6 (5.5-11.7)	10,800
	16-24	. (-.)	.	14.4 (1.0-27.8)	1,900	8.1 (0.2-16.0)	1,900
	25-34	5.4 (0.0-12.3)	800	9.0 (0.0-19.1)	1,200	7.1 (1.1-13.1)	2,000
	35-44	7.8 (0.0-16.9)	900	5.4 (0.0-12.5)	600	6.7 (0.8-12.5)	1,500
	45-54	13.2 (0.0-31.9)	1,400	9.9 (0.9-18.8)	900	11.7 (0.5-22.8)	2,300
	55-64	8.5 (0.0-18.8)	700	10.9 (0.0-22.1)	800	9.6 (2.0-17.3)	1,400
	65-74	8.3 (0.0-19.8)	400	19.3 (1.9-36.6)	900	13.4 (3.1-23.6)	1,300
	75+	5.0 (0.0-14.8)	100	5.8 (0.0-16.8)	300	5.5 (0.0-13.5)	500
NSW	All Ages	9.1 (8.4-9.9)	208,300	12.3 (11.6-12.9)	288,700	10.7 (10.2-11.2)	498,500
	16-24	11.8 (9.4-14.2)	43,600	15.0 (12.9-17.1)	53,400	13.3 (11.7-14.9)	97,100
	25-34	12.0 (9.5-14.5)	52,400	10.7 (9.1-12.3)	47,900	11.3 (9.9-12.8)	100,300
	35-44	6.8 (5.4-8.2)	30,600	10.3 (8.9-11.7)	47,300	8.6 (7.6-9.6)	78,300
	45-54	8.1 (6.5-9.6)	32,300	13.8 (12.2-15.4)	56,000	11.0 (9.8-12.1)	88,500
	55-64	7.4 (6.0-8.8)	22,000	12.9 (11.4-14.4)	37,100	10.2 (9.1-11.2)	59,700
	65-74	8.3 (6.8-9.7)	16,600	13.0 (11.3-14.6)	28,300	10.7 (9.6-11.8)	45,100
	75+	8.4 (6.4-10.4)	10,800	10.5 (8.7-12.2)	18,700	9.6 (8.3-10.9)	29,600

**Note:** Estimates are based on 421 respondents in South Eastern Sydney and 35,333 in NSW. For this indicator 1 (0.24%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 36 (0.1%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those respondents who had symptoms of asthma or had taken treatment for asthma in the last 12 months. The questions used to define the indicator were 'Have you ever been told by a doctor or at a hospital that you have asthma?' and 'Have you had symptoms of asthma or taken treatment for asthma in the last 12 months?'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Diabetes or high blood sugar by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	4.1 (1.5-6.8)	2,600	3.4 (1.2-5.6)	2,100	3.8 (2.1-5.5)	4,800
	16-24	. (-.)	.	3.4 (0.0-10.1)	500	1.9 (0.0-5.7)	500
	25-34	1.3 (0.0-3.8)	200	. (-.)	.	0.7 (0.0-2.0)	200
	35-44	. (-.)	.	. (-.)	.	. (-.)	.
	45-54	2.6 (0.0-7.6)	300	. (-.)	.	1.4 (0.0-4.2)	300
	55-64	8.9 (0.0-20.7)	700	8.2 (0.0-16.6)	600	8.5 (1.2-15.9)	1,300
	65-74	16.9 (0.0-34.6)	900	12.3 (0.0-24.6)	500	14.8 (3.6-26.0)	1,400
	75+	21.0 (0.0-43.8)	600	10.2 (0.0-21.3)	600	13.9 (3.2-24.5)	1,200
NSW	All Ages	7.2 (6.6-7.7)	163,400	5.5 (5.1-5.9)	128,900	6.3 (6.0-6.7)	294,200
	16-24	0.9 (0.3-1.5)	3,300	1.8 (0.9-2.7)	6,400	1.4 (0.8-1.9)	9,900
	25-34	2.7 (1.5-3.8)	11,600	1.8 (1.2-2.5)	8,100	2.2 (1.6-2.9)	19,700
	35-44	3.5 (2.5-4.6)	16,000	2.6 (1.9-3.3)	12,000	3.1 (2.5-3.7)	28,000
	45-54	8.0 (6.6-9.5)	32,000	5.3 (4.3-6.3)	21,400	6.7 (5.8-7.6)	54,100
	55-64	16.0 (13.9-18.0)	47,200	9.9 (8.5-11.3)	28,500	13.0 (11.7-14.3)	76,200
	65-74	17.6 (15.4-19.7)	35,200	14.2 (12.4-16.0)	30,800	15.8 (14.4-17.2)	66,300
	75+	14.3 (11.9-16.8)	18,200	12.1 (10.3-13.9)	21,700	13.0 (11.6-14.5)	40,100

**Note:** Estimates are based on 422 respondents in South Eastern Sydney and 35,248 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 8 (0.02%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who either had diabetes or high blood sugar levels but did not have gestational diabetes. The questions used to define the indicator were 'Have you ever been told by a doctor or at a hospital that you have diabetes?', 'Have you ever been told by a doctor or at a hospital that you have high sugar levels in your blood or urine?' and if female 'Were you pregnant when you were first told that you had diabetes or high blood sugar?' and 'Have you ever had diabetes or high blood sugar apart from when you were pregnant?'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice All natural teeth missing by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	5.2 (1.9-8.5)	3,300	5.6 (3.2-8.1)	3,500	5.4 (3.3-7.4)	6,800
	16-24	. (-.)	.	. (-.)	.	. (-.)	.
	25-34	. (-.)	.	. (-.)	.	. (-.)	.
	35-44	. (-.)	.	. (-.)	.	. (-.)	.
	45-54	7.3 (0.0-18.5)	800	. (-.)	.	4.0 (0.0-10.2)	800
	55-64	4.3 (0.0-10.7)	300	2.1 (0.0-5.0)	100	3.3 (0.0-6.9)	500
	65-74	9.3 (0.0-19.7)	500	26.4 (8.3-44.6)	1,200	17.2 (6.9-27.5)	1,700
	75+	59.1 (30.5-87.8)	1,700	40.0 (23.1-57.0)	2,200	46.6 (30.9-62.3)	3,900
NSW	All Ages	4.5 (4.1-4.9)	103,700	7.8 (7.3-8.2)	182,900	6.2 (5.9-6.5)	288,800
	16-24	0.3 (0.0-0.7)	1,200	1.0 (0.1-1.9)	3,600	0.7 (0.2-1.1)	4,800
	25-34	0.2 (0.0-0.5)	1,100	0.4 (0.0-0.9)	2,000	0.3 (0.1-0.6)	3,100
	35-44	0.6 (0.2-1.1)	2,900	1.3 (0.7-1.8)	5,800	1.0 (0.6-1.3)	8,700
	45-54	3.5 (2.5-4.5)	14,100	3.9 (3.0-4.8)	15,900	3.7 (3.0-4.4)	30,000
	55-64	6.8 (5.5-8.1)	20,200	11.8 (10.4-13.2)	33,900	9.4 (8.4-10.4)	55,100
	65-74	15.0 (13.1-16.9)	30,200	25.3 (23.1-27.5)	55,300	20.4 (18.9-21.9)	85,900
	75+	26.7 (23.3-30.2)	34,100	37.1 (34.5-39.7)	66,400	32.8 (30.7-34.8)	101,100

**Note:** Estimates are based on 422 respondents in South Eastern Sydney and 35,355 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 14 (0.04%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those respondents who had all of their natural teeth missing. The question used to define the indicator was 'Are any of your natural teeth missing? (Natural teeth does not include dentures but includes wisdom teeth)'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Overweight and obesity by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	48.7 (39.3-58.2)	30,800	29.1 (22.4-35.7)	17,100	39.3 (33.2-45.3)	47,900
	16-24	33.4 (8.9-57.9)	3,500	15.2 (1.2-29.3)	2,000	23.4 (9.7-37.0)	5,500
	25-34	70.0 (52.9-87.0)	10,400	22.8 (7.3-38.4)	2,600	49.4 (34.4-64.4)	13,000
	35-44	36.2 (15.8-56.7)	4,200	25.5 (8.2-42.9)	2,500	31.3 (17.8-44.8)	6,700
	45-54	38.6 (16.6-60.5)	4,100	39.2 (21.5-56.9)	3,400	38.9 (24.4-53.3)	7,500
	55-64	63.1 (41.0-85.1)	4,900	41.3 (24.6-58.0)	2,900	52.7 (38.5-66.8)	7,800
	65-74	53.8 (32.2-75.4)	2,800	42.3 (20.5-64.2)	1,600	49.0 (33.3-64.6)	4,400
	75+	34.9 (2.5-67.2)	900	41.0 (23.0-58.9)	2,100	38.9 (22.7-55.1)	3,000
NSW	All Ages	54.5 (53.3-55.8)	1,223,800	40.0 (39.0-41.0)	891,000	47.3 (46.5-48.1)	2,121,200
	16-24	30.3 (26.9-33.6)	108,600	18.3 (15.9-20.7)	61,200	24.6 (22.5-26.8)	171,400
	25-34	50.6 (47.2-54.1)	217,500	33.4 (30.9-36.0)	140,400	42.2 (40.0-44.3)	359,000
	35-44	61.1 (58.1-64.1)	272,300	39.0 (36.5-41.6)	170,100	50.1 (48.1-52.2)	442,900
	45-54	64.5 (61.6-67.3)	254,200	48.6 (46.2-51.0)	188,600	56.6 (54.7-58.5)	443,400
	55-64	66.2 (63.5-68.9)	193,300	53.0 (50.6-55.3)	146,800	59.8 (58.0-61.6)	341,500
	65-74	61.1 (58.4-63.8)	121,000	55.3 (52.8-57.8)	114,400	58.1 (56.2-59.9)	236,200
	75+	45.5 (41.7-49.2)	56,900	42.2 (39.3-45.0)	69,500	43.5 (41.2-45.8)	126,800

**Note:** Estimates are based on 408 respondents in South Eastern Sydney and 33,809 in NSW. For this indicator 14 (3.43%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 1557 (4.61%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those with a Body Mass Index (BMI) of 25 or higher. The questions used to define the indicator were 'How tall are you without shoes?' and 'How much do you weigh without clothes or shoes?'. The BMI is calculated as follows: BMI = weight(kg)/height<sup>2</sup>(m). Categories for this indicator include overweight (BMI between 25 and 29.9) and obese (BMI of 30 and over).

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice High and very high psychological distress by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	12.3 (6.5-18.1)	7,600	13.3 (8.3-18.4)	8,300	12.8 (9.0-16.7)	15,900
	16-24	23.0 (4.2-41.8)	2,400	24.6 (8.2-40.9)	3,300	23.9 (11.5-36.3)	5,700
	25-34	11.3 (0.3-22.4)	1,600	. (-.)	.	6.0 (0.1-11.8)	1,600
	35-44	12.0 (0.0-27.7)	1,300	12.2 (0.2-24.2)	1,300	12.1 (2.1-22.1)	2,600
	45-54	3.6 (0.0-8.6)	400	21.9 (8.0-35.8)	1,900	11.7 (4.5-18.9)	2,200
	55-64	21.2 (0.0-42.7)	1,500	9.2 (0.1-18.2)	700	15.2 (3.1-27.3)	2,200
	65-74	6.3 (0.0-18.1)	300	9.3 (0.0-20.1)	400	7.7 (0.0-15.7)	700
	75+	. (-.)	.	15.2 (3.3-27.1)	800	9.8 (1.8-17.7)	800
NSW	All Ages	10.4 (9.7-11.2)	235,900	13.9 (13.2-14.7)	324,900	12.2 (11.7-12.7)	562,000
	16-24	11.8 (9.5-14.0)	43,500	19.6 (17.0-22.2)	69,700	15.6 (13.9-17.3)	113,600
	25-34	12.2 (10.0-14.4)	52,600	13.9 (12.1-15.7)	61,800	13.0 (11.6-14.5)	114,500
	35-44	10.8 (9.1-12.6)	48,200	15.0 (13.3-16.8)	68,600	12.9 (11.7-14.1)	116,800
	45-54	10.7 (9.1-12.3)	42,300	14.2 (12.6-15.8)	57,000	12.5 (11.3-13.6)	99,700
	55-64	9.7 (7.9-11.4)	28,400	11.8 (10.2-13.4)	33,700	10.7 (9.5-11.8)	62,100
	65-74	6.1 (4.8-7.5)	12,200	8.4 (7.1-9.6)	18,100	7.3 (6.4-8.2)	30,500
	75+	7.0 (5.0-9.0)	8,700	9.2 (7.6-10.9)	16,100	8.2 (7.0-9.5)	24,800

**Note:** Estimates are based on 414 respondents in South Eastern Sydney and 34,985 in NSW. For this indicator 8 (1.93%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 384 (1.1%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those with a Kessler 10 (K10) score of 22 or above. The K10 is a 10-item questionnaire about the level of anxiety and depressive symptoms in the most recent 4-week period. K10 scores for respondents aged 65 years and over were derived using 6 questions from the K10 questionnaire.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Difficulties getting health care when needing it by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	7.9 (3.2-12.6)	4,600	8.4 (4.4-12.4)	5,000	8.1 (5.0-11.2)	9,600
	16-24	9.8 (0.0-23.1)	1,000	3.1 (0.0-7.7)	400	6.2 (0.0-12.8)	1,400
	25-34	. (-.)	.	8.6 (0.0-18.9)	1,000	4.2 (0.0-9.3)	1,000
	35-44	10.4 (0.0-24.4)	1,200	14.5 (0.8-28.2)	1,500	12.3 (2.4-22.1)	2,700
	45-54	15.9 (0.6-31.2)	1,300	15.8 (2.5-29.1)	1,400	15.9 (5.8-25.9)	2,700
	55-64	10.2 (0.0-23.8)	800	8.7 (0.0-17.3)	600	9.5 (1.3-17.7)	1,400
	65-74	6.2 (0.0-14.6)	300	. (-.)	.	3.3 (0.0-7.9)	300
	75+	. (-.)	.	3.1 (0.0-7.3)	200	2.0 (0.0-4.8)	200
NSW	All Ages	11.6 (10.8-12.3)	253,200	14.6 (13.9-15.3)	333,000	13.1 (12.6-13.6)	587,100
	16-24	5.2 (3.8-6.7)	18,800	9.9 (8.1-11.7)	34,200	7.5 (6.3-8.6)	53,000
	25-34	12.8 (10.6-15.0)	53,300	16.0 (14.2-17.8)	70,000	14.4 (13.0-15.8)	123,300
	35-44	13.6 (11.7-15.5)	59,500	19.5 (17.6-21.4)	87,700	16.6 (15.2-17.9)	147,200
	45-54	14.3 (12.4-16.1)	53,900	16.9 (15.2-18.6)	66,700	15.6 (14.4-16.9)	121,000
	55-64	11.8 (10.2-13.5)	33,400	14.1 (12.6-15.6)	38,600	12.9 (11.8-14.0)	72,100
	65-74	13.2 (11.4-15.0)	25,300	11.3 (9.8-12.7)	23,300	12.2 (11.0-13.3)	48,800
	75+	7.2 (5.5-9.0)	8,900	7.5 (6.1-9.0)	12,600	7.4 (6.3-8.5)	21,700

**Note:** Estimates are based on 403 respondents in South Eastern Sydney and 33,854 in NSW. For this indicator 19 (4.71%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 1515 (4.48%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who had difficulties getting health care when they needed it. It excludes those who said they do not need health care. The question used to define the indicator was 'Do you have any difficulties getting health care when you need it?'

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Emergency department care rated as excellent, very good or good by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	75.1 (47.4-100)	5,300	59.3 (37.4-81.1)	5,100	66.4 (48.8-84.0)	10,400
	16-24	62.6 (0.7-100)	1,100	. (-.)	.	27.8 (0.0-66.6)	1,100
	25-34	100 (100-100)	1,200	62.5 (0.0-100)	800	81.0 (44.5-100)	2,000
	35-44	100 (100-100)	900	100 (100-100)	700	100 (100-100)	1,700
	45-54	21.6 (0.0-60.0)	300	79.3 (49.5-100)	1,700	56.5 (19.5-93.6)	2,000
	55-64	100 (100-100)	600	77.3 (38.3-100)	700	86.6 (61.1-100)	1,300
	65-74	100 (100-100)	800	57.0 (0.0-100)	300	83.5 (53.1-100)	1,000
	75+	100 (100-100)	400	100 (100-100)	900	100 (100-100)	1,200
NSW	All Ages	78.7 (76.1-81.4)	256,800	76.6 (74.2-78.9)	241,900	77.7 (75.9-79.4)	500,200
	16-24	74.1 (67.2-81.1)	48,600	67.1 (59.8-74.4)	36,100	70.8 (65.8-75.8)	85,000
	25-34	73.6 (66.6-80.7)	46,600	69.9 (63.9-75.9)	49,100	71.7 (67.1-76.3)	95,600
	35-44	77.2 (70.8-83.5)	44,800	78.0 (72.7-83.2)	43,100	77.6 (73.5-81.7)	88,100
	45-54	74.7 (67.3-82.1)	35,000	76.3 (70.6-81.9)	33,600	75.4 (70.7-80.0)	68,800
	55-64	86.0 (80.2-91.7)	32,700	81.6 (76.5-86.7)	28,000	84.0 (80.1-87.8)	61,000
	65-74	90.6 (86.7-94.4)	29,800	87.0 (82.5-91.5)	25,100	89.0 (86.0-91.9)	55,100
	75+	89.7 (84.5-94.9)	19,400	91.7 (88.2-95.2)	27,000	90.9 (87.9-93.9)	46,600

**Note:** Estimates are based on 52 respondents in South Eastern Sydney and 5,672 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 45 (0.79%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those attending an emergency department in the last 12 months who rated the care as excellent, very good, or good for their most recent visit. The question used to define the indicator was 'In the last 12 months, have you attended a hospital emergency department or casualty for your own medical care?' and 'Overall, what do you think of the care you received at this emergency department? Was it excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Hospital care rated as excellent, very good or good by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	85.5 (63.0-100)	6,600	91.2 (79.0-100)	10,500	88.9 (77.2-100)	17,100
	16-24	. (-.-)	.	58.4 (1.1-100)	1,000	52.2 (1.2-100)	1,000
	25-34	100 (100-100)	1,800	100 (100-100)	2,600	100 (100-100)	4,400
	35-44	100 (100-100)	1,400	100 (100-100)	2,900	100 (100-100)	4,200
	45-54	26.1 (0.0-73.1)	300	92.9 (78.3-100)	1,500	63.5 (17.8-100)	1,800
	55-64	100 (100-100)	800	82.7 (52.0-100)	1,000	89.7 (70.1-100)	1,800
	65-74	100 (100-100)	1,000	100 (100-100)	400	100 (100-100)	1,300
	75+	100 (100-100)	1,300	100 (100-100)	1,200	100 (100-100)	2,500
NSW	All Ages	92.4 (90.6-94.2)	247,800	89.9 (88.3-91.6)	329,500	91.0 (89.8-92.2)	579,300
	16-24	90.4 (83.9-96.9)	29,200	84.7 (78.1-91.3)	35,400	87.2 (82.5-91.9)	64,700
	25-34	90.5 (85.2-95.9)	33,400	89.7 (85.9-93.4)	90,000	89.9 (86.8-93.0)	123,400
	35-44	87.1 (79.3-94.9)	26,700	89.1 (84.8-93.3)	54,100	88.5 (84.6-92.3)	81,200
	45-54	90.3 (84.6-95.9)	38,700	91.1 (87.4-94.8)	42,400	90.7 (87.4-94.0)	81,400
	55-64	95.5 (93.0-98.0)	40,800	88.8 (84.7-92.9)	33,500	92.4 (90.0-94.8)	74,800
	65-74	94.8 (92.1-97.4)	39,900	92.4 (89.5-95.3)	33,700	93.5 (91.5-95.5)	73,800
	75+	96.2 (93.5-99.0)	39,100	94.4 (92.1-96.7)	40,500	95.3 (93.5-97.1)	80,000

**Note:** Estimates are based on 69 respondents in South Eastern Sydney and 5,451 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 9 (0.17%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those admitted to hospital in the last 12 months who rated the care as excellent, very good, or good for their most recent overnight stay. The questions used to define the indicator were 'In the last 12 months, have you stayed for at least 1 night in hospital?' and 'Overall, what do you think of the care you received at this hospital? Was it excellent, very good, good, fair, or poor'.

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Attended a community event at least once in the last 6 months by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	40.7 (30.4-50.9)	19,800	50.9 (42.6-59.2)	23,400	45.6 (38.9-52.3)	43,200
	16-24	27.0 (4.8-49.3)	2,300	48.7 (28.6-68.7)	5,000	38.8 (22.7-55.0)	7,300
	25-34	35.7 (13.7-57.7)	4,400	57.8 (35.4-80.2)	4,500	44.3 (28.2-60.5)	8,900
	35-44	55.3 (30.6-80.0)	4,300	60.9 (40.2-81.6)	4,800	58.1 (42.0-74.2)	9,100
	45-54	51.8 (26.6-77.0)	4,500	60.9 (41.4-80.4)	4,200	55.8 (39.3-72.3)	8,700
	55-64	35.2 (10.2-60.1)	1,900	36.5 (19.0-53.9)	1,800	35.8 (20.3-51.3)	3,700
	65-74	40.8 (15.5-66.0)	1,500	40.6 (18.0-63.2)	1,700	40.7 (23.8-57.5)	3,200
	75+	39.6 (2.6-76.6)	900	34.7 (16.3-53.1)	1,400	36.5 (18.6-54.3)	2,200
NSW	All Ages	53.7 (52.3-55.1)	860,100	61.5 (60.4-62.6)	1,017,900	57.5 (56.6-58.4)	1,881,900
	16-24	50.1 (46.3-54.0)	131,100	59.0 (55.4-62.6)	152,500	54.4 (51.7-57.0)	284,300
	25-34	51.7 (47.9-55.4)	160,500	61.0 (58.1-63.9)	193,200	56.3 (53.9-58.6)	353,800
	35-44	65.1 (61.9-68.4)	208,300	72.8 (70.3-75.3)	236,700	68.9 (66.8-70.9)	445,800
	45-54	57.5 (54.4-60.7)	163,000	63.6 (61.0-66.2)	180,700	60.4 (58.4-62.5)	344,200
	55-64	48.4 (45.3-51.5)	97,900	58.7 (56.1-61.3)	116,100	53.4 (51.4-55.4)	214,900
	65-74	45.9 (42.8-49.0)	64,300	53.9 (51.1-56.8)	82,000	50.0 (47.9-52.1)	147,000
	75+	41.7 (37.4-45.9)	34,900	46.9 (43.9-50.0)	56,700	44.5 (42.0-47.0)	92,000

**Note:** Estimates are based on 337 respondents in South Eastern Sydney and 25,602 in NSW. For this indicator 0 (0%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 44 (0.17%) were 'not stated' (Don't know or Refused) in NSW.

The indicator includes those who have attended at least one community event in the last six months. The question used was "In the past six months, how often have you attended a local community event such as a church or school fete, school concert, or a street fair?".

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Most people can be trusted by age




Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	74.5 (66.1-83.0)	35,000	67.6 (59.8-75.4)	30,400	71.1 (65.3-77.0)	65,400
	16-24	83.3 (64.6-100)	6,600	87.3 (75.0-99.6)	9,000	85.6 (74.9-96.2)	15,600
	25-34	77.9 (61.3-94.5)	9,300	50.5 (27.2-73.9)	3,900	67.1 (52.5-81.6)	13,200
	35-44	78.6 (59.1-98.1)	6,100	76.7 (58.3-95.2)	6,000	77.7 (64.2-91.1)	12,200
	45-54	83.8 (65.8-100)	6,900	56.5 (36.4-76.6)	3,900	71.4 (57.2-85.6)	10,800
	55-64	43.2 (17.5-68.9)	2,300	62.6 (44.0-81.2)	2,800	52.0 (35.1-69.0)	5,000
	65-74	60.3 (35.0-85.5)	2,300	66.1 (43.0-89.2)	2,600	63.3 (46.1-80.4)	4,900
	75+	75.1 (45.9-100)	1,600	57.3 (37.4-77.3)	2,100	63.7 (46.3-81.1)	3,700
NSW	All Ages	70.2 (68.9-71.5)	1,101,100	65.3 (64.1-66.4)	1,045,900	67.6 (66.8-68.5)	2,154,500
	16-24	65.8 (62.2-69.5)	168,400	57.3 (53.7-60.9)	144,600	61.5 (58.9-64.1)	314,400
	25-34	66.6 (63.0-70.1)	202,900	60.7 (57.8-63.7)	186,100	63.6 (61.3-65.9)	389,400
	35-44	69.6 (66.4-72.7)	218,200	66.3 (63.6-69.0)	210,300	67.9 (65.8-70.0)	430,000
	45-54	74.5 (71.7-77.3)	208,300	69.3 (66.8-71.9)	192,000	71.9 (70.0-73.8)	401,100
	55-64	71.8 (69.0-74.7)	143,200	67.0 (64.5-69.6)	128,100	69.3 (67.4-71.3)	272,200
	65-74	72.6 (69.8-75.5)	98,900	68.5 (65.7-71.4)	99,900	70.5 (68.5-72.5)	200,000
	75+	76.5 (72.7-80.3)	61,200	75.1 (72.4-77.9)	84,800	75.8 (73.5-78.0)	147,400

**Note:** Estimates are based on 323 respondents in South Eastern Sydney and 24,837 in NSW. For this indicator 14 (4.33%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 809 (3.26%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who strongly agree, or agree that most people can be trusted. The question used was "Most people can be trusted. Do you strongly agree, agree, disagree or strongly disagree?".

# Report of the New South Wales Population Health Survey

## South Eastern Sydney Division of General Practice Visit neighbours by age



Division	Age (years)	Males (95% CI)	Males (est. no.)	Females (95% CI)	Females (est. no.)	Persons (95% CI)	Persons (est. no.)
South Eastern Sydney	All Ages	51.6 (41.3-61.9)	25,100	52.7 (44.2-61.1)	24,000	52.1 (45.4-58.8)	49,100
	16-24	78.0 (53.1-100)	6,700	48.0 (28.2-67.9)	5,000	61.6 (44.9-78.4)	11,600
	25-34	53.3 (31.9-74.7)	6,500	35.7 (14.5-56.9)	2,800	46.4 (30.7-62.2)	9,300
	35-44	48.9 (23.7-74.1)	3,800	82.2 (67.8-96.6)	6,500	65.7 (49.8-81.5)	10,300
	45-54	18.1 (2.3-34.0)	1,600	44.6 (24.6-64.5)	3,100	29.9 (16.4-43.3)	4,600
	55-64	54.3 (29.0-79.5)	3,000	67.2 (50.2-84.2)	3,400	60.4 (45.0-75.9)	6,300
	65-74	50.2 (25.3-75.2)	1,900	50.6 (27.3-73.8)	2,100	50.4 (33.4-67.4)	4,000
NSW	75+	75.6 (50.0-100)	1,700	35.8 (17.9-53.6)	1,300	50.7 (32.7-68.8)	3,000
	All Ages	68.1 (66.7-69.4)	1,089,900	63.8 (62.6-64.9)	1,054,800	65.9 (65.0-66.7)	2,153,700
	16-24	74.6 (71.3-77.9)	195,200	64.0 (60.5-67.5)	165,500	69.3 (66.9-71.7)	363,100
	25-34	66.8 (63.3-70.3)	207,900	63.6 (60.7-66.4)	201,400	65.2 (62.9-67.4)	410,100
	35-44	65.9 (62.6-69.3)	210,400	66.7 (64.1-69.4)	216,600	66.4 (64.3-68.5)	428,800
	45-54	66.5 (63.4-69.6)	188,500	61.1 (58.4-63.8)	173,500	63.7 (61.7-65.8)	362,800
	55-64	68.0 (65.0-71.0)	137,200	63.5 (60.9-66.1)	125,400	65.7 (63.8-67.7)	263,900
	65-74	67.2 (64.2-70.3)	94,200	65.4 (62.7-68.2)	99,600	66.2 (64.1-68.3)	194,600
75+	67.6 (63.5-71.6)	56,600	60.2 (57.2-63.3)	72,800	63.1 (60.7-65.6)	130,400	

**Note:** Estimates are based on 336 respondents in South Eastern Sydney and 25,605 in NSW. For this indicator 1 (0.3%) were 'not stated' (Don't know or Refused) in South Eastern Sydney and 41 (0.16%) were 'not stated' (Don't know or Refused) in NSW. The indicator includes those who visited someone in their neighbourhood at least once in the past week. The question used was "How often have you visited someone in your neighbourhood in the past week?"