

Influenza Surveillance Weekly Report

Week 19: 7 to 13 May 2018


Key Points

- ▶ Influenza activity continued to be low across all NSW local health districts
- ▶ Influenza A was circulating at higher levels than influenza B

Activity compared to the previous week – NSW local health districts

Local Health District	Confirmed Influenza		NSW Emergency Departments (60) All Respiratory/Fever/Unspecified infections		
	Cases	Trend ¹	Presentations	Trend ¹	% of LHD ED presentations ²
Central Coast	1	▶	293	▶	11%
Far West	2	▶	42	▶	7%
Hunter New England	5	▶	724	▲	12%
Illawarra Shoalhaven	3	▶	242	▶	10%
Mid North Coast	2	▶	244	▶	13%
Murrumbidgee	0	▶	228	▶	12%
Nepean Blue Mountains	2	▶	255	▲	14%
Northern NSW	4	▶	232	▶	11%
Northern Sydney	12	▶	459	▲	11%
South Eastern Sydney	9	▶	665	▶	12%
South Western Sydney	5	▶	685	▶	13%
Southern NSW	1	▶	70	▶	11%
Sydney	6	▶	375	▶	12%
Western NSW	1	▶	172	▶	11%
Western Sydney	9	▶	712	▲	16%
New South Wales	62	▶	5402	▶	12%

Confirmed influenza by NSW local health district and local area (SA2)³



Summary for this reporting week:

- ▶ [Hospital surveillance](#) – ILI presentations to EDs well below the seasonal threshold
- ▶ [Laboratory surveillance](#) – low influenza laboratory test positive rate (1.3%)
- ▶ [Community surveillance](#) – low influenza activity across all LHDs
- ▶ [National surveillance](#) – low influenza activity nationally

Hospital Surveillance

NSW emergency department (ED) presentations for respiratory illness

Source: PHREDSS⁴

For the week ending 13 May 2018:

- The daily index of increase for *influenza-like illness* (ILI)⁵ presentations across NSW was low at 1.8 and remains well below the seasonal threshold of 15.
- Presentations for *All respiratory illness, fever and unspecified infections* increased but remained low and were within the usual range for this time of year (Figure 1, Table 1). Some of the increase is due to *bronchiolitis* presentations which increased within the usual range for this time of year (Figure 4 and Table 1), and *asthma* presentations (Table 1).
- The proportion of *All respiratory illness, fever and unspecified infections* presentations to all unplanned ED presentations was low at 12.4 per 100 presentations and steady (Figure 2).
- There were increases in presentations for *All respiratory illness, fever and unspecified infections* this week in the Nepean Blue Mountains, Hunter New England, Northern Sydney and Western Sydney LHDs. The stable trend in influenza notifications in these areas suggests this is unrelated to influenza activity. There is no indication of any sustained increase in respiratory illness activity in the coming weeks.
- ILI presentations resulting in admission increased and were slightly above the usual range (Figure 3, Table 1).
- ED presentations for *pneumonia*⁶ were steady and within the usual range, as were admissions for *pneumonia* (Table 1).
- *Pneumonia* and ILI presentations requiring admission to critical care increased but were within the usual range for this time of year (Table 1).

Figure 1: Total weekly counts of ED visits for *All respiratory illness, fever and unspecified infections*, all ages, from 1 January – 13 May, 2018 (black line), compared with the 5 previous years (coloured lines).




Figure 2: Total weekly counts of ED visits for *All respiratory illness, fever and unspecified infections*, all ages, as a rate per 100 ED visits, from 1 January – 13 May, 2018 (black line), compared with the range of season rate curves for the 5 previous years (white zone) aligned to the PHREDSS season start in 2017 (week 26).




Figure 3: Total weekly counts of ED visits for *influenza-like-illness* that were admitted, all ages, from 1 January – 13 May, 2018 (black line), compared with the 5 previous years (coloured lines).




Figure 4 Total weekly counts of ED visits for *bronchiolitis*, all ages, from 1 January – 13 May, 2018 (black line), compared with the 5 previous years (coloured lines).




Table 1 Weekly emergency department respiratory illness summary, week ending 13 May 2018.

Data source	Diagnosis or problem category	Trend since last week	Comparison with usual range*	Significantly elevated age groups	Significant elevated severity indicators**	Comment
ED presentations 60 NSW hospitals	Influenza-like illness (ILI)	Steady (31)	Within (26–48)			The NSW daily index of increase for ILI presentations was 1.8.
	ILI admissions	Increased (8)	Above (1-6)			
	Pneumonia	Steady (388)	Within (329–470)			
	Pneumonia admissions	Steady (289)	Within (255–313)			
	Pneumonia and ILI critical care admissions	Increased (23)	Within (19–34)			
	Asthma	Increased (524)	Within (505–688)			
	Bronchiolitis	Increased (284)	Within (243–332)			Bronchiolitis is a disease of infants.
	All respiratory illness, fever and unspecified infections	Increased (5,385)	Within (4,786–5,695)			
Ambulance	Breathing problems	Decreased (1,880)	Within (1,502–1,901)	65+ years (1059)		

Notes:*The usual range is the range of weekly counts for the same week in the previous five years for ED presentations and for ambulance Triple (000) calls.

Key for trend since last week: Non-bold and green=decreased or steady; Non-bold and orange=increased

Key for comparison with usual range: Non-bold and green =usual range; Non-bold and orange=above usual range, but not significantly above five-year mean; **Bold** and yellow=within usual range, but significantly above five-year mean; **Bold** and red = above the usual range and significantly above five-year mean (ED).

Counts are statistically significant (shown in **bold**) if they are at least five standard deviations above the five-year mean. The 'daily index of increase' is statistically significant above a threshold of 15. LHD = Local Health District.

**Severity indicators include: Admission or admission to a critical care ward (CCW); Triage category 1; Ambulance arrival and Death in ED.

FluCAN (The Influenza Complications Alert Network)

In 2009, the [FluCAN](#) surveillance system was created to be a rapid alert system for severe respiratory illness requiring hospitalisation. Data is provided on patients admitted with influenza confirmed by polymerase chain reaction (PCR) testing.


In NSW, three hospitals participate in providing weekly FluCAN data: Westmead Hospital, John Hunter Hospital and the Children's Hospital at Westmead.

During week 19 there was one influenza B admission in a NSW sentinel hospital (Figure 5).

Since 1 April 2018, there have been six hospital admissions reported for influenza; four due to influenza A and two due to influenza B (Figure 5).

Of these admissions, four were paediatric cases (<16 years of age) and two were adults. None of the cases have been admitted to a critical care ward.

Figure 5: FluCAN – Confirmed influenza hospital admissions in NSW, 1 April – 13 May, 2018.



Laboratory Surveillance

For the week ending 13 May 2018 the number and proportion of respiratory specimens reported by NSW sentinel laboratories ⁷ which tested positive for influenza A or influenza B was low (Table 2, Figure 5).

Overall, 1.3% of tests for respiratory viruses were positive for influenza (Figure 5), the same as the previous week. Influenza A strains remained more common than B strains (Figure 6).

Rhinovirus and respiratory syncytial virus (RSV) were the most common respiratory viruses identified (Table 2).

Table 2: Summary of testing for influenza and other respiratory viruses at NSW laboratories, 1 January to 13 May 2018.

Month ending	Total Tests	TEST RESULTS															
		Influenza A						Influenza B		Adeno	Parainf 1, 2 & 3	RSV	Rhino	HMPV **	Entero		
		Total		H3N2		H1N1 pdm09		A (Not typed)								Total	
Total	(%)	Total	(%A)	Total	(%A)	Total	(%A)	Total	(%)								
28/01/2018	12819	483	(3.8%)	26	(5.4%)	38	(7.9%)	414	(85.7%)	507	(4.0%)	404	599	492	1601	325	196
25/02/2018	14540	531	(3.7%)	46	(8.7%)	35	(6.6%)	448	(84.4%)	503	(3.5%)	374	552	846	2498	221	284
01/04/2018*	22518	524	(2.3%)	52	(9.9%)	49	(9.4%)	423	(80.7%)	424	(1.9%)	703	1057	2022	4775	306	485
29/04/2018	19888	247	(1.2%)	22	(8.9%)	35	(14.2%)	190	(76.9%)	147	(0.7%)	640	869	2669	3634	277	415
Week ending																	
6/05/2018	5278	47	(0.9%)	4	(8.5%)	2	(4.3%)	41	(87.2%)	21	(0.4%)	174	222	691	902	72	95
13/05/2018	5666	50	(0.9%)	5	(10.0%)	3	(6.0%)	38	(76.0%)	22	(0.4%)	155	212	780	1114	71	109

Notes: * Five-week reporting period. ** Human metapneumovirus

Figure 6: Weekly influenza positive test results by type and sub-type reported by NSW sentinel laboratories, 1 January to 13 May 2018





Figure 7: Percentage of laboratory tests positive for influenza A and influenza B by week, 1 January 2013 to 13 May 2018, New South Wales.



Influenza notifications by Local Health District (LHD)

In the week ending 13 May there were 62 notifications of influenza confirmed by polymerase chain reaction (PCR) testing, lower than the 71 notifications in the previous week.

Influenza notification rates were low and stable across all NSW LHDs (Table 3).

Table 3: Weekly notifications of laboratory-confirmed influenza by Local Health District.

Local Health District	Week ending 13 May 2018		Week ending 6 May 2018	
	Number of notifications	Rate per 100 000 population	Number of notifications	Rate per 100 000 population
Central Coast	1	0.29	0	0
Far West	2	6.55	0	0
Hunter New England	5	0.53	4	0.43
Illawarra Shoalhaven	3	0.73	5	1.21
Mid North Coast	2	0.89	3	1.33
Murrumbidgee	0	0	1	0.41
Nepean Blue Mountains	2	0.51	6	1.54
Northern NSW	4	1.29	1	0.32
Northern Sydney	12	1.3	13	1.41
South Eastern Sydney	9	0.96	14	1.49
South Western Sydney	5	0.5	7	0.69
Southern NSW	1	0.46	2	0.92
Sydney	6	0.9	9	1.35
Western NSW	1	0.36	4	1.43
Western Sydney	9	0.9	10	1

Notes: * All data are preliminary and may change as more notifications are received. Excludes notifications based on serology. For further information see the [influenza notifications data page](#).

Influenza outbreaks in institutions

There were no respiratory outbreaks reported this week.

In the year to date there have been 11 laboratory confirmed influenza outbreaks in institutions reported to NSW public health units, including ten in residential care facilities (Table 3, Figure 8). Six have been due to influenza A and five were due to influenza B.


In the ten influenza outbreaks affecting residential care facilities, at least 70 residents were reported to have had ILI symptoms and 13 required hospitalisation. Overall, there have been two deaths in residents reported which were linked to these outbreaks, both of whom were noted to have other significant co-morbidities.

Table 3: Reported influenza outbreaks in NSW institutions, January 2011 to May 2018.

Year	2011	2012	2013	2014	2015	2016	2017	2018*
No. of outbreaks	4	39	12	120	103	279	588	11

Notes: * Year to date.

Figure 8: Reported influenza outbreaks in NSW residential care facilities by month, 2014 to 2018.




The Australian Sentinel Practices Research Network (ASPREN)

ASPREN is a network of sentinel general practitioners (GPs) run through the Royal Australian College of General Practitioners and the University of Adelaide which has collected de-identified information on influenza-like illness (ILI) and other conditions seen in general practice since 1991.

Participating GPs in the program report on the proportion of patients presenting with an ILI. The number of GPs participating on a weekly basis may vary.

In week 19 there were ASPREN reports received from 78 NSW GPs. The reported consultation rate for ILI per 1000 consultations was low at 2.82 (Figure 9). For further information see the [ASPREN website](#).

Figure 9: ASPREN – NSW and National GP ILI rates per 1000 consultations – 2018 to week 19, compared to 2017.



FluTracking.net


FluTracking.net is an online health surveillance system to detect epidemics of influenza. It is a project of the University of Newcastle, the Hunter New England Local Health District and the Hunter Medical Research Institute.

Participants complete a simple online weekly survey which is used to generate data on the rate of ILI symptoms in communities.

In week 19 FluTracking received reports for 11,271 people in NSW with the following results:

- 1.5% of respondents reported fever and cough, slightly higher than the previous week and below the five year annual mean (1.3%) (Figure 10).
- Among respondents who reported being vaccinated for influenza in 2018, 1.4% reported fever and cough compared to the 1.6% rate reported among unvaccinated respondents (Figure 10).
- 0.9% of respondents reported fever, cough and absence from normal duties, similar to the previous week (0.8%).

Figure 10: FluTracking – Percent of NSW participants reporting fever and cough by vaccination status.



Notes: Participants are not considered vaccinated until at least two weeks has elapsed since their recorded time of vaccination.

For further information on the project and how to participate please see the [FluTracking](#) website.

National and International Influenza Surveillance

National Influenza Surveillance

Although national influenza surveillance reports are not produced at this time of year, many jurisdictions are reporting low influenza activity. Total national reports of laboratory-confirmed influenza in April were similar or lower than 2017.

For further information on the National Notifiable Disease Surveillance System, which includes laboratory-confirmed influenza reports, see: <http://www9.health.gov.au/cda/source/cda-index.cfm>.

Global Influenza Update

The latest [WHO global update on 14 May 2018](#) provides data up to 29 April. Influenza activity returned to inter-seasonal levels in most of the countries in the temperate zone of the northern hemisphere except for some countries in Eastern Europe. Activity increased in some countries in tropical America. In the temperate zone of the southern hemisphere, influenza activity increased but remained below the seasonal thresholds. Worldwide, seasonal influenza subtypes A and B accounted for approximately the same proportion of influenza detections. Follow the link for the [WHO influenza surveillance reports](#).

Influenza at the human-animal interface

WHO publishes regular updated risk assessments of human infections with avian and other non-seasonal influenza viruses at [Influenza at the human-animal interface](#), with the most recent report published on 2 March 2018. These reports provide information on human cases of infection with non-seasonal influenza viruses, such as H5 and H7 clade viruses, and outbreaks among animals.

Since the previous update, new human infections with avian influenza A(H7N4), A(H7N9) and A(H9N2) viruses were reported. The overall risk assessment for these viruses remains unchanged. Whenever avian influenza viruses are circulating in poultry, sporadic infections and small clusters of human cases are possible in people exposed to infected poultry or contaminated environments, therefore sporadic human cases would not be unexpected.

Other sources of information on avian influenza and the risk of human infection include:

- US CDC [Avian influenza](#)
- European CDC (ECDC) [Avian influenza](#)
- Public Health Agency of Canada [Avian influenza H7N9](#).

Composition of 2018 Australian influenza vaccines

The WHO Consultation on the Composition of Influenza Vaccines for the 2018 Southern Hemisphere was held in Melbourne on 25-27 September 2017.

Following the Consultation, WHO announced its recommendations for the composition of quadrivalent vaccines for use in the 2018 Southern Hemisphere influenza season, which includes changes in the influenza A(H3N2) component, as follows:

- an A/Michigan/45/2015 (H1N1)pdm09-like virus
- an A/Singapore/INF1HM-16-0019/2016 (H3N2)-like virus⁸
- a B/Phuket/3073/2013-like virus (Yamagata lineage)
- a B/Brisbane/60/2008-like virus (Victoria lineage).⁹

The WHO consultation on the composition of influenza vaccines for the Northern Hemisphere 2018-19 influenza season was held in February 2018. WHO announced its recommendations for the composition of quadrivalent vaccines for use in the 2018-19 Northern Hemisphere influenza season, which includes changes in the influenza A(H3N2) and influenza B (Victoria lineage) components.

More details about the most recent influenza vaccine recommendations can be found at: <http://www.who.int/influenza/vaccines/virus/en/>.

Report Notes:

¹ Notes for trend comparisons with the previous week:

		Trend in Cases	Trend in Presentations
▶	Stable	<10% change or <20 cases change	<10% change or <40 presentations change
▼	Decrease	10% or greater decrease	10% or greater decrease
▲	Increase	10-20% increase	10-20% increase
▲	Higher increase	>20% increase	>20% increase

² *All Respiratory, fever and unspecified infections* presentations as a percentage of all unplanned emergency department presentations in participating hospitals in the local health district.

³ NSW Local Health Districts and SA2: Influenza notification maps use NSW Local Health District Boundaries and Australian Bureau of Statistics (ABS) statistical area level 2 (SA2) of place of residence of cases are shown. Note that place of residence is used as a surrogate for place of acquisition for cases; the infection may have been acquired while the person was in another area.

⁴ NSW Health Public Health Rapid, Emergency Disease and Syndromic Surveillance system, CEE, NSW Ministry of Health. Comparisons are made with data for the preceding 5 years. Includes unplanned presentations to 60 NSW emergency departments, which accounted for 83% of all NSW ED presentations in the 2016/2017 financial year. The coverage is lower in rural EDs. Data is continuously updated.

⁵ The ED 'ILI' syndrome includes provisional diagnoses selected by a clinician of 'influenza-like illness' or 'influenza' (including 'pneumonia with influenza'), avian and other new influenza viruses.

⁶ The ED 'Pneumonia' syndrome includes provisional diagnoses selected by a clinician of 'viral, bacterial, atypical or unspecified pneumonia', 'SARS', or 'legionnaire's disease'. It excludes the diagnosis 'pneumonia with influenza'.

⁷ Preliminary laboratory data is provided by participating sentinel laboratories on a weekly basis and are subject to change. Point-of-care test results have been included since August 2012 but serological diagnoses are not included. Participating sentinel laboratories: Pathology North (Hunter, Royal North Shore Hospital), Pathology West (Nepean, Westmead), South Eastern Area Laboratory Services, Sydney South West Pathology Service (Liverpool, Royal Prince Alfred Hospital), The Children's Hospital at Westmead, Australian Clinical Labs, Douglas Hanly Moir Pathology, Laverty Pathology, Medlab, SydPath, VDRLab

⁸ This replaces A/Hong Kong/4801/2014 (H3N2)-like virus used in the 2017 seasonal influenza vaccines.

⁹ This B/Brisbane strain had been part of the WHO recommendations for 2017 southern hemisphere trivalent influenza vaccines but has been replaced by the B/Phuket strain for 2018 trivalent vaccines.