

# Influenza Surveillance Weekly Report

**Week 22: 27 May to 2 June 2019**


## Key Points

- ▶ Influenza activity continued to be high across all NSW local health districts, consistent with the annual influenza season.
- ▶ Respiratory presentations to NSW emergency departments decreased slightly but remained high in most districts, and were within the usual range for influenza seasons overall.
- ▶ Influenza A strains predominated but influenza B strain activity is also increasing.

## Activity compared to the previous week – NSW local health districts

Local Health District	Confirmed Influenza Notifications		NSW Emergency Departments (67) All Respiratory/Fever/Unspecified infections		
	Cases	Trend <sup>1</sup>	Presentations	Trend <sup>1</sup>	% of LHD ED presentations <sup>2</sup>
Central Coast	57	▶	408	▶	15%
Far West	11	▶	40	▶	10%
Hunter New England	97	▼	904	▼	14%
Illawarra Shoalhaven	87	▲	399	▶	14%
Mid North Coast	31	▲	300	▼	15%
Murrumbidgee	86	▶	342	▶	17%
Nepean Blue Mountains	135	▲	304	▶	15%
Northern NSW	53	▲	255	▶	12%
Northern Sydney	408	▲	569	▼	13%
South Eastern Sydney	311	▲	931	▶	16%
South Western Sydney	220	▶	974	▶	17%
Southern NSW	32	▶	265	▶	16%
Sydney	199	▲	505	▶	16%
Western NSW	102	▲	355	▶	14%
Western Sydney	516	▲	1032	▶	20%
New South Wales	2345	▲	7582	▶	15%

## Confirmed influenza by NSW local health district and local area (SA2)<sup>3</sup>



## Summary for this reporting week:

- ▶ [Hospital surveillance](#) – ILI presentations to EDs remain on an increasing trend
- ▶ [Laboratory surveillance](#) – the influenza laboratory test positive rate was only slightly higher (16.9%). Influenza A strains predominated but B strains are increasing
- ▶ [Community surveillance](#) – influenza activity increased across the majority of LHDs and was above the usual range across all LHDs
- ▶ [Death surveillance](#) – 49 influenza-related deaths have been reported to date in 2019. People who die with influenza may have other underlying illnesses, and surveillance captures only a proportion of people who die from influenza
- ▶ [National surveillance](#) – high influenza activity for this time of year

## Hospital Surveillance


### NSW emergency department (ED) presentations for respiratory illness

Source: PHREDSS<sup>4</sup>


For the week ending 2 June 2019:

- Presentations for *All respiratory illness, fever and unspecified infections* decreased this week but remain above the usual range for this time of year (Figure 1, Table 1). The proportion of these presentations to all unplanned ED presentations increased slightly and was higher than the previous week at 15.4 per 100 presentations and within the seasonal range (Figure 2).
- Presentations were significantly elevated across all ages and in the majority of NSW local health districts (LHD) (Table 1).
- The daily index of increase for *influenza-like illness* (ILI)<sup>5</sup> presentations across NSW increased further to 47.9 (up from 43.2 last week). The seasonal threshold of 15 was exceeded on 21 April (Week 16), marking the start of the PHREDSS ILI season.
- ILI presentations resulting in admission increased and remained above the usual range for this time of year (Figure 3, Table 1). Admissions were elevated across all age groups.
- ED presentations and admissions for pneumonia both decreased, however both remain above the usual range for this time of year (Table 1).
- *Pneumonia* and ILI presentations requiring admission to critical care remained steady but were above the usual range for this time of year (Figure 4, Table 1).
- ED presentations for *Bronchiolitis* decreased and were within the usual range for this time of year (Table 1).


**Figure 1:** Total weekly counts of ED visits for *All respiratory illness, fever and unspecified infections*, all ages, 1 January – 2 June 2019 (black line), compared with the 5 previous years (coloured lines).




**Figure 2:** Total weekly counts of ED visits for *All respiratory illness, fever and unspecified infections*, all ages, as a rate per 100 ED visits, 1 January – 2 June 2019 (black line), compared with the range of season rate curves for the 5 previous years (white zone) aligned to the PHREDSS season start in 2019 (week 16).



**Figure 3:** Total weekly counts of ED visits for *influenza-like-illness* that were admitted, all ages, 1 January – 2 June 2019 (black line), compared with the 5 previous years (coloured lines).



**Figure 4:** Total weekly counts of ED presentations for influenza-like illness and pneumonia, *that were admitted to a critical care ward*, all ages, 1 January – 2 June 2019 (black line), compared with the 5 previous years (coloured lines).



**Table 1:** Weekly emergency department respiratory illness summary, week ending 2 June 2019.

Data source	Diagnosis or problem category	Trend since last week	Comparison with usual range*	Significantly elevated age groups	Significant elevated severity indicators**	Comment
ED presentations 60 NSW hospitals	Influenza-like illness (ILI)	Increased (309)	<b>Above (27–52)</b>	<b>5-16 years (60)</b> <b>17-34 years (99)</b> <b>65+ years (47)</b> <b>0-4 years (32)</b> <b>35-64 years (71)</b>	<b>Ambulance arrival (44)</b>	The NSW daily index of increase for ILI presentations was 47.9.
	ILI admissions	Increased (72)	<b>Above (4–10)</b>	<b>5-16 years (8)</b> <b>65+ years (29)</b> <b>17-34 years (19)</b> <b>0-4 years (7)</b> <b>35-64 years (9)</b>	<b>Critical Care (4)</b> <b>Ambulance arrival (29)</b>	
	Pneumonia	Decreased (583)	Above (434–532)	<b>5-16 years (60)</b>		
	Pneumonia admissions	Decreased (381)	Above (321–377)			
	Pneumonia and ILI critical care admissions	Steady (36)	Above (21–35)			
	Asthma	Decreased (580)	Within (568–664)			
	Bronchiolitis	Decreased (320)	Within (271–370)			Bronchiolitis is a disease of infants.
	All respiratory illness, fever and unspecified infections	Decreased (7,564)	<b>Above (5,744–6,227)</b>	<b>5-16 years (1,144)</b> <b>17-34 years (1,013)</b> <b>65+ years (1,597)</b> <b>35-64 years (1,215)</b> <b>0-4 years (2,595)</b>	<b>Admission (2,601)</b> <b>Ambulance arrival (1,636)</b>	
Ambulance	Breathing problems	Decreased (2,299)	<b>Above (1,629–1,943)</b>	<b>65+ years (1,297)</b>		

Notes: \*The usual range is the range of weekly counts for the same week in the previous five years for ED presentations and for ambulance Triple (000) calls.

Key for trend since last week: Non-bold and green=decreased or steady; Non-bold and orange=increased

Key for comparison with usual range: Non-bold and green =usual range; Non-bold and orange=above usual range, but not significantly above five-year mean; Bold and yellow=within usual range, but significantly above five-year mean; Bold and red = above the usual range and significantly above five-year mean (ED).

Counts are statistically significant (shown in bold) if they are at least five standard deviations above the five-year mean.

The 'daily index of increase' is statistically significant above a threshold of 15. LHD = Local Health District.

\*\*Severity indicators include: Admission or admission to a critical care ward (CCW); Triage category 1; Ambulance arrival and Death in ED.

## FluCAN (The Influenza Complications Alert Network)


In 2009, the FluCAN surveillance system was created to be a rapid alert system for severe respiratory illness requiring hospitalisation. Data is provided on patients admitted with influenza confirmed by polymerase chain reaction (PCR) testing.

In NSW, three hospitals participate in providing weekly FluCAN data: Westmead Hospital, John Hunter Hospital and the Children's Hospital at Westmead.

During week 22 there were 10 influenza admissions to NSW sentinel hospitals (Figure 6).

Since 1 April 2019, there have been 71 hospital admissions reported for influenza; 48 due to influenza A (including 12 A(H1N1) and 12 A(H3)) and 23 due to influenza B (Figure 6). Of these admissions, 64 were paediatric cases (<16 years of age) and seven were in adults. No cases have been admitted to a critical care ward.

**Figure 6:** FluCAN – Confirmed influenza hospital admissions in NSW, 1 April – 2 June, 2019\*.



**Note:** \* Admissions data are subject to change as new information is received. Westmead Hospital data is not available so far for 2019.

## Laboratory Surveillance

For the week ending 2 June 2019 the number and proportion of respiratory specimens reported by NSW sentinel laboratories<sup>6</sup> which tested positive for influenza A or influenza B increased only slightly but remained higher than expected for this time of year (Table 2, Figure 7).

Overall, 16.9% of tests for respiratory viruses were positive for influenza (Figure 7), higher than the previous week (16.3%).

Influenza A strains remained more common than B strains, with influenza A(H3N2) strains now more common than A(H1N1) strains (Table 2, Figures 7-8).


Rhinovirus remained the most common respiratory virus identified, followed by influenza then respiratory syncytial virus (RSV), which is a common cause of bronchiolitis in infants (Table 2).

**Table 2:** Summary of testing for influenza and other respiratory viruses at NSW laboratories, 1 January to 2 June 2019.


Month ending	Total Tests	TEST RESULTS															
		Influenza A								Influenza B		Adeno	Parainf 1, 2 & 3	RSV	Rhino	HMPV **	Entero
		Total		H3N2		H1N1 pdm09		A (Not typed)		Total							
		Total	(%)	Total	(%A)	Total	(%A)	Total	(%A)	Total	(%)	Total	Total	Total	Total	Total	Total
3/02/2019*	23496	2055	(8.7%)	111	(5.4%)	161	(7.8%)	1777	(86.5%)	129	(0.5%)	730	902	920	3171	270	485
3/03/2019*	25351	2232	(8.8%)	144	(6.5%)	134	(6.0%)	1954	(87.5%)	145	(0.6%)	710	926	1448	5053	162	693
31/03/2019	31863	2664	(8.4%)	132	(5.0%)	198	(7.4%)	2334	(87.6%)	302	(0.9%)	967	1408	2583	5866	172	843
28/04/2019	34720	2957	(8.5%)	144	(4.9%)	158	(5.3%)	2652	(89.7%)	491	(1.4%)	1003	1422	3799	7148	208	1109
28/04/2019	61942	6303	(10.2%)	262	(4.2%)	112	(1.8%)	5929	(94.1%)	2270	(3.7%)	1528	1337	4695	11729	312	1206
<b>Week ending</b>																	
5/05/2019	9906	700	(7.1%)	49	(7.0%)	24	(3.4%)	627	(89.6%)	208	(2.1%)	274	261	899	1531	51	225
12/05/2019	10336	815	(7.9%)	36	(4.4%)	17	(2.1%)	762	(93.5%)	297	(2.9%)	279	271	828	1703	54	232
19/05/2019	11786	1176	(10.0%)	52	(4.4%)	26	(2.2%)	1098	(93.4%)	401	(3.4%)	295	242	891	2179	40	243
26/05/2019	13885	1651	(11.9%)	51	(3.1%)	25	(1.5%)	1575	(95.4%)	612	(4.4%)	336	281	1018	2948	76	245
2/06/2019	16029	1961	(12.2%)	74	(3.8%)	20	(1.0%)	1867	(95.2%)	752	(4.7%)	344	282	1059	3368	91	261

Notes: \* Five-week reporting period. \*\* Human metapneumovirus

**Figure 7:** Weekly influenza positive test results by type and sub-type reported by NSW sentinel laboratories, 1 January to 2 June 2019.



**Figure 8:** Percentage of laboratory tests positive for influenza A and influenza B by week, 1 January 2014 to 2 June 2019, New South Wales.



### Influenza notifications by Local Health District (LHD)

In the week ending 2 June there were 2345 notifications of influenza confirmed by polymerase chain reaction (PCR) testing, higher than the 1934 (revised) notifications reported in the previous week. Rates were higher than usual for this time of year.

Influenza notification rates increased across the majority of the State, with the exception of Hunter New England and Southern NSW LHDs. Rates were higher than usual in all jurisdictions for this time of year. Rates were highest in Western Sydney, Northern Sydney, Far West and Western NSW and Nepean Blue Mountains LHDs (Table 3).

**Table 3:** Weekly notifications of laboratory-confirmed influenza by local health district.

Local Health District	Week ending 02 Jun 2019		Week ending 26 May 2019	
	Number of notifications	Rate per 100 000 population	Number of notifications	Rate per 100 000 population
Central Coast	57	16.36	58	16.64
Far West	11	36.59	9	29.94
Hunter New England	97	10.29	125	13.26
Illawarra Shoalhaven	87	20.91	60	14.42
Mid North Coast	31	13.88	13	5.82
Murrumbidgee	86	28.97	83	27.96
Nepean Blue Mountains	135	35.05	104	27
Northern NSW	53	17.27	23	7.49
Northern Sydney	408	43.15	347	36.7
South Eastern Sydney	311	32.81	217	22.89
South Western Sydney	220	21.57	216	21.18
Southern NSW	32	14.94	35	16.35
Sydney	199	28.98	152	22.14
Western NSW	102	35.96	77	27.15
Western Sydney	516	50.23	415	40.4

Notes: \* All data are preliminary and may change as more notifications are received. For further information see the [influenza notifications data page](#).

### Influenza outbreaks in institutions

There were six influenza outbreaks in institutions reported this week. One outbreak was in a private hospital, with the rest in residential care facilities. All were due to influenza A.

In the year to date there have been 62 laboratory confirmed influenza outbreaks in institutions reported to NSW public health units, including 50 in residential care facilities (Table 4, Figure 9). Fifty-nine of the outbreaks have been due to influenza A, two were due to influenza B and one involved both A and B strains.

In the 50 influenza outbreaks affecting residential care facilities, at least 481 residents were reported to have had ILI symptoms and 53 required hospitalisation. Overall, there have been 14 deaths<sup>1</sup> in residents reported which were linked to these outbreaks, all of whom were noted to have other significant co-morbidities.

<sup>1</sup> Deaths associated with institutional outbreaks are also included in the [Deaths surveillance](#) section if laboratory-confirmed.


NSW public health units advise institutions on how to manage their influenza outbreaks. NSW Health also provides influenza antiviral treatment to help control outbreaks when requested and appropriate. NSW Health have provided 763 courses of antivirals so far this year.

**Table 4:** Reported influenza outbreaks in NSW residential care facilities, January 2014 to 2 June 2019.

Year	2014	2015	2016	2017	2018	2019*
Number of outbreaks	121	103	252	543	42	50

Note: \* Year to date.

**Figure 9:** Reported influenza outbreaks in NSW residential care facilities by month, 2014 to 2 June 2019.



## The Australian Sentinel Practices Research Network (ASPREN)


ASPREN is a network of sentinel general practitioners (GPs) run through the Royal Australian College of General Practitioners and the University of Adelaide which has collected de-identified information on influenza-like illness (ILI) and other conditions seen in general practice since 1991.

Participating GPs in the program report on the proportion of patients presenting with an ILI. The number of GPs participating on a weekly basis may vary.

In week 22 there were ASPREN reports received from 72 NSW GPs. The reported consultation rate for ILI per 1000 consultations was increased at 8.7 (Figure 10), lower than the previous week (12.82), but higher than usual for this time of year.

For further information see the [ASPREN website](#).

**Figure 10:** ASPREN – NSW and National GP ILI rates per 1000 consultations – 2019 to the week ending 2 June, compared to 2018 weekly rates.



## FluTracking.net

FluTracking.net is an online health surveillance system to detect epidemics of influenza. It is a project of the University of Newcastle, the Hunter New England Local Health District and the Hunter Medical Research Institute. Participants complete a simple online weekly survey which is used to generate data on the rate of ILI symptoms in communities.


In week 22 FluTracking received reports for 13,803 people in NSW with the following results:

- 2.0% of respondents reported fever and cough, lower than the previous week (2.1%) and similar to the five year annual mean (2.1%) (Figure 11).
- Among respondents who reported being vaccinated for influenza in 2019, 1.9% reported fever and cough lower than the 2.1% rate reported among unvaccinated respondents (Figure 11).
- 1.3% of respondents reported fever, cough and absence from normal duties, lower than the previous week (1.4%).



the 25-34 years age-group, one death in a person in the 45-54 years age-group, with the remaining deaths in people aged 55 years or older. An additional 1400 (7.71%) death certificates mentioned pneumonia.

**Figure 12:** Rate of deaths classified as *influenza or pneumonia* per 100 000 NSW population, 2014 – 10 May, 2019



Source: NSW Registry of Births, Deaths and Marriages.

\* Notes on interpreting death data:

- (a) Deaths registration data is routinely reviewed for deaths attributed to pneumonia or influenza. While pneumonia has many causes, a well-known indicator of seasonal and pandemic influenza activity is an increase in the number of death certificates that mention pneumonia or influenza as a cause of death.
- (b) The predicted seasonal baseline estimates the predicted rate of pneumonia or influenza deaths in the absence of influenza epidemics. If deaths exceed the epidemic threshold, then it may be an indication that influenza is beginning to circulate widely and may be more severe.
- (c) The number of deaths mentioning “Pneumonia or influenza” is reported as a rate per 100,000 NSW population (rather than a rate per total deaths reported).
- (d) Deaths referred to a coroner during the reporting period may not be available for analysis, particularly deaths in younger people which are more likely to require a coronial inquest. Influenza-related deaths in younger people may be under-represented in these data as a result.
- (e) The interval between death and death data availability is usually at least 14 days, and so these data are at least two weeks behind reports from emergency departments and laboratories and subject to change.

### Influenza-related deaths with laboratory confirmation

For the year to 2 June 2019 there have been 49 laboratory-confirmed influenza deaths (Table 5). This includes six people who died in this reporting week. All of the newly reported deaths were in people aged 60 years or older.

In 2019, 15 of the deaths occurred in May, 12 were in April, 3 were in March, 10 were in February and 9 were in January.

These influenza death data come from all sources, including public health units, aged care facility outbreaks, the NSW Coroner’s Office, and the NSW Registry of Births, Deaths and Marriages, but

are only included here where there has been laboratory confirmation of influenza infection for that person. Data are subject to change as new information is received.

**Table 5:** Laboratory-confirmed influenza deaths by age-group and year, NSW, 2017 to 2 June 2019 (by date of death).

Age-group	Year		
	2017	2018	2019*
0-4 years	2	2	0
5-19 years	4	0	0
20-64 years	44	6	8
65+ years	509	32	41
<b>Total</b>	<b>559</b>	<b>40</b>	<b>49</b>

Notes: \*Year to date.

## Government-funded vaccine distribution

---

NSW Health commenced distributing National Immunisation Program and NSW Government Program influenza vaccines on 1 April 2019.

National Immunisation Program (NIP) vaccines include vaccines for people aged 65 years and over, pregnant women, Aboriginal people aged 6 months and over, and people 6 months and over with medical conditions pre-disposing them to severe influenza.

NSW Government Program vaccines are for health care workers in NSW Health facilities and all children from 6 months to under 5 years of age not covered under the NIP.

As of 2 June, 2.2 million doses had been distributed to general practitioners, Aboriginal medical services, hospitals, aged care facilities, and childhood vaccination clinics across NSW.

For more information about the 2019 Influenza Vaccination Program see:

<https://www.health.nsw.gov.au/immunisation/Pages/flu.aspx> .

## National and International Influenza Surveillance

---

### National Influenza Surveillance

The fortnightly *Australian Surveillance Report No.2*, with data up to 19 May 2019, noted:

- **Activity** –Currently, influenza and influenza-like illness (ILI) activity are high for this time of year compared to previous years. At the national level, notifications of laboratory-confirmed influenza have increased in the past fortnight and it is likely these figures will be revised upwards due to backlogs in data entry. Influenza A was the most common respiratory virus detected in patients presenting with ILI to sentinel general practices this fortnight.
- **Severity** –There is no indication of the potential severity of the 2019 season at this time.
- **Impact** – There is no indication of the potential impact on society of the 2019 season at this time.
- **Virology** – In the year to date and in the past fortnight, the majority of confirmed influenza cases reported nationally were influenza A (87%). Where subtyping data were available, influenza A(H3N2) was the dominant influenza A subtype in the past fortnight.

For further information see the [Australian Influenza Surveillance Reports](#).

### Global Influenza Update

The latest [WHO global update on 27 May 2019](#) provides data up to 12 May 2019. In the temperate zones of the southern hemisphere, influenza detections increased overall. In summary:

- In Australia and New Zealand influenza detections were predominantly influenza A(H3N2) and influenza B viruses.
- In South Africa, predominantly influenza A(H3N2) viruses were detected.
- In South America, influenza A(H1N1)pdm09 viruses predominated.
- In Southern Asia, influenza activity was low overall.
- In the Caribbean, Central American countries, and the tropical countries of South America, influenza and RSV activity were low in general.
- In Eastern, West and Middle Africa, influenza activity was low across reporting countries.
- In North America and Europe, influenza activity was low overall.
- In North Africa, influenza detections were low across reporting countries.
- In Western Asia, influenza activity was low overall, but with continued detections in a few countries on the Arabian Peninsula.
- In East Asia, decreased but continued influenza activity was reported.

Worldwide, seasonal influenza A viruses accounted for the majority of detections.

Follow the link for the [WHO influenza surveillance reports](#).

### **Influenza at the human-animal interface**

WHO publishes regular updated risk assessments of human infections with avian and other non-seasonal influenza viruses at [Influenza at the human-animal interface](#), with the most recent report published on 9 April 2019. These reports provide information on human cases of infection with non-seasonal influenza viruses, such as H5 and H7 clade viruses, and outbreaks among animals.

Since the previous update, new human infections with avian influenza A(H7N9) and A(H9N2) viruses were reported. The overall risk assessment for these viruses remains unchanged. Other sources of information on avian influenza and the risk of human infection include:

- US CDC [Avian influenza](#)
- European CDC (ECDC) [Avian influenza](#)
- Public Health Agency of Canada [Avian influenza H7N9](#).

## **Composition of influenza vaccines in 2019**

---

### **WHO influenza vaccine strain recommendations – Southern Hemisphere, 2019**

The [WHO recommendations](#) for the composition of trivalent vaccines included changes in the influenza A(H3N2) component and the influenza B (Victoria lineage), as follows:

- an A/Michigan/45/2015 (H1N1)pdm09-like virus
- an A/Switzerland/8060/2017 (H3N2)-like virus
- a B/Colorado/06/2017-like virus (B/Victoria lineage)

It was recommended that quadrivalent vaccines also contain a second B component, a B/Phuket/3073/2013-like virus (B/Yamagata lineage).

### **Australian influenza vaccine strain recommendations – 2019 influenza season**

The Australian Influenza Vaccine Committee (AIVC) recommendation for the Australian trivalent vaccine includes a B/Yamagata lineage virus (a B/Phuket/3073/2013-like virus), rather than a

B/Victoria lineage virus, based on circulating influenza B viruses at the time of the recommendation. The Therapeutic Goods Administration (TGA) accepted the [AIVC recommendations](#) for 2019.

Information on NSW seasonal influenza vaccination activities in 2019, including free vaccine for all children aged 6 months to less than 5 years can be found at:

<https://www.health.nsw.gov.au/immunisation/Pages/flu.aspx> .

### **WHO influenza vaccine strain recommendations – Northern Hemisphere, 2019-20**

The WHO Consultation on the Composition of Influenza Vaccines for Use in the 2019-20 Northern Hemisphere Influenza Season was held in Beijing on 18-20 February 2019.

From this meeting it was recommended that egg based quadrivalent vaccines for use in the 2019-2020 northern hemisphere influenza season contain the following:

- an A/Brisbane/02/2018 (H1N1)pdm09-like virus;
- an A/Kansas/14/2017 (H3N2)-like virus;
- a B/Colorado/06/2017-like virus (B/Victoria/2/87 lineage); and
- a B/Phuket/3073/2013-like virus (B/Yamagata/16/88 lineage).

It was also recommended that the influenza B virus component of trivalent vaccines for use in the 2019-2020 northern hemisphere influenza season should be a B/Colorado/06/2017-like virus of the B/Victoria/2/87-lineage.

In light of recent changes in the proportions of genetically and antigenically diverse A(H3N2) viruses, the recommendation for the A(H3N2) component was announced on 21 March. More details about the most recent influenza vaccine recommendations can be found at:

<http://www.who.int/influenza/vaccines/virus/en/> .

## Report Notes:

---

<sup>1</sup> Notes for trend comparisons with the previous week:

		Trend in Cases	Trend in Presentations
▶	Stable	<10% change or <20 cases change	<10% change or <40 presentations change
▼	Decrease	10% or greater decrease	10% or greater decrease
▲	Increase	10-20% increase	10-20% increase
▲	Higher increase	>20% increase	>20% increase

<sup>2</sup> All Respiratory, fever and unspecified infections presentations as a percentage of all unplanned emergency department presentations in participating hospitals in the local health district.

<sup>3</sup> NSW Local Health Districts and SA2: Influenza notification maps use NSW Local Health District Boundaries and Australian Bureau of Statistics (ABS) statistical area level 2 (SA2) of place of residence of cases are shown. Note that place of residence is used as a surrogate for place of acquisition for cases; the infection may have been acquired while the person was in another area.

<sup>4</sup> NSW Health Public Health Rapid, Emergency Disease and Syndromic Surveillance system, CEE, NSW Ministry of Health. Comparisons are made with data for the preceding 5 years. Includes unplanned presentations to 67 NSW emergency departments, which accounted for 83% of all NSW ED presentations in the 2016/2017 financial year. The coverage is lower in rural EDs. Data is continuously updated.

<sup>5</sup> The ED 'ILI' syndrome includes provisional diagnoses selected by a clinician of 'influenza-like illness' or 'influenza' (including 'pneumonia with influenza'), avian and other new influenza viruses.

<sup>6</sup> Preliminary laboratory data is provided by participating sentinel laboratories on a weekly basis and are subject to change. Point-of-care test results have been included since August 2012 but serological diagnoses are not included. Participating sentinel laboratories: Pathology North (Hunter, Royal North Shore Hospital), Pathology West (Nepean, Westmead), South Eastern Area Laboratory Services, Sydney South West Pathology Service (Liverpool, Royal Prince Alfred Hospital), The Children's Hospital at Westmead, Australian Clinical Labs, Douglas Hanly Moir Pathology, Laverty Pathology, Medlab, SydPath, VDRLab