



Invasive Group A Streptococcus (iGAS) infection



If someone close to you has an iGAS infection

Visit your doctor if you have any of the following symptoms in the next 30 days

Skin sores or



Sore throat or



Fever or chills



Time to visit your doctor



If you or someone you care for is really sick or gets sick very quickly, has difficulty breathing, becomes confused, or has blue or pale skin, call **Triple Zero (000)** or go to a hospital emergency department.

Infection can spread through close contact

Kissing and touching



Coughing and sneezing



Sharing clothes and towels with someone with symptoms



Sharing drinks, vapes and/or cigarettes



Staying in same bed with someone with symptoms



What can I do to prevent iGAS infections?

Wash hands...			Visit your doctor...		To stop the spread...	
after coughing and sneezing	wash hands with soap and water	before eating	to get treatment for everyone in the household with skin sores, and cover any broken skin	if you may have been in contact with someone with scabies	keep wounds clean and covered until healed	stay at home if you have symptoms or until you have completed your first day of antibiotics

For further NSW Health iGAS information for close contacts, scan the QR Code or visit www.health.nsw.gov.au/Infectious/controlguideline/Documents/igas-appendix2.pdf

For non-urgent medical assistance call Healthdirect on 1800 022 222.



Acknowledgment

Centre for Disease Control, Department of Health Northern Territory Government. www.nt.gov.au/health

August 2025 © NSW Health. SHPN (HP NSW) 241060.