

# Patients presenting to General Practitioners who are also receiving treatment with long acting depot buprenorphine



## **Long acting depot buprenorphine has been approved for the treatment of opioid dependence in Australia**

The Therapeutic Goods Administration (TGA) in Australia has approved two long-acting injected depot buprenorphine medications: Buvidal™ and Sublocade™. Both products are listed on the PBS.

### **These formulations of buprenorphine are administered weekly or monthly**

Buvidal™ is a modified release formulation of buprenorphine which is administered via subcutaneous (SC) injection in *weekly* or *monthly* intervals.

Sublocade™ is an extended-release formulation of buprenorphine which is administered via subcutaneous injection in *monthly* intervals.

### **What does this mean for GPs?**

Patients with opioid dependence may request this product for treatment of opioid dependence. Increasing numbers of patients may present to GPs who are being treated with depot buprenorphine, which may have implications for patients' other treatment needs.

### **Depot buprenorphine may reduce the effects of opioid analgesia**

Buprenorphine, through its properties as a partial agonist at opioid receptors, may reduce the effects of opioid analgesics. Adequate analgesia may be more difficult to achieve when administering opioids to patients on depot buprenorphine. Opioids can be used for analgesia but doses will need to be titrated carefully against clinical response. Consultation with Pain Services or Drug and Alcohol Specialists is recommended.

### **There are potential drug-drug interactions**

There are a number of clinically relevant drug-drug interactions (DDIs) that could occur for patients on depot buprenorphine. Some of these include:

- Interactions with CNS depressants that increase the risk of overdose such as other opioids, alcohol, gabapentinoids, antihistamines, anti-psychotics and benzodiazepines.
- Except in emergency situations, opioid antagonists should generally not be used in patients using depot buprenorphine.
- The metabolism of CYP3A4 inhibitors and inducers, anticholinergics and diuretics may be altered. This includes rifampin, carbamazepine, phenytoin, phenobarbital; macrolide antibiotics such as erythromycin, azole-antifungal agents and antiretroviral medications such as ritonavir.

### **Depot buprenorphine is released slowly and can have an extended duration of effect**

Though depot buprenorphine is administered weekly or monthly, plasma concentrations of buprenorphine slowly reduce after the injection and may remain at levels that cause at least some opioid partial agonist effects for up to 20 weeks in some people, depending on dose and duration of treatment. There may therefore be a persisting though reducing effect on analgesia requirements and DDIs for many weeks after the drug is administered.

## **How will you know if your patient is on depot buprenorphine?**

In the first week after administration of Sublocade™ a small collection may be palpated. Mostly however, neither product is noticeable on external examination. Patients have been encouraged to carry and present a wallet card explaining their depot buprenorphine treatment status. However, patients might be reluctant to disclose that they are on depot buprenorphine due to previous experiences of stigmatisation or discrimination by health workers. Respectfully explaining why you need to know about medications including depot buprenorphine, and maintaining a non-judgemental approach to their drug use, helps improve the accuracy of the history obtained.

When a patient presents who has been receiving depot buprenorphine, clinicians should, as is usual with patients on opioid replacement therapy, confirm last dosing details in order to inform safe treatment. This includes recent dosing history, the date of last dose, the dose received and when the next dose is due.

## **Can GPs prescribe depot buprenorphine?**

Yes. The previous restriction limiting this product to specialist prescribers has now been removed. From April 21 2020 Buvidal® and Sublocade® may be prescribed by OTP accredited prescribers and non-OTP accredited prescribers (GPs).

The existing caps on patient numbers remain. That is, non-accredited prescribers are permitted to induct and manage up to twenty (20) opioid dependent patients on buprenorphine medicines, which can include subcutaneous or sublingual formulations. Accredited prescribers have a patient limit of 200 patients.

## **How do I arrange for the product to be delivered and administered?**

Depot buprenorphine must never be dispensed to patients and can only be administered by a health care professional.

GPs may have the product delivered to their practice where it can be administered to the patient by a nurse or by the GP. Storage requirements apply to buprenorphine as it is an s8 medication. GPs may also arrange with a pharmacy to order, receive and deliver the product to them for administration. Details about storage regulations and how to order and arrange delivery are available on the NSW Health webpage: <https://www.health.nsw.gov.au/aod/Pages/depot-buprenorphine.aspx>.

## **More information and support is available for GPs**

### ***By telephone***

- The NSW Drug and Alcohol Specialist Advisory Service (DASAS) is a 24/7 helpline that enables health professionals to get advice from specialist alcohol and other drug medical and clinical nurse consultants on any drug and alcohol clinical issues. DASAS clinicians will provide advice on clinical issues related to depot buprenorphine.

Phone DASAS: Sydney metropolitan: (02) 9361 8006; Regional and rural NSW: 1800 023 687

### ***On-line***

- Clinical Guidelines for use of Depot Buprenorphine have been published by the Ministry of Health and are available at <https://www.health.nsw.gov.au/aod/Pages/depot-bupe-guidelines.aspx>.
- The RACGP has made available a webinar: Introduction to Depot Buprenorphine in the Treatment of Opioid Dependent Patients. This webinar, presented by Professor Nick Lintzeris and Dr Hester Wilson, explains the clinical guidance, evidence base, pharmacology, and risks associated with depot buprenorphine, and how to implement it as an Opioid Agonist Treatment (OAT) for opioid dependency. The webinar can be accessed free of charge through their website: <https://www.racgp.org.au/education/professional-development/online-learning/webinars/drugs-and-alcohol/buprenorphine-in-the-treatment-of-opioid-dependent>.