


NSW Arbovirus Surveillance & Mosquito Monitoring Program, 2018-2019

Weekly Update: 9 November 2018



Health
Pathology

creating better health & justice systems
www.pathology.health.nsw.gov.au

Contents

Summary	2
Comment.....	2
Environmental Conditions	3
Rainfall	3
Three Month Rainfall & Temperature Forecast	3
Tidal	4
MVEV Climatic Models	5
Forbes' Hypothesis.....	5
Nichol's Hypothesis.....	5
Arboviral Isolates	6
Exotic Detections	6
Human Notifications	7
Monthly RRV notifications	9
Monthly BFV notifications.....	9
Mosquito Results	10
Inland	11
Coastal	12
Sydney	13
Sentinel Chicken Flocks	14

Summary

- **Climate:** light to moderate rainfall was recorded over the last week, which encompassed the entire state. For October, rainfall was below average in the southern ranges, but above average for most of the coast.
- **Three Month Forecast:** for November 2018 to January 2019, rainfall is predicted to be below average for most of NSW, although average along the coast. Maximum and minimum temperatures are predicted to exceed the average. According to the Bureau of Meteorology (BOM) as of 7 November 2018, the El Niño Alert continues and a positive Indian Ocean Dipole is now underway. These environmental parameters suggest that rainfall patterns are likely to be below average in the upcoming months.
- **Tides:** there was some hatching of *Aedes vigilax* associated with the recent rains, furthermore, the tides over 7-10 November 2018, have been above predicted, prompting a treatment at Sydney Olympic Park. The next high tides are due over 23-28 November 2018 and are forecast to reach over 1.9m. These tides should trigger *Aedes vigilax* hatching.
- **Murray Valley Encephalitis virus (MVEV) Models:** the data relevant to both the Forbes' and Nichols' hypotheses have been updated to October 2018. Neither model is suggestive of an MVEV epidemic.
- **Mosquito Numbers Inland:** mosquito numbers were low.
- **Mosquito Numbers Coast:** surveillance activities are due to begin in December.
- **Mosquito Numbers Sydney:** surveillance activities are due to begin in December, although one trapping night this week from Georges River produced a 'medium' catch (50-100 mosquitoes).
- **Arboviral Isolates:** there are no arboviral isolates to date.
- **Chicken Sentinel Flocks:** no report has yet been issued.
- **Human Notifications:** for the current fiscal year, there have been 149 Ross River virus (RRV) and 17 Barmah Forest virus (BFV) notifications, which is slightly below the previous four season average. However, notifications at this time of the year are usually either false positives or not recent infections.


Comment: this is the first report for the 2018-2019 surveillance season. Only a few sites began trapping this week, and for the inland, mosquito numbers were 'low'. One trapping night at the Georges River in Sydney produced 'medium' numbers. However in the nearby region of Ryde, there have been a number of mosquito complaints submitted to the Council (Craig Redfern, SEHO, *pers. comm.*). The recent rains have filled empty containers around the home, triggering hatching of species such as *Aedes notoscriptus*. The rains also initiated hatching of *Aedes vigilax* on the salt marshes. Thus people living near Parramatta River received the double effect of sudden increases in salt marsh and urban mosquito species.

The BOM continues with the El Niño Alert and the Indian Ocean Dipole is positive, while the three month forecast is largely for dry and hot conditions ahead.

Environmental Conditions

Rainfall

Rainfall across Australia for the week ending 8 November 2018 is depicted on the left and monthly rainfall deciles for October 2018 are on the right. Over the last week, there was light to moderate rainfall across the entire state. For October, rainfall was below average in the southern ranges, but above average for most of the coast. Maximum temperatures and minimum temperatures for October were 1-2 degrees above average for the state.



Three Month Rainfall & Temperature Forecast

For November 2018 to January 2019, rainfall is predicted to be below average for most of NSW, although average along the coast. Maximum and minimum temperatures are predicted to exceed the average. The following webpages contain graphics of the seasonal outlook:

www.bom.gov.au/climate/outlooks/#/rainfall/median (Rainfall outlook).


www.bom.gov.au/climate/outlooks/#/temperature/summary (Max & min temperature outlook).

According to the BOM as of 7 November 2018, the El Niño Alert continues. A positive Indian Ocean Dipole (IOD) is now underway (positive IODs are associated with dry conditions across Australia). These indices suggest that rainfall patterns are likely to be below average in the upcoming months.

For more information: www.bom.gov.au/climate/enso/ and, <http://www.bom.gov.au/climate/iod/>

Tidal

Tidal information is relevant for the prediction of the activity of the salt marsh mosquito, *Aedes vigilax*. Typically for NSW, tides of over 1.8m, as measured at Sydney, can induce hatching of *Aedes vigilax* larvae and the graph below of predicted tide heights can provide some indication of when this is likely to occur. Note this trigger height varies between regions, thus at Batemans Bay, a tide height over 0.8m can initiate egg hatching.



The rains along the coast initiated some saltmarsh mosquito hatching around the Sydney area and no control was able to be undertaken. The current series of high tides are greater than predicted and treatments of mosquito habitat at Sydney Olympic Park will be occurring (C. Webb, Department of Medical Entomology, *pers. comm.*).

The next series of high tides will be during 23-28 November 2018 and these are forecast to reach over 1.9m. This is likely to trigger *Aedes vigilax* hatching.

Actual tide heights can vary by 0.3m (or more in unusual circumstances) due to variations in atmospheric pressure, rainfall, wind and other climatic phenomena. Sea level rise with climate change may also result increased tide heights. Thus predicted tide height should be used as a gauge only for potential *Aedes vigilax* activity. The larvae of the saltmarsh mosquito relies on a inundation/drying cycle for the mudflats in which it lives; continual wet weather prevents the drying cycles thereby reducing larval production.

MVEV Climatic Models

Three predictive environmental based models for MVEV activity have been developed; the Forbes (which relies on rainfall in the river catchment basins of Eastern Australia), Nichols (based on the Southern Oscillation), and the Bennett theory (based on the Indian Ocean Dipole). The latter theory has low reliability and is not considered below. Note that all the predictive models have been developed on a limited data set and do not always forecast activity. There can also be unusual environmental conditions that may lead to the introduction of the virus to southeastern Australia, such as the movement of low pressure cells from the north to the south of the country during 2008 and 2011. Vertical transmission of the virus (from adult to the egg in *Aedes* species) can result in restricted activity following localised heavy precipitation (as per 2003 at Menindee).

i. Forbes' Hypothesis

Rainfall was not above Decile 7 in all of the river catchment basins in eastern Australia for the last quarter of 2017, the first quarter of 2018, nor the last quarter of 2018 (Table 1). Thus Forbes' hypothesis for an MVEV outbreak has not been fulfilled.

Table 1. Rainfall indices for the main catchment basins of eastern Australia as per Forbes' hypothesis, relevant to the 2018-2019 season. Note that a value of 1 equals Decile 7 rainfall.

Catchment Basin	Oct-Dec 2017	Jan-Mar 2018	Oct-Dec 2018*	Jan-Mar 2019
Darling River	0.93	0.52	0.85	
Lachlan/Murrumbidgee/Murray Rivers	1.15	0.70	0.44	
Northern Rivers	0.81	1.07	0.77	
North Lake Eyre system	0.75	0.69	0.52	

*Data for October only

ii. Nichol's Hypothesis

Table 2. The seasonal atmospheric pressures (in mm) according to Nichol's hypothesis, relevant to the 2018-2019 season.

	Autumn 2018	Winter 2018	Spring 2018*
2018 Value	1009.27	1011.8	1012.30
Pre past MVEV seasons	<1009.74	<1012.99	<1009.99

*Data for October only

The Spring period pertaining to the Nichol's hypothesis is not in line with past MVEV active years.

Arboviral Isolates

LOCATION - Site	Date Trapped	Mosquito Species	Virus
Nil to date			

Human Notifications

Weekly notifications of human mosquito-borne disease infections are available from the NSW Ministry of Health, Communicable Disease Weekly Report and summarized in the Table below* (www.health.nsw.gov.au/Infectious/reports/Pages/CDWR.aspx).

Table 4. Notifications of mosquito-borne disease in NSW, 2018-2019*

Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
7-Jul-18	12	1	10	1	0	0	24
14-Jul-18	9	1	2	3	0	0	15
21-Jul-18	5	2	3	2	0	0	12
28-Jul-18	8	1	6	3	0	0	18
4-Aug-18	4	0	8	3	0	0	15
11-Aug-18	8	3	6	1	0	0	18
18-Aug-18	12	0	3	1	0	0	16
25-Aug-18	8	1	2	1	0	0	12
1-Sep-18	8	0	0	0	0	0	8
8-Sep-18	5	1	2	2	0	0	10
15-Sep-18	13	4	1	5	0	0	23
22-Sep-18	5	0	5	1	0	0	11
29-Sep-18	7	2	5	2	0	0	16
6-Oct-18	10	0	2	1	0	0	13
13-Oct-18	9	0	2	4	0	0	15
20-Oct-18	7	0	5	2	1	0	15
27-Oct-18	11	0	7	1	0	0	19
3-Nov-18							
10-Nov-18							
17-Nov-18							
24-Nov-18							
1-Dec-18							
8-Dec-18							
15-Dec-18							
22-Dec-18							
29-Dec-18							

RRV = Ross River virus; BFV = Barmah Forest virus; DENV = Dengue virus; CHIKV = Chikungunya virus; ZIKV = Zika virus.

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland.

*The data in this table is updated once available from the NSW Ministry of Health.

Comment: It should also be noted that notifications are for NSW residents and that infection may have been acquired elsewhere and that winter notifications of RRV are likely to be false positives.

Table 4 cont. Notifications of mosquito-borne disease in NSW, 2018-2019*

Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
5-Jan-19							
12-Jan-19							
19-Jan-19							
26-Jan-19							
2-Feb-19							
9-Feb-19							
16-Feb-19							
23-Feb-19							
2-Mar-19							
9-Mar-19							
16-Mar-19							
23-Mar-19							
30-Mar-19							
6-Apr-19							
13-Apr-19							
20-Apr-19							
27-Apr-19							
4-May-19							
11-May-19							
18-May-19							
25-May-19							
1-June-19							
8-June-19							
15-June-19							
22-June-19							
29-June-19							
Total	141	16	69	33	1	0	260

RRV = Ross River virus; BFV = Barmah Forest virus; DENV = Dengue virus; CHIKV = Chikungunya virus; ZIKV = Zika virus.

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland.

*The data in this table is updated once available from the NSW Ministry of Health.

Table 5. Ross River virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2013 to November 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2014-2015	38	50	46	67	59	90	117	305	431	264	102	50	1,619
2015-2016	54	61	53	61	70	54	42	60	78	79	52	16	680
2016-2017	12	11	20	17	38	216	429	274	200	142	174	89	1,622
2017-2018	29	37	52	56	37	31	30	39	51	74	96	70	602
2018 - 2019	32	41	31	41	5								150
Ave [†]	33	40	43	50	51	98	155	170	190	140	106	56	1,132

*updated 9 November 2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).

[†]Average for 2014-15 to 2017-18.

Table modified from: <http://www1.health.nsw.gov.au/IDD/#/ROSS>

Table 6. Barmah Forest virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2014 to November 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2014-2015	10	3	11	11	8	4	12	17	43	43	16	11	189
2015-2016	6	9	7	9	6	3	4	5	2	3	10	2	66
2016-2017	4	3	0	0	1	9	9	5	8	6	24	24	93
2017-2018	8	10	6	8	8	6	5	12	8	10	8	7	96
2018 - 2019	4	6	5	1	1								17
Ave [†]	7	6	6	7	6	6	8	10	15	16	15	11	113


*updated 9 November 2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).

[†]Average for 2014-15 to 2017-18.

Table modified from: <http://www1.health.nsw.gov.au/IDD/#/BF>

Mosquito Results

Mosquito abundance is best described in relative terms, and in keeping with the terminology from previous NSW Arbovirus Surveillance and Mosquito Monitoring Program Annual Reports, mosquito numbers are depicted in the tables below as:



Each location represents the average for all trapping sites at that location.

Inland

Location	Mosquito	Oct-17				Nov				Dec				Jan-18				Feb				Mar				
		7	14	21	28	4	11	18	25	2	9	16	23	30	6	13	20	27	3	10	17	24	3	10	17	24
Albury	<i>Cx. annul</i>																									
	Total Mosq.																									
Bourke	<i>Cx. annul</i>																									
	Total Mosq.																									
Griffith	<i>Cx. annul</i>																									
	Total Mosq.																									
Leeton	<i>Cx. annul</i>																									
	Total Mosq.																									
Macquarie Marshes	<i>Cx. annul</i>																									
	Total Mosq.																									
Mathoura	<i>Cx. annul</i>																									
	Total Mosq.																									
Wagga	<i>Cx. annul</i>																									
	Total Mosq.																									

Coastal

Location	Mosquito	Oct-17				Nov					Dec				Jan-18				Feb				Mar				
		7	14	21	28	4	11	18	25	2	9	16	23	30	6	13	20	27	3	10	17	24	3	10	17	24	31
Ballina	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Coffs Harbour	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Gosford	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Lake Macquarie	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Port Macquarie	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Tweed	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Wyong	<i>Ae. vigilax</i>																										
	Total Mosq.																										

Sydney

Location	Mosquito	Nov-18				Dec					Jan-19				Feb				Mar					Apr			
		4	11	18	25	2	9	16	23	30	6	13	20	27	3	10	17	24	3	10	17	24	31	7	14	21	28
Banks-town	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Blacktown	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Georges River	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Hawkes-bury	<i>Cx. annul</i>																										
	Total Mosq.																										
Hills Shire	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Penrith	<i>Ae. vigilax</i>																										
	Total Mosq.																										
Sydney Olympic Park	<i>Ae. vigilax</i>																										
	Total Mosq.																										

Sentinel Chicken Flocks

Location	Oct-17					Nov				Dec					Jan-18				Feb				Mar			
	7	14	21	28	4	11	18	25	2	9	16	23	30	6	13	20	27	3	10	17	24	3	10	17	24	31
Deniliquin																										
Dubbo																										
Forbes																										
Griffith																										
Hay																										
Leeton																										
Macquarie Marshes																										
Menindee																										
Moree																										

N= Negative for MVEV & Kunjin virus

Prepared by: Stephen Doggett, Manager, Department of Medical Entomology, NSW Health Pathology (ICPMR), Westmead Hospital NSW 2145. Email: Stephen.Doggett@health.nsw.gov.au

Please note that these results remain the property of the NSW Ministry of Health and may not be used or disseminated to unauthorized persons or organizations without permission.

SHPN: (EH) 180675