## NSW Childhood Immunisation Schedule

## **Primary and Community Health**





Diphtheria-tetanus-pertussishepatitis B-polio-Haemophilus influenzae type b



Infanrix Hexa or Vaxelis

Pneumococcal



Prevenar 13

Rotavirus



Rotarix

Meningococcal B (Aboriginal\* children only)



Bexsero

months

Diphtheria-tetanus-pertussis-hepatitis B-polio-*Haemophilus influenzae* type b



PRESCRIPTION ONLY MEDICINE

Uniform to MACO O GLOBAN

Company of the Company of t

At risk children ≥ 6 months require an additional dose of Prevenar 13 (Pneumococcal). Refer to the online Australian Immunisation Handbook for the list of conditions.

Aboriginal<sup>#</sup> children ≥ 6 months with certain medical conditions may require an additional dose of Meningococcal B. Refer to the online Australian Immunisation Handbook for the list of conditions.

Infanrix Hexa or Vaxelis

12 months

Meningococcal ACWY



Nimenrix

Pneumococcal



Prevenar 13

Measles-mumps-rubella



Priorix or MMR II

Meningococcal B (Aboriginal\* children only)



Bexsero

TRIPACEL®

months

 $\infty$ 

za vaccination is recommended for children aged 6 months to less than 5 years

Measles-mumps-rubella-varicella



Priorix-Tetra

Haemophilus influenzae type b



Act-HIB

Diphtheria-tetanus-pertussis



Infanrix **or** Tripacel

/ears

Diphtheria-tetanus-pertussis-polio



- Countries of the Coun

PRESCRIPTION ONLY MEDICINE
KEEP OUT OF REACH OF CHILDREN

One provided yearing to final

One

Quadracel or Infanrix IPV

At risk children require a dose of Pneumovax 23 (pneumococcal) at 2-12 months after Prevenar 13 or at age 4 years whichever is later.

5 years‡

ΛΙ

6 months to less than 5 years\*

\*Annual influen

Influenza vaccines

Vaxigrip Tetra



Flucelvax Quad



Vaxigrip Tetra



All children aged 6 months to less than 5 years are recommended to receive the annual influenza vaccine (NIP funded)\*.

Children aged 6 months to less than 9 years of age require 2 doses at least 4 weeks apart in their first year of influenza vaccination.