

NSW Immunisation Schedule


Visit the NSW Health website to view the current NSW Immunisation Schedule at: health.nsw.gov.au/schedule

Aboriginal Immunisation Health Workers

Aboriginal Immunisation Health Workers help Aboriginal people keep up to date with vaccines. This protects family and community from harmful diseases. Find out more at: health.nsw.gov.au/aboriginal-immunisation

Save the Date to Vaccinate

Vaccines work best when given at the right time. Find out when your baby's next vaccination is due health.nsw.gov.au/vaccinate



More information

NSW Health Website

health.nsw.gov.au/immunisation

MumBubVax

mumbubvax.org.au/Hepatitis-B-vaccine

Australian Immunisation Register

servicesaustralia.gov.au/australian-immunisation-register

NSW Antenatal Pertussis Vaccination Program

health.nsw.gov.au/protectnewborns

Details for your local aboriginal immunisation health worker:

Name:

Number:

Email:

Working days/time:

NSW Health

Important vaccines for Pregnancy



Some diseases are very dangerous to you and your baby if you are pregnant.

They can lead to miscarriage, premature birth or make your baby very sick when they are born.

Vaccines protect you and your baby from these diseases.

You pass the protection to your baby in the womb.

Your health care worker can help you get any of the vaccines you need.

If you're not sure if you're up to date with vaccines, talk to your health care worker.

Before you become pregnant

Make sure you're up to date with vaccines that protect against these diseases:

- Measles, Mumps and Rubella
- Chicken Pox (Varicella)

After vaccination, you should avoid getting pregnant for 28 days.

While you are pregnant

Hepatitis B

You will have a blood test for hepatitis B early in your pregnancy. If you have hepatitis B, your baby will need two needles within 12 hours of birth.



Whooping Cough (pertussis)

Get a whooping cough vaccine between 20 and 32 weeks of pregnancy. If you get this vaccine in pregnancy your baby will have protection when they are born.

COVID-19

COVID-19 vaccination protects you and your baby from getting sick. Make sure you're fully protected. It's safe to get a COVID-19 vaccine at any time during your pregnancy, including a COVID-19 booster shot. It is recommended that you advise your health care provider that you are pregnant (if they are not already aware) so that they can select the most appropriate COVID-19 vaccine for you.

Flu (influenza)

Get a flu vaccine any time in pregnancy. This protects you while you are pregnant and the baby when they are born.

After baby is born

It is safe for you to receive routine vaccinations immediately after birth, even if you are breast feeding. You should have the whooping cough vaccine before you leave hospital if you were not vaccinated during your pregnancy.

Your baby's first vaccination, hepatitis B, is recommended just after birth and the next scheduled vaccinations are due when your baby is 6 weeks of age.

Babies born to mothers with hepatitis B infection need a blood test 3 months after completing the course of hepatitis B vaccine to ensure they are protected. You and your doctor will be sent a reminder letter from NSW Health.

