



Safety Notice 002/17

Identification of Post-Injection Syndrome Olanzapine Pamoate Long Acting Injection

8 February 2017

- Distributed to:**
- Chief Executives
 - Directors of Clinical Governance
 - Directors of Mental Health Services

- Action required by:**
- Directors of Clinical Governance
 - Directors of Mental Health Services
 - Directors of Emergency Departments

- We recommend you also inform:**
- Drug and Therapeutic Committees
 - Directors of Pharmacy
 - Managers of Community Mental Health Services

- Expert Reference Group**
- Content reviewed by:
- Mental Health Branch
 - Chief Psychiatrist, NSW
 - Clinical Excellence Commission

Clinical Excellence Commission

Tel. 02 9269 5500
 Fax. 02 9269 5599
 Email: CEC-PatientSafety@health.nsw.gov.au

Internet Website
<http://www.health.nsw.gov.au/quality/sabs>

Intranet Website
<http://internal.health.nsw.gov.au/quality/sabs/>

Review date

February 2020

Background

Olanzapine Long Acting Injection (LAI) is an atypical antipsychotic used in the maintenance treatment of schizophrenia. A rare serious adverse event related to the use of olanzapine LAI is post-injection syndrome (PIS). It is reported to occur in 0.07% of injections¹. Non-recognition of PIS symptoms has resulted in the death of a patient.

PIS results from inadvertent intravascular injection of olanzapine, causing a range of olanzapine overdose-type symptoms. Post injection syndrome is not dose, frequency or time point specific, and the risk of occurrence exists following every administration. In most cases of PIS (84%) the initial signs and symptoms occur within the first hour after injection, but onset after 3 hours has been reported¹. Full recovery usually occurs within 24-72 hours².

The signs and symptoms of PIS include sedation (ranging from mild sedation to deep sleep and unconsciousness), and/or delirium (including confusion/confused state, disorientation, anxiety and agitation). Other symptoms include dizziness, weakness, altered speech/dysarthria, altered gait, muscle spasms, possible seizures and hypertension^{1,2}.

Higher doses and therefore a larger final volume for injection and low body mass index (BMI) may present a higher risk for PIS; however, PIS has occurred in patients who do not have these risk factors¹.

PIS has not been reported with other long acting antipsychotic injections.

References

1. Olanzapine depot injection (Zyprexa Relprevv) for schizophrenia
<http://www.nps.org.au/publications/health-professional/nps-radar/2009/december-2009/olanzapine-depot>
2. Post-injection delirium/sedation syndrome in patients with schizophrenia treated with olanzapine long-acting injection, I: analysis of cases
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2895589/>
3. Protocol for the use of Olanzapine Pamoate Long Acting Injection. Sydney Local Health District. 2013

- Suggested Actions by Local Health Districts:**
1. Distribute this Safety Notice to all relevant clinical staff
 2. Ensure that Emergency Department clinicians are aware of olanzapine LAI post-injection syndrome and its management
 3. If a patient with a mental health condition presents to an emergency department with symptoms of sedation and/or delirium, PIS is considered
 4. Ensure that staff administering olanzapine LAI are appropriately trained in its administration¹
 5. Ensure that staff administering olanzapine LAI are aware of the signs and symptoms of PIS
 6. That services administering olanzapine LAI have a local guideline in place that stipulates:
 - a. 30 minutely monitoring of the consumers alertness for 3 hours following injection¹
 - b. assessment by a medical officer prior to discharge to ensure no signs and symptoms of PIS are displayed
 - c. consumers are escorted home by a responsible person or staff member post administration
 - d. awareness by consumer/responsible person of the possible signs and symptoms of PIS and the need for urgent medical attention if they occur. An agreed management plan should be in place
 - e. That services administering olanzapine LAI have access to emergency services for treatment of PIS