

Trends in NSW residents born in the Netherlands

Indicator	Year	Males % (95% CI)	Females % (95% CI)	Persons % (95% CI)
Trends in health behaviours				
Risk alcohol drinking	1997-1998	53.6 (42.4-64.8)	34.0 (21.1-46.8)	45.3 (36.6-54.0)
	2002-2005	51.1 (38.5-63.8)	31.1 (18.9-43.4)	41.2 (32.4-50.1)
	2006-2009	30.3 (14.7-45.8)	18.6 (9.5-27.7)	25.1 (15.3-34.8)
Screening mammogram in the last 2 years (50 to 69 years)	1997-1998		77.0 (62.2-91.9)	
	2002-2005		75.2 (56.6-93.8)	
	2006-2009		76.8 (54.3-99.3)	
Pap test in the last 2 years (20 to 69 years)	1997-1998		79.1 (62.5-95.7)	
	2002-2005		70.4 (47.3-93.4)	
	2006-2009		72.8 (52.1-93.4)	
Hysterectomy (20 to 69 years)	1997-1998		25.0 (12.3-37.7)	
	2002-2005		25.9 (9.1-42.8)	
	2006-2009		35.4 (17.1-53.7)	
Vaccinated against influenza in the last 12 months (50 years and over)	1997-1998	26.9 (16.2-37.6)	47.9 (31.9-63.8)	35.7 (25.8-45.5)
	2002-2005	55.1 (41.8-68.3)	49.6 (38.0-61.2)	52.3 (43.5-61.2)
	2006-2009	46.2 (32.5-60.0)	51.5 (38.9-64.0)	48.6 (39.1-58.0)
Vaccinated against pneumococcal disease in the last 5 years (50 years and over)	2002-2005	29.9 (16.1-43.8)	25.2 (15.6-34.8)	27.5 (19.0-36.9)
	2006-2009	34.1 (21.2-47.1)	31.2 (19.6-43.8)	32.8 (24.0-41.0)
Live in homes with a smoke alarm or detector	1997-1998			64.2 (51.7-72.6)
	2002-2005			80.4 (73.2-86.9)
	2006-2009			89.7 (80.8-98.6)
Two or more serves of fruit a day	1997-1998	46.0 (35.0-56.9)	63.1 (52.5-77.8)	53.8 (45.4-62.3)
	2002-2005	56.6 (44.1-65.1)	55.0 (43.1-66.8)	55.8 (47.2-64.4)
	2006-2009	42.0 (26.5-57.5)	64.8 (52.9-76.7)	52.8 (42.7-62.8)
Five or more serves of vegetables a day	1997-1998	8.5 (2.7-14.2)	5.2 (0.0-10.8)	7.1 (3.0-11.2)
	2002-2005	3.6 (0.7-8.5)	5.3 (1.9-10.6)	5.4 (2.5-8.4)
	2006-2009	2.7 (0.0-5.8)	3.2 (4.8-21.6)	7.6 (3.2-12.0)
Three or more serves of vegetables a day	1997-1998	28.5 (18.8-38.1)	36.2 (23.8-48.6)	31.7 (24.0-39.3)
	2002-2005	44.6 (31.1-57.5)	45.7 (34.2-57.1)	45.1 (36.5-53.7)
	2006-2009	33.8 (22.7-50.9)	43.6 (31.4-55.7)	39.9 (30.4-49.5)
Usually consumes lower fat or skim milk	1997-1998	53.2 (41.9-64.4)	62.1 (49.5-74.7)	56.8 (48.2-65.3)
	2002-2005	46.4 (33.7-59.1)	55.7 (43.9-67.5)	51.1 (42.4-59.7)
	2006-2009	49.6 (34.2-64.9)	53.2 (41.2-65.3)	51.3 (41.3-61.3)
Adequate physical activity	1997-1998	60.0 (45.1-75.0)	35.3 (18.7-51.9)	50.1 (38.1-62.0)
	2002-2005	43.8 (31.1-56.5)	55.9 (43.8-68.0)	50.5 (41.4-59.6)
	2006-2009	43.3 (27.8-58.9)	49.2 (36.4-62.1)	46.1 (35.7-56.5)
Current smoking	1997-1998	20.5 (11.5-29.5)	22.7 (11.2-34.3)	21.4 (14.3-28.5)
	2002-2005	14.9 (6.0-23.7)	19.5 (8.0-31.1)	17.2 (9.8-24.6)
	2006-2009	5.2 (0.4-10.1)	14.0 (5.1-22.9)	9.1 (4.2-14.0)
Live in smoke-free households	1997-1998			75.4 (68.2-82.6)
	2002-2005			85.5 (80.4-90.7)
	2006-2009			90.1 (84.6-95.5)
Bans smoking in car	2002-2005			84.9 (76.6-93.1)
	2006-2009			91.0 (85.9-96.2)
Trends in health status				
Excellent, very good or good self-rated health status	1997-1998	82.8 (74.6-91.1)	75.4 (63.6-87.1)	79.8 (72.9-86.6)
	2002-2005	75.3 (64.7-85.9)	85.3 (78.8-91.8)	80.3 (74.0-86.6)
	2006-2009	86.2 (77.8-94.7)	80.0 (70.7-89.3)	83.4 (77.1-89.7)
Current asthma	1997-1998	4.7 (0.4-9.0)	18.6 (6.7-30.6)	10.4 (4.6-16.2)
	2002-2005	4.2 (0.0-8.7)	10.7 (4.4-17.1)	7.5 (3.6-11.4)
	2006-2009	8.0 (1.8-14.2)	7.0 (0.4-13.7)	7.5 (3.0-12.0)
Blood pressure measured in last 2 years	1997-1998	88.3 (80.7-95.8)	98.3 (96.3-100)	92.4 (87.7-97.0)
	2002-2005	89.6 (80.4-98.8)	100 (100-100)	95.3 (91.0-99.6)
	2006-2009	81.0 (56.2-100)	97.4 (92.2-100)	88.2 (73.4-100)
Ever had high blood pressure	1997-1998	25.7 (16.3-35.1)	24.9 (12.2-37.6)	25.4 (17.7-33.0)
	2002-2005	33.1 (18.6-47.7)	32.4 (19.7-45.1)	32.7 (23.1-42.3)
	2006-2009	24.8 (3.7-45.9)	55.5 (27.5-83.6)	38.4 (19.1-57.6)
Cholesterol measured in last 2 years	1997-1998	55.9 (44.8-67.0)	52.1 (38.3-66.0)	54.4 (45.7-63.1)
	2002-2005	74.4 (60.3-88.5)	72.7 (56.1-89.3)	73.4 (62.3-84.6)
	2006-2009	74.0 (48.8-99.2)	85.3 (63.4-100)	78.7 (61.2-96.3)
Ever had high cholesterol	1997-1998	31.2 (19.5-42.9)	27.0 (15.0-39.0)	29.5 (21.0-38.1)
	2002-2005	26.1 (12.6-39.5)	42.2 (28.2-56.2)	34.7 (24.8-44.6)
	2006-2009	16.9 (0.0-36.5)	68.5 (44.8-92.2)	38.8 (18.4-59.1)
Diabetes or high blood glucose	1997-1998	6.9 (1.1-12.7)	12.5 (3.0-21.9)	9.2 (4.0-14.3)
	2002-2005	18.0 (8.2-27.8)	3.7 (0.1-7.3)	10.8 (5.5-16.2)
	2006-2009	13.0 (5.0-20.9)	13.5 (5.5-21.5)	13.2 (7.5-18.9)
High and very high psychological distress	1997-1998	4.6 (0.6-8.6)	15.3 (6.6-23.9)	9.0 (4.7-13.3)
	2002-2005	5.3 (0.2-10.3)	14.7 (2.0-27.4)	10.5 (2.9-18.1)
	2006-2009	3.6 (0.0-8.9)	5.2 (0.9-9.5)	4.3 (0.8-7.8)
Visited a dental professional within the last 12 months	2002-2005	47.9 (34.9-61.0)	52.5 (40.3-64.7)	50.2 (41.3-59.2)
	2006-2009	63.8 (47.3-80.3)	66.4 (53.4-79.4)	64.9 (54.1-75.8)
All natural teeth missing	1997-1998	24.2 (11.2-37.2)	23.5 (7.4-39.6)	23.9 (13.8-34.0)
	2002-2005	16.9 (7.8-25.9)	17.5 (10.3-24.7)	17.2 (11.4-22.9)
	2006-2009	6.7 (0.3-13.0)	12.6 (4.4-20.7)	9.3 (4.2-14.4)

Indicator	Year	Males % (95% CI)	Females % (95% CI)	Persons % (95% CI)
Overweight	1997-1998	43.6 (32.4-54.8)	27.5 (15.8-39.3)	37.1 (28.6-45.5)
	2002-2005	43.9 (30.9-56.9)	40.2 (27.5-52.8)	41.9 (32.8-51.0)
	2006-2009	35.6 (20.3-50.8)	29.1 (18.5-39.7)	32.7 (23.0-42.3)
Obese	1997-1998	16.9 (8.1-25.7)	10.5 (4.1-16.8)	14.3 (8.4-20.2)
	2002-2005	19.1 (8.3-30.0)	13.3 (5.8-20.9)	16.0 (9.5-22.5)
	2006-2009	14.4 (6.2-22.5)	28.3 (17.0-39.5)	20.7 (13.5-27.8)
Overweight or obese	1997-1998	60.5 (49.7-71.3)	38.0 (25.1-50.9)	51.4 (42.6-60.2)
	2002-2005	63.0 (50.6-75.5)	53.5 (40.3-66.7)	57.8 (48.5-67.1)
	2006-2009	49.9 (34.2-65.7)	57.4 (45.0-69.8)	53.3 (43.0-63.6)
Trends in health services				
Private health insurance	1997-1998	49.0 (38.0-60.0)	31.6 (19.7-43.5)	41.9 (33.5-50.3)
	2002-2005	56.3 (44.1-68.5)	50.2 (38.5-61.8)	53.2 (44.7-61.7)
	2006-2009	61.0 (48.5-73.6)	39.2 (28.3-50.0)	51.1 (42.2-60.1)
Difficulties getting health care when needing it	1997-1998	9.1 (3.8-14.5)	15.7 (6.9-24.6)	11.8 (7.0-16.7)
	2002-2005	17.6 (8.8-26.5)	18.6 (6.9-30.4)	18.1 (10.8-25.5)
	2006-2009	9.8 (3.3-16.2)	13.4 (6.3-20.6)	11.4 (6.6-16.2)
Emergency department presentation in the last 12 months	1997-1998	9.5 (4.1-14.8)	9.8 (3.9-15.8)	9.6 (5.6-13.6)
	2002-2005	9.1 (1.6-16.7)	8.4 (3.1-13.6)	8.7 (4.2-13.3)
	2006-2009	11.6 (4.7-18.5)	24.0 (14.8-33.1)	17.2 (11.4-23.0)
Hospital admission in the last 12 months	1997-1998	14.1 (7.3-20.9)	10.2 (3.7-16.7)	12.5 (7.6-17.3)
	2002-2005	12.7 (3.5-21.8)	16.2 (5.1-27.3)	14.4 (7.2-21.7)
	2006-2009	13.3 (6.6-20.0)	20.3 (11.7-28.9)	16.5 (11.0-21.9)
Visited a general practice in the last 12 months	1997-1998	80.8 (72.3-89.4)	89.5 (81.2-97.8)	84.4 (76.3-90.5)
	2006-2009	82.8 (70.1-95.5)	86.1 (77.8-94.3)	84.2 (76.2-92.3)
	Trends in social capital			
Most people can be trusted	2002-2005	68.7 (56.7-80.7)	67.0 (54.1-79.9)	67.7 (58.8-76.7)
	2006-2009	89.7 (81.0-98.5)	63.6 (51.4-75.9)	75.9 (67.6-84.2)
	2002-2005	68.4 (56.1-80.7)	45.8 (32.9-58.7)	56.0 (46.7-65.3)
Feel safe walking down their street after dark	2006-2009	83.7 (73.9-93.6)	61.3 (49.3-73.2)	72.0 (63.8-80.3)
	2002-2005	73.0 (60.7-85.4)	71.6 (60.8-82.4)	72.3 (64.2-80.4)
	2006-2009	81.1 (65.9-96.2)	85.6 (77.1-94.0)	83.4 (74.7-92.0)
Visited neighbours in the last week	2002-2005	62.9 (50.8-75.1)	68.8 (58.2-79.4)	66.2 (58.1-74.2)
	2006-2009	72.3 (58.0-86.5)	54.7 (42.2-67.1)	63.1 (53.4-72.7)
	2002-2005	82.9 (73.5-92.2)	82.6 (73.5-91.7)	82.7 (76.2-89.3)
Ran into friends and acquaintances when shopping in local area	2006-2009	70.7 (56.9-84.6)	87.8 (80.7-94.8)	79.6 (71.9-87.3)
	2002-2005	72.8 (60.5-85.1)	80.0 (70.5-89.5)	76.8 (69.1-84.5)
	2006-2009	77.7 (64.9-90.4)	84.3 (74.6-93.9)	81.1 (73.2-89.0)
Would feel sad to leave their neighbourhood	2002-2005	72.8 (60.5-85.1)	80.0 (70.5-89.5)	76.8 (69.1-84.5)
	2006-2009	77.7 (64.9-90.4)	84.3 (74.6-93.9)	81.1 (73.2-89.0)

Note: Indicators include adults 16 years and over unless specified.

Source: New South Wales Population Health Survey 2006-2009 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

WARNING: Estimates Not for Publication
Please check HealthStats NSW for more information

Trends in NSW residents born in Australia

Indicator	Year	Males % (95% CI)	Females % (95% CI)	Persons % (95% CI)
Trends in health behaviours				
Risk alcohol drinking	1997-1998	55.5 (54.3-56.8)	39.5 (38.5-40.6)	47.2 (46.4-48.1)
	2002-2005	44.0 (42.8-45.2)	33.0 (32.1-33.9)	38.4 (37.6-39.1)
	2006-2009	44.7 (43.3-46.1)	29.4 (28.4-30.4)	36.8 (35.9-37.7)
Screening mammogram in the last 2 years (50 to 69 years)	1997-1998		77.4 (75.6-79.1)	
	2002-2005		76.2 (74.0-78.3)	
	2006-2009		78.6 (76.6-80.7)	
Pap test in the last 2 years (20 to 69 years)	1997-1998		78.9 (77.3-80.4)	
	2002-2005		74.9 (73.3-76.4)	
	2006-2009		75.5 (73.7-77.2)	
Hysterectomy (20 to 69 years)	1997-1998		13.6 (12.9-14.3)	
	2002-2005		12.4 (11.6-13.3)	
	2006-2009		12.7 (11.7-13.6)	
Vaccinated against influenza in the last 12 months (50 years and over)	1997-1998	36.7 (34.9-38.5)	42.0 (40.5-43.6)	39.6 (38.4-40.8)
	2002-2005	47.3 (45.8-48.9)	53.7 (52.5-54.9)	50.8 (49.8-51.7)
	2006-2009	45.3 (43.8-46.8)	53.0 (51.8-54.2)	49.4 (48.4-50.4)
Vaccinated against pneumococcal disease in the last 5 years (50 years and over)	2002-2005	22.9 (21.7-24.1)	28.9 (27.9-30.0)	26.1 (25.3-26.9)
	2006-2009	29.7 (28.4-31.1)	36.1 (34.9-37.2)	33.2 (32.3-34.1)
Live in homes with a smoke alarm or detector	1997-1998			64.7 (63.9-65.5)
	2002-2005			75.9 (75.1-76.6)
	2006-2009			93.0 (92.5-93.4)
Two or more serves of fruit a day	1997-1998	37.6 (36.4-38.8)	50.3 (49.2-51.3)	44.1 (43.3-44.9)
	2002-2005	41.7 (40.5-42.9)	54.1 (53.1-55.0)	48.1 (47.3-48.8)
	2006-2009	49.4 (48.0-50.8)	59.4 (58.3-60.4)	54.6 (53.7-55.4)
Five or more serves of vegetables a day	1997-1998	8.0 (7.4-8.6)	9.9 (9.3-10.6)	9.0 (8.6-9.4)
	2002-2005	6.6 (6.1-7.2)	10.9 (10.4-11.5)	8.8 (8.4-9.2)
	2006-2009	7.4 (6.8-8.1)	14.3 (13.6-15.0)	11.0 (10.5-11.5)
Three or more serves of vegetables a day	1997-1998	30.1 (29.0-31.2)	41.5 (40.4-42.5)	36.0 (35.2-36.7)
	2002-2005	32.1 (31.1-33.2)	48.3 (47.4-49.3)	40.4 (39.7-41.2)
	2006-2009	35.2 (33.9-36.5)	54.2 (53.1-55.3)	45.0 (44.1-45.8)
Usually consumes lower fat or skim milk	1997-1998	38.3 (37.1-39.5)	54.2 (53.2-55.3)	46.5 (45.7-47.3)
	2002-2005	37.7 (36.6-38.9)	53.1 (52.2-54.1)	45.6 (44.9-46.4)
	2006-2009	41.9 (40.6-43.3)	55.8 (54.7-56.9)	49.1 (48.2-50.0)
Adequate physical activity	1997-1998	53.9 (52.1-55.6)	43.6 (42.1-45.1)	48.5 (47.4-49.7)
	2002-2005	53.6 (52.3-54.8)	43.5 (42.4-44.5)	48.4 (47.5-49.2)
	2006-2009	62.5 (61.1-63.9)	50.4 (49.3-51.5)	56.3 (55.4-57.2)
Current smoking	1997-1998	26.7 (25.6-27.8)	22.8 (21.9-23.6)	24.7 (24.0-25.4)
	2002-2005	24.1 (23.1-25.1)	20.4 (19.6-21.2)	22.2 (21.6-22.8)
	2006-2009	20.2 (19.0-21.3)	16.9 (16.1-17.8)	18.5 (17.8-19.2)
Live in smoke-free households	1997-1998			70.9 (70.2-71.7)
	2002-2005			83.1 (82.6-83.7)
	2006-2009			89.2 (88.7-89.8)
Bans smoking in car	2002-2005			82.6 (81.9-83.3)
	2006-2009			88.5 (87.9-89.1)
Trends in health status				
Excellent, very good, or good self-rated health status	1997-1998	85.8 (85.1-86.6)	85.0 (84.3-85.7)	85.4 (84.9-85.9)
	2002-2005	81.6 (80.7-82.5)	80.0 (79.3-80.7)	80.8 (80.2-81.3)
	2006-2009	82.8 (81.9-83.7)	78.9 (78.1-79.7)	80.8 (80.2-81.4)
Current asthma	1997-1998	10.0 (9.2-10.7)	12.9 (12.2-13.7)	11.5 (11.0-12.0)
	2002-2005	12.5 (11.7-13.3)	16.5 (15.7-17.2)	14.5 (14.0-15.1)
	2006-2009	10.2 (9.3-11.0)	13.8 (13.1-14.6)	12.1 (11.5-12.6)
Blood pressure measured in last 2 years	1997-1998	82.2 (81.3-83.2)	91.5 (90.9-92.2)	87.0 (86.4-87.6)
	2002-2005	84.2 (82.9-85.4)	91.3 (90.5-92.1)	87.9 (87.1-88.6)
	2006-2009	84.9 (82.5-87.3)	92.4 (91.2-93.7)	88.8 (87.4-90.1)
Ever had high blood pressure	1997-1998	17.0 (16.2-17.9)	17.1 (16.4-17.9)	17.1 (16.5-17.7)
	2002-2005	23.2 (22.0-24.4)	24.2 (23.2-25.2)	23.7 (22.9-24.5)
	2006-2009	29.5 (27.0-31.9)	33.4 (31.5-35.3)	31.5 (30.0-33.0)
Cholesterol measured in last 2 years	1997-1998	46.6 (45.4-47.8)	44.2 (43.2-45.3)	45.4 (44.6-46.2)
	2002-2005	53.6 (51.9-55.2)	53.7 (52.3-55.0)	53.6 (52.6-54.7)
	2006-2009	59.6 (56.7-62.6)	61.8 (59.5-64.0)	60.7 (58.9-62.6)
Ever had high cholesterol	1997-1998	21.9 (20.7-23.1)	21.8 (20.7-22.8)	21.8 (21.1-22.6)
	2002-2005	24.7 (23.2-26.3)	24.1 (22.9-25.3)	24.4 (23.5-25.4)
	2006-2009	27.4 (25.0-29.9)	26.7 (24.8-28.5)	27.0 (25.5-28.6)
Diabetes or high blood glucose	1997-1998	4.5 (4.1-5.0)	3.7 (3.4-4.1)	4.1 (3.8-4.4)
	2002-2005	7.1 (6.6-7.6)	5.5 (5.2-5.9)	6.3 (6.0-6.6)
	2006-2009	8.0 (7.4-8.6)	6.4 (6.0-6.8)	7.2 (6.8-7.6)
High and very high psychological distress	1997-1998	8.5 (7.9-9.2)	12.1 (11.4-12.8)	10.4 (9.9-10.9)
	2002-2005	9.5 (8.8-10.3)	13.3 (12.6-14.1)	11.5 (10.9-12.0)
	2006-2009	9.1 (8.3-9.9)	12.1 (11.4-12.8)	10.7 (10.1-11.2)
Visited a dental professional within the last 12 months	2002-2005	51.5 (50.3-52.7)	56.6 (55.6-57.6)	54.1 (53.3-54.9)
	2006-2009	55.5 (54.1-57.0)	59.5 (58.4-60.6)	57.6 (56.6-58.5)
All natural teeth missing	1997-1998	5.6 (4.9-6.2)	11.3 (10.5-12.2)	8.6 (8.0-9.1)
	2002-2005	4.6 (4.2-4.9)	7.9 (7.5-8.3)	6.3 (6.0-6.5)
	2006-2009	3.9 (3.5-4.2)	6.5 (6.1-7.0)	5.2 (4.9-5.5)

Indicator	Year	Males % (95% CI)	Females % (95% CI)	Persons % (95% CI)
Overweight	1997-1998	38.0 (36.8-39.1)	23.2 (22.3-24.0)	30.5 (29.7-31.2)
	2002-2005	40.6 (39.3-41.8)	25.5 (24.6-26.4)	32.9 (32.2-33.7)
	2006-2009	41.8 (40.4-43.2)	26.9 (26.0-27.9)	34.3 (33.4-35.1)
Obese	1997-1998	12.1 (11.3-12.8)	12.0 (11.4-12.7)	12.1 (11.6-12.6)
	2002-2005	16.5 (15.6-17.4)	16.7 (16.0-17.5)	16.6 (16.0-17.2)
	2006-2009	19.8 (18.7-20.9)	20.1 (19.2-20.9)	19.9 (19.2-20.6)
Overweight or obese	1997-1998	50.1 (48.8-51.3)	35.2 (34.2-36.2)	42.5 (41.7-43.3)
	2002-2005	57.1 (55.8-58.4)	42.2 (41.1-43.2)	49.5 (48.7-50.4)
	2006-2009	61.6 (60.2-63.0)	47.0 (45.9-48.1)	54.2 (53.3-55.1)
Trends in health services				
Private health insurance	1997-1998	42.8 (41.5-44.0)	43.4 (42.4-44.5)	43.1 (42.3-43.9)
	2002-2005	54.6 (53.4-55.7)	55.5 (54.5-56.4)	55.0 (54.3-55.8)
	2006-2009	56.5 (55.3-57.8)	57.8 (56.8-58.7)	57.2 (56.4-57.9)
Difficulties getting health care when needing it	1997-1998	8.6 (8.0-9.2)	11.5 (10.9-12.1)	10.1 (9.6-10.5)
	2002-2005	11.7 (11.0-12.4)	15.6 (14.9-16.2)	13.7 (13.2-14.2)
	2006-2009	14.4 (13.6-15.2)	19.7 (19.0-20.5)	17.1 (16.6-17.7)
Emergency department presentation in the last 12 months	1997-1998	16.5 (15.6-17.4)	12.8 (12.1-13.4)	14.6 (14.0-15.1)
	2002-2005	15.8 (14.9-16.6)	14.1 (13.5-14.8)	14.9 (14.4-15.4)
	2006-2009	18.3 (17.3-19.3)	16.5 (15.8-17.2)	17.4 (16.8-18.0)
Hospital admission in the last 12 months	1997-1998	11.7 (11.0-12.5)	15.6 (14.8-16.3)	13.7 (13.2-14.2)
	2002-2005	12.5 (11.8-13.2)	15.8 (15.1-16.5)	14.2 (13.7-14.7)
	2006-2009	12.7 (11.9-13.4)	17.1 (16.3-17.8)	14.9 (14.4-15.5)
Visited a general practice in the last 12 months	1997-1998	84.9 (84.0-85.7)	91.9 (91.4-92.5)	88.5 (88.0-89.0)
	2006-2009	81.3 (80.2-82.5)	90.4 (89.7-91.0)	86.0 (85.3-86.6)
	Trends in social capital			
Most people can be trusted	2002-2005	72.6 (71.4-73.8)	68.7 (67.7-69.7)	70.6 (69.9-71.4)
	2006-2009	73.8 (72.5-75.1)	72.5 (71.5-73.6)	73.2 (72.3-74.0)
	2002-2005	81.8 (80.8-82.7)	57.6 (56.6-58.7)	69.4 (68.7-70.1)
Feel safe walking down their street after dark	2006-2009	85.0 (84.1-85.9)	61.5 (60.4-62.6)	73.1 (72.3-73.8)
	2002-2005	77.3 (76.2-78.4)	74.5 (73.6-75.5)	75.9 (75.2-76.6)
	2006-2009	77.9 (76.7-79.1)	75.0 (74.0-76.0)	76.4 (75.6-77.2)
Area has a reputation for being a safe place	2002-2005	70.2 (69.0-71.3)	65.1 (64.1-66.1)	67.5 (66.8-68.3)
	2006-2009	67.3 (65.9-68.7)	63.7 (62.6-64.8)	65.5 (64.6-66.4)
	2002-2005	82.5 (81.5-83.5)	85.3 (84.6-86.0)	84.0 (83.3-84.6)
Ran into friends and acquaintances when shopping in local area	2006-2009	81.6 (80.5-82.7)	86.0 (85.2-86.8)	83.9 (83.2-84.5)
	2002-2005	70.8 (69.6-72.0)	77.0 (76.1-77.9)	74.0 (73.3-74.8)
	2006-2009	72.0 (70.7-73.3)	77.5 (76.5-78.5)	74.8 (74.0-75.6)
Would feel sad to leave their neighbourhood	2002-2005	70.8 (69.6-72.0)	77.0 (76.1-77.9)	74.0 (73.3-74.8)
	2006-2009	72.0 (70.7-73.3)	77.5 (76.5-78.5)	74.8 (74.0-75.6)

Note: Indicators include adults 16 years and over unless specified.

Source: New South Wales Population Health Survey 2006-2009 (HOIST). Centre for Epidemiology and Research, NSW Department of Health.

WARNING: Estimates New South Wales
Please check HealthStats NSW for more details.