

Influenza Weekly Epidemiology Report, NSW

22 to 28 September 2012

Produced by: Population Health Division, NSW Ministry of Health.

This report describes the surveillance for influenza and other respiratory pathogens, undertaken by NSW Health to date. This includes data from a range of surveillance systems.

For weekly communicable disease surveillance updates refer to the Communicable Disease Weekly Report at <http://www.health.nsw.gov.au/publichealth/infectious/index.asp>.

1. Summary

For the week ending 28 September 2012:

- The influenza-like illness (ILI) presentation rate to selected emergency departments (ED) decreased further this week. The rate was below the usual range for this time of year and well below the peak of activity seen in mid-July.
- ED admissions to critical care units for ILI and pneumonia decreased this week and were within the usual range for this time of year.
- Laboratory testing data showed that influenza A activity continued to decline from its peak in late June, while influenza B activity has also declined having reached its peak mid September.
- As of 7 September, the population death rate for influenza and pneumonia increased slightly but was below the epidemic threshold.

2. Emergency Department (ED) presentations


Data from 59 NSW emergency departments (ED) are included. Comparisons are made with data for the preceding five years. Recent counts are subject to change.

Source: NSW Health Public Health Real-time Emergency Department Surveillance System (PHREDSS) managed by the Centre for Epidemiology and Evidence, NSW Ministry of Health.

Presentations for influenza-like illness and other respiratory illness


- The total number of patients presenting to ED with influenza-like illness (ILI) decreased compared to the previous week (rate of 1.0 cases per 1000 presentations) and was below the usual range for this time of year (Figure 1 and Table 1).
- The total number of admissions from ED to critical care units for ILI and pneumonia decreased this week and are within the usual range for this time of year (Figure 2).

Figure 1: Total weekly counts of Emergency Department visits for influenza-like illness, from January – September 2012 (black line), compared with each of the 5 previous years (coloured lines), for 59 NSW hospitals.*



* **Note:** Excludes 2009 data to enable comparison of 2012 data with data from previous non-pandemic years. Data shown includes ED presentations up to 30 September 2012.

Figure 2: Total weekly counts of Emergency Department visits for pneumonia and influenza-like illness, which were subsequently admitted to a critical care ward, from January – September 2012 (black line), compared with each of the 5 previous years (coloured lines), for 59 NSW hospitals.



* **Note:** Data shown includes ED presentations up to 30 September 2012.

Table 1: Weekly ED and Ambulance Respiratory Activity Summary. Includes 59 NSW EDs and Sydney Ambulance Division.

Data source	Diagnosis or problem category	Trend since last week	Overall comparison with usual range for time of year	Statistically significant age groups (if any)	Statistically significant local increase (if any)	Action other than this report (if any)	Comment
ED presentations, 59 NSW hospitals*	Influenza like illness (ILI)	Decreased	Below				
	Pneumonia	Increased	Usual				
	Pneumonia and ILI admissions	Steady	Usual				
	Pneumonia and ILI critical care admissions	Decreased	Usual				
	Bronchiolitis	Steady	Above				
	Respiratory, fever and unspecified infections	Decreased	Usual				
	Asthma	Steady	Usual				
	Total presentations	Steady	Above	9% above 2011 counts			
Ambulance calls, Sydney region	Breathing problems	Decreased	Usual				

Notes on Table 1:

- (1) Statistically significant increases are shown in bold.
- (2) This report summarises activity from 59 Emergency Departments (EDs) across NSW and the Sydney Ambulance Operations Region. It provides information on general respiratory activity. Recent activity counts are subject to change.
- (3) This is a routine general report for information on respiratory activity, and is additional to public health situation reports that advise of unusual increases in activity in particular provisional ED diagnosis groupings or Ambulance problem categories. It is prepared by the Centre for Epidemiology and Intelligence.

3. Laboratory testing summary for influenza

For the week ending 28 September 2012:

- A total of 1092 tests for respiratory viruses were performed at sentinel NSW laboratories (Table 2) with 8.0% testing positive for influenza.
- Influenza A: 22 specimens (2.0%) tested positive (Table 2, Figure 4). Of these:
 - 7 (31.8%) tested positive for influenza A(H3N2)
 - Two tested positive for influenza A(pH1N1). The remainder tested negative to influenza A(pH1N1) and are assumed to have been A(H3N2)
- Influenza B: 65 specimens (6.0%) tested positive (Table 2, Figure 4).
- The proportion of respiratory specimens positive for influenza A decreased further compared to the previous week, influenza B activity decreased as well compared to the previous weeks.

Influenza activity continues to decline and is no longer the dominant respiratory virus identified by NSW sentinel laboratories.


Table 2: Summary of testing for influenza and other respiratory viruses at NSW laboratories, 1 January to 28 September 2012.

Month ending	Total Tests	Influenza A		A(H3N2)		A(pH1N1) *		Influenza B		Adeno.	Parainf. 1, 2 & 3	RSV	Rhino.	Entero.	HMPV**
		Total	(%)	Total	(%Flu A) *	Total	(%Flu A) *	Total	(%)						
27/01/2012	1617	14	(0.9%)	6	(42.9%)	4	(28.6%)	7	(0.4%)	37	60	38	119	64	36
02/03/2012*	2520	31	(1.2%)	12	(38.7%)	1	(3.2%)	15	(0.6%)	44	65	156	224	128	30
30/03/2012	2573	36	(1.4%)	25	(69.4%)	3	(8.3%)	16	(0.6%)	59	79	269	263	114	40
27/04/2012	2857	46	(1.6%)	31	(67.4%)	5	(10.9%)	11	(0.4%)	65	63	422	231	114	28
1/06/2012	4394	209	(4.8%)	166	(79.4%)	2	(1.0%)	30	(0.7%)	91	76	574	463	170	31
29/06/2012	5704	1316	(23.1%)	613	(46.6%)	2	(0.2%)	84	(1.5%)	96	68	558	535	16	53
27/07/2012	6818	1552	(22.8%)	982	(63.3%)	5	(0.3%)	159	(2.3%)	138	70	551	552	13	88
31/08/2012*	7781	915	(11.8%)	556	(60.8%)	10	(1.1%)	344	(4.4%)	165	145	515	577	34	189
Week ending															
7/09/2012	1466	65	(4.4%)	38	(58.5%)	3	(4.6%)	103	(7.0%)	40	48	86	122	4	52
14/09/2012	1348	52	(3.9%)	33	(63.5%)	0	(0.0%)	98	(7.3%)	36	56	54	105	11	51
21/09/2012	1190	39	(3.3%)	16	(41.0%)	1	(2.6%)	107	(9.0%)	46	41	57	109	2	58
28/09/2012	1092	22	(2.0%)	7	(31.8%)	2	(9.1%)	65	(6.0%)	40	56	42	111	7	42

* Subset of influenza A positive tests; ** HMPV = Human metapneumovirus

Note: Data is provided by laboratories on a weekly basis. Includes point of care tests as of 10 August 2012. Influenza laboratory diagnoses using virology are reported by South Eastern Area Laboratory Services (SEALS), Institute of Clinical Pathology and Medical Research (ICPMR), The Children’s Hospital at Westmead (CHW), Sydney South West Area Services (SSWPS), Pacific Laboratory Medicine Services (PaLMS), Royal Prince Alfred Hospital (RPAH), Hunter Area Pathology Service (HAPS), St Vincent’s (SydPath), Nepean, Douglas Hanley Moir (DHM) , VDRLab.

Figure 4: Percent of respiratory samples positive for influenza A or influenza B, 1 January 2007 – 28 September 2012, New South Wales.



Note: Data is provided by laboratories on a weekly basis. Includes point of care tests as of 10 August 2012. Influenza laboratory diagnoses using virology are reported by South Eastern Area Laboratory Services (SEALS), Institute of Clinical Pathology and Medical Research (ICPMR), The Children’s Hospital at Westmead (CHW), Sydney South West Pathology Services (SSWPS), Pacific Laboratory Medicine Services (PaLMS), Royal Prince Alfred Hospital (RPAH), Hunter Area Pathology Services (HAPS) , St Vincent’s (SydPath) , Nepean (no data between Oct 2010 to June 2011), Douglas Hanley

Moir (DHM) , VDRLab from 5 March 2010 , Lavery (data from 1 April 2010 to February 2011) and St Vincent's (data since November 2010).

Laboratory-confirmed Influenza outbreaks in residential care facilities

There were no respiratory outbreaks in residential care facilities reported this week associated with influenza.

In the year to date (up to week ending 28 September), there have been 38 laboratory confirmed influenza outbreaks in institutions reported to NSW Public Health Units (Table 3). All but three outbreaks occurred in an aged care facility. At least 764 residents were reported to have had ILI symptoms and 57 required hospitalisation. Twenty-seven deaths in residents linked to the outbreaks have been reported, all of whom were noted to have other significant co-morbidities.

Table 3. Reported influenza outbreaks in NSW institutions, 2005-2012.

Year	2005	2006	2007	2008	2009	2010	2011	2012*
No. of outbreaks	5	2	25	9	1	2	4	38

*Preliminary data up to 28 September 2012. These data are subject to change as more information is obtained.

Respiratory outbreaks in aged care facilities were uncommon from 2009 to 2011, and this is thought to be due to the predominance of the influenza A(pH1N1) strain in these years, against which people in older age-groups appeared to have higher levels of protection.

4. Deaths with pneumonia or influenza reported on the death certificate


Deaths registration data is routinely reviewed for deaths attributed to pneumonia or influenza. While pneumonia has many causes, a well-known indicator of seasonal and pandemic influenza activity is an increase in the number of death certificates that mention pneumonia or influenza as a cause of death.

The predicted seasonal baseline estimates the predicted rate of influenza or pneumonia deaths in the absence of influenza epidemics. If deaths exceed the epidemic threshold, then it may be an indication that influenza is beginning to circulate widely.

For the week ending 7 September:

- There were 1.62 pneumonia or influenza deaths per 100,000 NSW population, which is below the epidemic threshold of 1.67 per 100,000 population (Figure 5).*
- Between 1 July and 31 August 2012, out of 10194 deaths there were 27 death certificates mentioning influenza, and 1345 mentioning pneumonia. The majority of these influenza and pneumonia deaths were in persons aged greater than 65 years.
- The updated data on pneumonia and influenza deaths indicates that the rate of deaths in this category was above the epidemic threshold for most of July. As expected, the increase in the death rate has mirrored the increases seen in laboratory isolations of influenza and Emergency Department ILI activity, but with a delay of one to two weeks.

Figure 5: Rate of deaths classified as influenza and pneumonia (by NSW Registered Death Certificates) per 100,000 NSW population, 2007 - 2012.



Source: NSW Registry of Births, Deaths and Marriages.

*** Notes on interpreting death data:**

- (1) The number of deaths mentioning "Pneumonia or influenza" is reported as a rate per 100,000 NSW population. Using the NSW population provides a more stable and reliable denominator than deaths from all causes. This is because pneumonia and influenza are known to contribute to increases in deaths from non-respiratory illnesses, such as deaths due to ischaemic heart disease. As the number of these deaths will increase with rises in influenza activity, the actual effect of influenza on mortality rates will be obscured if all-cause mortality is used as the denominator. This limitation is avoided by using the NSW population, which is relatively constant throughout the year, as the denominator.
- (2) Deaths referred to a coroner during the reporting period may not be available for analysis. Deaths in younger people may be more likely to require a coronial inquest. Therefore influenza-related deaths in younger people may be under-represented in these data.
- (3) The interval between death and death data availability is usually at least 7 days, and so these data are one week behind reports from emergency departments and laboratories. In addition, previous weekly rates may also change due to longer delays in reporting some deaths.

5. National and International Influenza Surveillance and Links

Novel Swine-Origin triple reassortant H3N2 viruses in the United States

Up to 28 September, the US CDC reported that multiple human infections with variant influenza A (H3N2v) viruses had been identified across 11 US states, bringing the total to 306 cases since it was first reported July 2011(see updated case counts at: <http://www.cdc.gov/flu/swineflu/h3n2v-case-count.htm>). Since July 2012, 16 cases have required hospitalisation and there has been one death reported.

As a result of enhanced surveillance activities for H3N2v, one infection with an influenza A (H1N1) variant (H1N1v) virus has been detected in Missouri in a patient who became ill after contact with swine. The patient has recovered from their illness. Confirmatory testing at CDC identified H1N1v with the matrix (M) gene from the 2009 H1N1 influenza virus in specimens collected from this

patient. Cases of H1N1v have been detected previously, and the current case marks the second report of H1N1v with the M gene from the 2009 H1N1 virus.

The H1N1v case, in addition to the H3N2v cases outlined above and the H1N2v cases reported previously, brings the total number of variant influenza virus infections detected since July 2012 to 309.

Links to Other Influenza Surveillance

Australian Influenza Surveillance Reports:

<http://www.health.gov.au/internet/main/publishing.nsf/Content/cda-ozflu-2012.htm>

World Health Organization Influenza Updates:

<http://www.who.int/csr/disease/influenza/en/index.html>

WHO Collaborating Centre for Reference and Research on Influenza (Melbourne):

<http://www.influenzacentre.org/index.htm>