

# NSW Health Influenza Surveillance Report

Week 37: 6 to 13 September 2015

## Summary:

- **Influenza activity continues to trend down, but variation remains across the state.**
- **The impact on emergency departments decreased but remains high in some local health districts and in the 5-34 year age group.**
- **Influenza B strains continuing to predominate.**
- **Based on previous seasons, influenza activity will likely continue at lower levels during September.**

## In this reporting week:

- [Hospital surveillance](#) – presentations to NSW emergency departments for influenza-like illness (ILI) decreased again this week. Bronchiolitis presentations increased slightly this week and remain above the usual range for this time of year.
- [Laboratory surveillance](#) – the proportion of respiratory samples positive for influenza was moderate to high at 28.9%, but continues to trend down. Influenza B viruses continue to predominate.
- [Community surveillance](#) – influenza notifications across all but one local health district (South Western Sydney LHD) decreased this week are expected to decline further in the coming weeks. Data collected from ASPREN, FluTracking and eGPS showed declines in seasonal ILI activity. Three new outbreaks were reported in residential aged care facilities.
- [National and international influenza surveillance](#) – across Australia, seasonal influenza activity appears to have peaked in recent weeks with the exception of South Australia where activity continues to rise. The timing and magnitude of the peak is similar to 2014.

## About this report:

Health Protection NSW collects and analyses surveillance data on influenza and other respiratory viruses. Surveillance reports are produced weekly commencing in May, and continuing until the end of the influenza season. Monthly reports are produced throughout the rest of the year.

The influenza surveillance reports include data from a range of surveillance systems and sources concerned with Emergency Department illness surveillance, laboratory (virological) surveillance, and community illness surveillance. Pneumonia and influenza mortality data are also monitored and reported upon periodically.

For further information on influenza see the [NSW Health Influenza website](#).

# 1. Hospital Surveillance


## NSW emergency department (ED) presentations for influenza-like illness (ILI) and other respiratory illnesses

Source: PHREDSS<sup>1</sup>

For the week ending 13 September 2015:

- ILI presentations<sup>2</sup> decreased this week. Activity remained well above the usual range of activity seen in recent years (Figure 1 and Table 1), although trending down.
- The index of increase for ILI presentations was 24.6 on 13 September, higher than the previous week (44.6). The index crossed the threshold level of 15 on 26 June and peaked at 64.2 on 19 August (higher than the peak of 50.7 seen in 2014).
- The proportion of ILI presentations to all ED presentations is moderate at 4.0 per 1000 presentations up from the previous week (6.2 per 1000 presentations). Presentations were particularly elevated among persons aged 5 to 34 years, and in Western Sydney and Mid North Coast LHDs, and at The Children's Hospital at Westmead, Lithgow, Royal North Shore and Bankstown Hospital's (Table 1).
- ED presentations for pneumonia<sup>3</sup> continued to decrease but remained above the usual range for this time of year. Presentations were notably elevated in South Western Sydney LHD and Moree Hospital (Figure 2 and Table 1).
- Pneumonia or ILI presentations which resulted in admission decreased, but remained above the usual range for this time of year. Admissions to critical care decreased and were within the usual range (Figure 3 and Table 1).
- The category combining all respiratory, fever and unspecified infection presentations decreased but remained above the usual range for this time of year; presentations were elevated in all age groups. Presentations were increased in South Western Sydney, Northern Sydney, Western Sydney, Sydney and Western NSW LHDs (Table 1).
- Bronchiolitis presentations increased this week and remained above the usual range for this time of year. Presentations were elevated at Singleton Hospital (Table 1).

**Figure 1:** Total weekly counts of ED visits for influenza-like illness, from January – 13 September 2015 (black line), compared with each of the 5 previous years (coloured lines).




<sup>1</sup> NSW Health Public Health Rapid, Emergency Disease and Syndromic Surveillance system. Managed by the Centre for Epidemiology and Evidence, NSW Ministry of Health. Data from 59 NSW emergency departments are included. Comparisons are made with data for the preceding five years. Recent counts are subject to change. This includes data from 59 NSW emergency departments (EDs), representing approximately 85% of metropolitan ED presentations and approximately 60% of rural ED presentations.


<sup>2</sup> ILI is when the treating ED doctor makes a provisional clinical diagnosis of ILI Syndrome, which includes: 'influenza-like illness' or 'influenza' (including 'pneumonia with influenza').

<sup>3</sup> Pneumonia is when there is a provisional clinical diagnosis of Pneumonia Syndrome, which includes: 'viral, bacterial or unspecified pneumonia', 'SARS', or 'legionnaire's disease'. Excludes 'pneumonia with influenza'.

**Figure 2:** Total weekly counts of ED presentations for pneumonia, from January – 13 September 2015 (black line), compared with each of the 5 previous years (coloured lines).



**Figure 3:** Total weekly counts of ED presentations for pneumonia or influenza-like illness and admitted to a critical care ward, from January – 13 September 2015 (black line), compared with each of the 5 previous years (coloured lines).



**Table 1:** Weekly ED and Ambulance Respiratory Activity Summary for the week ending 13 September 2015. Includes data from 59 NSW EDs and the NSW Ambulance Division. \*

Data source	Diagnosis or problem category	Trend since last week	Comparison with usual range for time of year	Statistically significant age groups (if any)	Statistically significant local increase (if any)	Action other than this report (if any)	Comment
ED presentations, 59 NSW hospitals	Influenza like illness (ILI)	Decreased	<b>Above</b>	5-34 years	Western Sydney, Mid North Coast LHDs, and at The Children's Hospital at Westmead and Wollongong, Lithgow, Royal North Shore and Bankstown Hospital's	Daily ILI Situation reports have been sent since 27 August to support LHD planning	
	Pneumonia	Decreased	<b>Above</b>		South Western Sydney LHD Moree Hospital		
	Pneumonia and ILI admissions	Decreased	Above				
	Pneumonia and ILI critical care admissions	Decreased	Usual				
	Bronchiolitis	Increased	Above		Singleton Hospital		Bronchiolitis is a disease of infants.
	Respiratory illness, fever or unspecified infections	Decreased	<b>Above</b>	<b>All age groups</b>	South Western Sydney, Northern Sydney, Western Sydney, Sydney and Western NSW LHD's		
	Asthma	Increased	Above				
Ambulance Triple Zero (000) calls, NSW	Breathing problems	Steady	<b>Above</b>				

\* **Notes on Table 1:** Statistically significant increases are shown in bold. Recent activity counts are subject to change. This is a routine general report for information on respiratory activity and is additional to public health situation reports that advise of unusual increases in activity in particular provisional ED diagnosis groupings or Ambulance problem categories.


## FluCAN (The Influenza Complications Alert Network)

In 2009, the [FluCAN](#) surveillance system was created to be a rapid alert system for severe respiratory illness requiring hospitalisation. Data is provided on patients admitted with influenza confirmed by polymerase chain reaction (PCR) testing.

In NSW, three hospitals participate in providing weekly FluCAN data: Westmead Hospital, John Hunter Hospital and the Children's Hospital at Westmead.

- During week 37 there were 10 (1 adult and 9 children) influenza admissions in NSW sentinel hospitals (Figure 4).
- Since 1 April 2015, there have been 223 hospital admissions reported for influenza; 92 with influenza A, 127 with influenza B and one dual infection and 3 unknown (Figure 4).
- Of these admissions, 87 were paediatric (<16 years of age) cases and 136 were in adults. Seventeen cases were admitted to ICU/HDU.

**Figure 4:** FluCAN – weekly number of confirmed influenza hospital admissions in NSW, April – September 2015.



## 2. Laboratory Surveillance

For the week ending 13 September 2015 the number and proportion of respiratory specimens reported by NSW sentinel laboratories [4] which tested positive for influenza A or influenza B continued to decrease compared to the activity levels seen in the previous week suggesting that the peak of the season has passed (Table 2 and Figures 5-6).

A total of 9,540 tests for respiratory viruses were reported with 28.9% testing positive for influenza viruses, slightly down from 35.5% in the previous week. Of these, Influenza B viruses continued to be identified twice as often as influenza A viruses.

[4]: Preliminary laboratory data is provided by participating sentinel laboratories on a weekly basis and are subject to change. Point-of-care test results have been included since August 2012 but serological diagnoses are not included.

**Participating sentinel laboratories:** South Eastern Area Laboratory Services, The Children's Hospital at Westmead, Sydney South West Pathology Service, Pacific Laboratory Medicine Service, Royal Prince Alfred Hospital, Hunter Area Pathology Service, Pathology West (Westmead & Nepean), Douglas Hanley Moir Pathology, VDRLab, Laverty Pathology, SydPath (St Vincent's), Medlab, and Laverty.

Influenza viruses were the leading respiratory viruses reported. Other viruses are circulating at usual levels for this time of year (Table 2).


**Table 2:** Summary of testing for influenza and other respiratory viruses at NSW laboratories, 1 January to 13 September, 2015.

Month ending	Total Tests	TEST RESULTS															
		Influenza A						Influenza B		Adeno	Parainf 1, 2 & 3	RSV	Rhino	Entero	HMPV **		
		Total	H3N2	H1N1 pdm09	A (Not typed)		Total										
Total	(%)	Total	(%A)	Total	(%A)	Total	(%A)	Total	(%)								
01/02/2015*	5920	182	(3.1%)	40	(22.0%)	11	(6.0%)	131	(72.0%)	55	(0.9%)	150	181	181	607	59	49
01/03/2015	6287	212	(3.4%)	72	(34.0%)	14	(6.6%)	126	(59.4%)	75	(1.2%)	128	83	271	842	24	29
29/03/2015	8577	242	(2.8%)	87	(36.0%)	21	(8.7%)	135	(55.8%)	108	(1.3%)	181	117	767	1084	52	34
03/05/2015*	12584	285	(2.3%)	125	(43.9%)	13	(4.6%)	147	(51.6%)	163	(1.3%)	257	187	1351	1443	59	78
31/05/2015	12244	128	(1.0%)	42	(32.8%)	9	(7.0%)	83	(64.8%)	200	(1.6%)	272	167	1276	1514	64	64
28/06/2015	15431	297	(1.9%)	56	(18.9%)	16	(5.4%)	225	(75.8%)	581	(3.8%)	378	183	1585	2027	96	135
02/08/2015*	22771	1125	(4.9%)	332	(29.5%)	141	(12.5%)	654	(58.1%)	2125	(9.3%)	721	273	1878	2484	149	425
30/08/2015*	32606	3717	(11.4%)	1428	(38.4%)	595	(16.0%)	1728	(46.5%)	7819	(24.0%)	747	295	1014	2369	69	445
Summary of weekly influenza positive test results																	
Week ending																	
06/09/2015	9933	1093	(11.0%)	379	(34.7%)	203	(18.6%)	511	(46.8%)	2429	(24.5%)	235	101	180	600	16	137
13/09/2015	9540	914	(9.6%)	375	(41.0%)	136	(14.9%)	403	(44.1%)	1838	(19.3%)	370	127	227	545	17	129


**Notes:**

- \* Five-week reporting period.
- \*\* Human metapneumovirus

**Figure 5:** Weekly influenza positive test results by type and sub-type reported by NSW sentinel laboratories, 1 January to 13 September 2015.



**Figure 6:** Percentage of laboratory tests positive for influenza A and influenza B by week, 1 January 2010 – 13 September 2015, New South Wales.



### 3. Community Surveillance

#### Influenza notifications by Local Health District (LHD)

In the week ending 13 September there were 2814 notifications of influenza confirmed by polymerase chain reaction (PCR) testing, down from 3371 notifications in the previous week.

Districts with the highest notification rates were Western Sydney, Northern Sydney, and South Western Sydney LHDs (Table 3). Influenza activity decreased across all LHDs with the exception of South Western Sydney.

**Table 3:** Weekly notifications of laboratory-confirmed influenza by Local Health District.

Local Health District	Week ending 13 Sep 2015		Previous 4 weeks	
	Number of notifications	Rate per 100 000 population	Average weekly notifications	Rate per 100 000 population
Central Coast	94	28.09	97	28.99
Far West	1	3.26	2	6.52
Hunter New England	257	28.25	301	33.12
Illawarra Shoalhaven	89	22.29	114	28.55
Mid North Coast	40	18.62	45	20.94
Murrumbidgee	46	19.26	85	35.72
Nepean Blue Mountains	136	36.95	227	61.67
Northern NSW	55	18.5	80	26.91
Northern Sydney	478	53.23	630	70.16
South Eastern Sydney	235	26.33	433	48.54
South Western Sydney	481	50.9	419	44.38
Southern NSW	27	13.14	45	21.74
Sydney	250	40.31	308	49.66
Western NSW	65	23.44	66	23.92
Western Sydney	560	60.42	619	66.78

**Note:** \* All data are preliminary and may change as more notifications are received. Excludes notifications based on serology.

## Influenza outbreaks in institutions

There were 3 influenza outbreaks reported in residential aged care facilities this week. One was due to influenza A and two had both A and B strains identified.

In the year to date, there have been 83 laboratory confirmed influenza outbreaks in institutions reported to NSW public health units (Table 4): 48 have been due to influenza A, 25 due to influenza B and 10 were combined A and B. At least 780 residents were reported to have had ILI symptoms and 116 required hospitalisation. Twenty-six deaths in residents linked to these outbreaks have been reported, all of whom were noted to have other significant co-morbidities.

People in older age-groups are at higher risk of infection from influenza A(H3N2) strains than from the influenza A(H1N1) strain. The influenza A(H3N2) strain predominated in 2012 and 2014 and was associated with an increase in influenza outbreaks in institutions, particularly residential aged care facilities (Table 4).

**Table 4.** Reported influenza outbreaks in NSW institutions, 2010 to 13 September 2015.

Year	2010	2011	2012	2013	2014	2015 *
No. of outbreaks	2	4	39	12	120	83

\* Year to date.

## Electronic General Practice Surveillance (eGPS)


eGPS is a primary care influenza surveillance system involving sentinel general practices within three NSW Local Health Districts (LHD): Northern Sydney (NS), South Eastern Sydney (SES) and Illawarra Shoalhaven (IS). The system monitors patient consultations for influenza-like illness (ILI) as an indicator of influenza activity. Consultations for ILI are identified each week by an automatic search of electronic records for validated combinations of ILI terms rather than diagnosis codes.

Data generated from eGPS should be interpreted with caution as they are not representative of all practices within the participating LHDs or across NSW.

In Week 37:

- There were 6 surveillance reports received from eGPS sentinel practices in NSW; no reports were received from Illawarra Shoalhaven this week.
- The average rate of ILI patient consultations decreased to 2.2% (range 1.2 – 2.7%), down from 2.9% in the previous week and within the usual range seen for this time of year (Figure 7).

**Figure 7.** Average rate of influenza-like presentations to sentinel general practices by week of consultation 2011-2015 (year to date).



## The Australian Sentinel Practices Research Network (ASPREN)

ASPREN is a network of sentinel general practitioners (GPs) run through the Royal Australian College of General Practitioners and the University of Adelaide which has collected de-identified information on influenza-like illness (ILI) and other conditions seen in general practice since 1991. Participating GPs in the program report on the proportion of patients presenting with an ILI. The number of GPs participating on a weekly basis may vary.

In week 37 there were 33 ASPREN reports received from NSW GPs. The overall consultation rate for ILI was moderate at 2.5 %, slightly down from the rate of 2.6% in the previous week.

For further information please see the [ASPREN](#) website.


## FluTracking.net

FluTracking.net is an online health surveillance system to detect epidemics of influenza. FluTracking is a project of the University of Newcastle, the Hunter New England Local Health District and the Hunter Medical Research Institute. It involves participants from around Australia completing a simple online weekly survey which is used to generate data on the rate of ILI symptoms in communities.

In week 37 FluTracking received reports for 6248 people in NSW with the following results:

- 3.2% of respondents reported fever and cough, lower than the previous week (3.5%) (Figure 8), and continuing a downward trend.
- 1.9 % of respondents reported fever, cough and absence from normal duties, the same as the previous week (data not shown).

**Figure 8:** FluTracking – weekly influenza-like illness reporting rate, NSW, 2010 – 2015.



For further information, including national estimates, please see the [FluTracking](#) website.



## 4. National and International Influenza Surveillance

### National Influenza Surveillance

The Australian Department of Health has reported up to 28 August 2015.

Across most jurisdictions, seasonal influenza activity appears to have peaked in recent weeks with the exception of South Australia where activity continues to rise. The timing and magnitude of the peak is similar to 2014.

- Influenza notification rates have been highest among those aged between 5 and 9 and over 85 years with a secondary peak in those aged 40-44 years.
- Influenza B continues to be the dominant influenza virus type nationally, comprising over two thirds of all notifications. In the Australian Capital Territory and Western Australia, influenza A continues to replace influenza B.
- All systems that monitor influenza-like illness (ILI) activity are reporting decreasing activity following a season peak in the week ending 23 August. Influenza is the primary cause of ILI in the community this fortnight however other respiratory viruses continue to circulate at elevated levels.
- Data for hospitalisations with confirmed influenza show high influenza activity which is typical for mid-season. Influenza B continues to account for more than half of admissions.
- The seasonal influenza vaccines appear to be a good match for circulating strains with 83% of samples matching the trivalent seasonal vaccine (TIV).

Follow the link for the [Australian Influenza Surveillance Reports](#) which provide the latest information on national influenza activity.

### Global Influenza Update

The World Health Organization (WHO) provides [weekly reports](#) of global influenza activity. As of 7 September 2015 (with data up to 23 August), global surveillance indicated that:

- In the Northern Hemisphere countries, respiratory virus activity remained low in general, and influenza activity continued at low, inter-seasonal levels. Influenza type A predominated in sporadic detections. A number of countries have also scaled down surveillance activity during the inter-seasonal period.
- In Eastern Africa, in countries with reported influenza activity, influenza type A predominated. In Western Africa, influenza activity decreased overall, with influenza B predominating in Ghana and influenza A in Côte d'Ivoire.
- In tropical countries of the Americas, Central America and the Caribbean, influenza activity remained at low levels, with the exception of Cuba, where high levels of influenza-like illness (ILI) and severe acute respiratory infections (SARI) were reported, associated with influenza A(H1N1)pdm09 and RSV viruses detections.
- In tropical Asia, countries in Southern Asia and South East Asia reported an overall low influenza activity though India reported a minor increase in activity with predominantly A(H1N1)pdm09. Influenza activity was still high in southern China with influenza A(H3N2) predominating.
- In temperate South America, ILI and SARI activity remained low and continued to decrease in general, except in Chile, where respiratory virus activity remained elevated. Influenza type A viruses predominated in the region.
- In South Africa, influenza activity decreased, with influenza type B predominating in recent weeks.
- In New Zealand, influenza activity may have peaked in the second week of August with influenza A(H3N2) and B predominating.

WHO reported global influenza laboratory data for the period 10 to 23 August 2015, which noted:

- Of the 32 226 specimens submitted for testing, 4246 were positive for influenza viruses, of which 3219 (76%) were typed as influenza A and 1027 (24%) as influenza B.
- Of the sub-typed seasonal influenza A viruses, 326 (12%) were influenza A (H1N1) and 2350 (88%) were influenza A(H3N2).
- Of the characterized B viruses, 126 (92%) belonged to the B/Yamagata lineage and 11 (8%) to the B/Victoria lineage.

## **Avian influenza Update**

### **Human infection with avian influenza A(H5) viruses**

WHO report that from 2003 through to 17 July 2015, 844 laboratory-confirmed human cases of avian influenza A(H5N1) virus infection have been officially reported to WHO from 16 countries. Of these cases, 449 have died.

Since the last WHO Influenza update on 23 June 2015, two new laboratory-confirmed human cases of avian influenza A(H5N1) virus infection were reported to WHO from Egypt.

### **Overall public health risk assessment for avian influenza A(H5) viruses:**

Whenever avian influenza viruses are circulating in poultry, sporadic infections and small clusters of human cases are possible in people exposed to infected poultry or contaminated environments, therefore sporadic human cases would not be unexpected.

### **Human infections with avian influenza A(H7N9) viruses in China**

A total of 677 laboratory-confirmed cases of human infection with avian influenza A(H7N9) viruses, including at least 275 deaths have been reported to WHO. The majority of recently reported human cases are associated with exposure to infected live poultry or contaminated environments, including markets where live poultry are sold. WHO advises that further sporadic human cases of avian influenza A(H7N9) infection are expected in affected and possibly neighbouring areas.

WHO is assessing the epidemiological situation and conducting further risk assessment based on the latest information. Overall, the public health risk from avian influenza A(H7N9) viruses has not changed.

### **Human infections with avian influenza A(H5N6) viruses in China**

One laboratory-confirmed case of human infection with avian influenza A(H5N6) virus was reported to WHO from China. The case developed symptoms on 6 July and was admitted to hospital on 9 July and, despite medical treatment, died on 10 July.

### **Overall public health risk assessment for avian influenza A(H9N2) viruses:**

Further human cases and small clusters could occur as this virus is circulating in poultry populations across Asia and Middle East. This virus does not seem to transmit easily between humans and tends to result in mild clinical disease; therefore the current likelihood of community-level spread and public health impact of this virus is considered low.

The latest WHO monthly risk assessment report for human infections with avian influenza A strains H5, H7, H9 is available here: [WHO Avian influenza monthly summary](#) .

Other sources of information on avian influenza and the risk of human infection include the following:

- US CDC [Avian influenza](#)
- European CDC (ECDC) [Avian influenza](#)
- Public Health Agency of Canada [Avian influenza H7N9](#) .

## Composition of 2015 Australian influenza vaccines

WHO changed its recommendations for the composition of trivalent vaccines for use in the 2015 influenza season (southern hemisphere winter) as follows:

- A/California/7/2009 (H1N1)pdm09-like virus;
- A/Switzerland/9715293/2013 (H3N2)-like virus <sup>a</sup>;
- B/Phuket/3073/2013-like virus (B/Yamagata lineage).

<sup>a</sup> A/South Australia/55/2014, A/Norway/466/2014 and A/Stockholm/6/2014 are A/Switzerland/9715293/2013-like viruses

It is recommended that quadrivalent vaccines containing two influenza B viruses contain the above three viruses and a B/Brisbane/60/2008-like (i.e. B/Victoria lineage) virus.

These changes from the previous vaccine recommendations (for the southern hemisphere in 2014 and the northern hemisphere in 2014-2015) reflect observed antigenic drift in circulating A(H3N2) and B/Yamagata lineage viruses. More details about the most recent recommendations can be found at: [http://www.who.int/influenza/vaccines/virus/recommendations/2015\\_south/en/](http://www.who.int/influenza/vaccines/virus/recommendations/2015_south/en/) .