NSW Arbovirus Surveillance & Mosquito Monitoring Program, 2017-2018

Weekly Update: 16/Feb/2018





creating better health & justice systems www.pathology.health.nsw.gov.au

Contents

Summary	2
Comment	2
Environmental Conditions	3
Rainfall	
Three Month Rainfall & Temperature Forecast	3
Tidal	4
MVEV Climatic Models	5
Forbes' Hypothesis	5
Nichol's Hypothesis	5
Arboviral Isolates	6
Exotic Detections	6
Human Notifications	7
Monthly RRV notifications	9
Monthly BFV notifications	9
Mosquito Results	10
Inland	11
Coastal	
Sydney	13
Sentinel Chicken Flocks	14



Summary

- **Climate**: over the last week, there was light rainfall along the coast. For January, rainfall was below average for the north east of the state and average elsewhere.
- Three Month Forecast: for March to May 2018, rainfall predictions for NSW are for average precipitation for most of the state, with a slight chance of exceeding the average along the coast and being below average in the far west. Maximum and minimum temperatures are expected to be around average, but above average in the west of the state. According to the BOM as 13/Feb/2018, the weak La Niña continues to decline and will end in autumn, hence the drier conditions of late.
- **Tidal**: a small series of high tides are occurring now over 16-17/Feb, with heights of 1.75m predicted. The tide height at Sydney has reached 1.83m and could trigger some *Aedes vigilax* hatching. Naturally, tides in other parts of the state may be subject to more variation.
- **MVEV models**: the data relevant to both the Forbes' and Nichols' hypotheses have been updated to the end of Jan 2018. Both models are not suggestive of an MVEV epidemic.
- **Mosquito Numbers Inland**: mosquito numbers continue to be lower than average, although still 'high' at Griffith.
- Mosquito Numbers Coast: collections were generally larger this week, with the first 'very high' collection of the season from Ballina, with almost 3,000 mosquitoes trapped at Lennox Heads (albeit, few Aedes vigilax). Coffs Harbour and Tweed produced 'high' mosquito numbers.
- **Mosquito Numbers Sydney**: Georges River produced a 'very high' catch, with almost 4,000 mosquitoes from Alfords Point, the majority being *Aedes vigilax*. Bankstown and Homebush both had 'high' catches, dominated by *Aedes vigilax*.
- **Arboviral Isolates**: there were two arboviral detections, both being Ross River virus and both from Griffith.
- Chicken Sentinel Flocks: there have been no seroconversions.
- Human Notifications: for the current fiscal year, there have been 270 RRV and 53 BFV notifications, this is well below the average compared with recent years (the averages to the end of January are 464RRV and 43BFV).


Comment: we have had further RRV detections from Griffith, but these are not unusual at this time of the year and mosquito numbers from the inland continue to be below normal. For the coast, we have seen an increase in mosquito numbers, especially *Aedes vigilax* at the Sydney sites. Arboviral notifications of local disease continue to be below average. On this, it is worth noting that more than 200 RRV notifications for this season were reported up to Nov 2016 and do not relate to the current seasons activity. Thus RRV cases for this season are, in actual fact, extremely low.



Environmental Conditions

Rainfall

Rainfall across Australia for the week ending 8/Feb/2018 is depicted on the left and monthly rainfall deciles for December 2017 are on the right. Over the last week, there was light rainfall along the coast and across the north of the state. For January, rainfall was below average for the north east of the state and average elsewhere. Maximum and minimum temperatures for January were well above average; up to five degrees higher than normal in the west of the state.



Three Month Rainfall & Temperature Forecast

For March to May 2018, rainfall predictions for NSW are for average precipitation for most of the state, with a slight chance of exceeding the average along the coast and being below average in the far west. Maximum and minimum temperatures are expected to be around average, but above average in the west of the state. The following pages contain graphics of the seasonal outlook:

<u>www.bom.gov.au/climate/outlooks/#/rainfall/median</u> (Rainfall outlook). <u>www.bom.gov.au/climate/outlooks/#/temperature/summary</u> (Max & min temperature outlook).


According to the BOM as of 13/Feb/2018, the weak La Niña continues to decline and will end in autumn. The Indian Ocean Dipole (IOD) remains neutral. This all suggests that rainfall patterns are likely to return to normal in the upcoming months.

For more information: <u>www.bom.gov.au/climate/enso/</u> and, <u>http://www.bom.gov.au/climate/iod/</u>



Tidal

Tidal information is relevant for the prediction of the activity of the salt marsh mosquito, *Aedes vigilax*. Typically for NSW, tides of over 1.8m, as measured at Sydney, can induce hatching of *Aedes vigilax* larvae and the graph below of predicted tide heights can provide some indication of when this is likely to occur. Note this height various between regions, thus at Batemans Bay, a tide height over 0.8m can trigger egg hatching.



Currently, there are a small series of high tides occurring now over 16-17/Feb, with heights of 1.75m predicted. The tide height at Sydney has reached 1.83m and could trigger some *Aedes vigilax* hatching. Tides in other parts of the state may be subject to more variation. The next larger series are due over 27/Feb to 4/Mar, with heights of 1.94m being predicted.

Note that actual tide heights can vary by 0.3m (or more in unusual circumstances) due to variations in atmospheric pressure, rainfall, wind and other climatic phenomena. Climate change will also result in much higher tide heights. Thus predicted tide height should be used as a gauge only for potential *Aedes vigilax* activity. The larvae of the saltmarsh mosquito relies on a inundation/drying cycle for the mudflats in which it lives; continual wet weather prevents the drying cycles thereby reducing larval production.



MVEV Climatic Models

Three predictive environmental based models for MVEV activity have been developed; the Forbes (which relies on rainfall in the river catchment basins of Eastern Australia), Nichols (based on the Southern Oscillation), and the Bennett theory (based on the Indian Ocean Dipole). The latter theory is poorly developed (and unreliable), and is not considered below. Note that all the predictive models have been developed on a limited data set and do not always forecast activity. There can also be unusual environmental conditions that may lead to the introduction of the virus to southeastern Australia, such as the movement of low pressure cells from the north to the south of the country during 2008 and 2011. Vertical transmission of the virus (from adult to the egg in *Aedes* species) can result in restricted activity following localized heavy precipitation (as per 2003 at Menindee).

i. Forbes' Hypothesis

Rainfall was not above Decile 7 in all of the river catchment basins in eastern Australia for the last quarter of 2016 or for the first quarter of 2017 (Table 1). For the last quarter of 2017, rainfall was above Decile 7 in only one catchment basin.

Table 1. Rainfall indices for the main catchment basins of eastern Australia as per Forbes' hypothesis, relevant to the 2017-2018 season. Note that a value of 1 equals Decile 7 rainfall.

Catchment Basin	Oct-Dec	Jan-Mar	Oct-Dec	Jan-Mar
Catchinent Basin	2016	2017	2017	2018*
Darling River	0.58	0.81	0.93	0.32
Lachlan/Murrumbidgee/Murray Rivers	0.92	1.01	1.15	3.00
Northern Rivers	0.98	1.03	0.81	1.29
North Lake Eyre system	1.09	0.73	0.75	0.47

*January data only

ii. Nichol's Hypothesis

Table 2. The seasonal atmospheric pressures (in mm) according to Nichol's hypothesis, relevant to the 2017-2018 season.

	Autumn 2017	Winter 2017	Spring 2017
2017 Value	1009.60	1013.23	1009.70
Pre past MVEV seasons	<1009.74	<1012.99	<1009.99

Only the Winter period pertaining to the Nichol's hypothesis is <u>not</u> in line with past MVEV active years.



Arboviral Isolates

LOCATION - Site	Date Trapped	Mosquito Species	Virus
GRIFFITH – Lake Wyangan	3/Jan/2018	Culex annulirostris	Ross River
GRIFFITH – Lake Wyangan	31/Jan/2018	Culex annulirostris	Ross River
GRIFFITH – Hanwood	5/Feb/2018	Culex annulirostris	Ross River

*Detection via Honey-Baited Cards, the mosquito species cannot be determined.

Exotic Detections

Over the last eight years there have been an increasing number of detections of exotic mosquitoes at major Australian ports. The main species have been the dengue mosquito, *Aedes aegypti*, and the Asian Tiger Mosquito, *Aedes albopictus*. Both of these pose a serious biosecurity risk to Australia being major vectors of several arboviruses including Dengue, Yellow Fever, Zika, and Chikungunya viruses.

The first detection of an exotic mosquito at Sydney Airport occurred in January 2016, and since then, further detections have occurred on a number of occasions. The majority of the detections have been *Aedes aegypti* and these were largely collected within the International Airport precinct.

On 7/Feb/2018, two female *Aedes aegypti* were detected in a BG trap located in a freight handling facility situated near the domestic terminal (see Map 2). Further detections occurred shortly afterwards in the same facility and these are outlined below.

Date	Result
7/Feb/2018	2º Aedes aegypti
10/Feb/2018	1ð Aedes aegypti
11/Feb/2018	1º Aedes aegypti

Detections of Aedes aegypti during February, 2018.

In response, the NSW Ministry of Health convened an emergency teleconference with key stakeholders, insecticidal treatments undertaken, surveillance measures where enhanced, and a survey of the facility and surrounds was undertaken to identify and potential mosquito breeding sites. This survey is now completed and the recommendations are in the process of being implemented.



Human Notifications

Weekly notifications of human mosquito-borne diseases infections are available from the NSW Ministry of Health, Communicable Disease Weekly Report and summarized in the Table below* (<u>www.health.nsw.gov.au/Infectious/reports/Pages/CDWR.aspx</u>).

Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
1-Jul-17	14	6	3	2	0	0	25
8-Jul-17	6	4	0	4	1	0	15
15-Jul-17	8	0	2	1	0	0	11
22-Jul-17	10	3	7	2	0	0	22
29-Jul-17	6	0	2	2	0	0	10
5-Aug-17	8	0	4	0	0	0	12
12-Aug-17	11	1	3	2	5	0	22
19-Aug-17	5	2	1	2	2	0	12
26-Aug-17	6	3	3	2	0	1	15
2-Sep-17	6	0	1	0	1	0	8
9-Sep-17	14	0	1	2	1	0	18
16-Sep-17	9	1	5	0	0	0	15
23-Sep-17	9	1	3	1	0	0	14
30-Sep-17	7	0	1	1	1	0	10
7-Oct-17	7	0	3	2	0	0	12
14-Oct-17	10	1	2	1	0	0	14
21-Oct-17	11	2	8	2	1	0	24
28-Oct-17	16	1	6	1	1	0	25
4-Nov-17	14	3	7	3	1	0	28
11-Nov-17	5	2	7	0	0	0	14
18-Nov-17	3	2	10	0	0	0	15
25-Nov-17	9	2	6	1	1	0	19
2-Dec-17	14	1	8	0	0	0	23
9-Dec-17	9	0	3	1	0	0	13
16-Dec-17	9	4	2	1	2	0	18
23-Dec-17	7	0	6	0	0	0	13
30-Dec-17	5	0	1	0	0	0	6
Total	238	39	105	33	17	1	433

Table 4. Notifications of Mosquito-Borne Disease in NSW, 2017-2018*

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland. *The data in this table is updated once available from the NSW Ministry of Health.

Comment: It should also be noted that notifications are for NSW residents and that infection may have been acquired elsewhere and that winter notifications of RRV are likely to be false positives.



Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
6-Jan-18	5	0	4	2	1	0	12
13-Jan-18	2	2	13	1	0	0	18
20-Jan-18	6	0	9	0	1	0	16
27-Jan-18	3	0	10	1	0	0	14
3-Feb-18	9	3	8	1	0	0	21
10-Feb-18	0	2	0	0	0	0	2
Total	263	46	149	38	19	1	488

Table 4 cont. Notifications of Mosquito-Borne Disease in NSW, 2017-2018*

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland. *The data in this table is updated once available from the NSW Ministry of Health.



Table 5. Ross River virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2013 to January 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2013- 2014	36	23	27	36	30	30	33	35	44	72	86	57	509
2014- 2015	38	50	46	67	59	90	117	305	431	264	102	50	1,619
2015- 2016	54	61	53	61	70	54	42	60	78	79	52	16	680
2016- 2017	12	11	20	17	38	216	429	274	200	142	174	89	1,622
2017- 2018	29	37	51	55	35	31	24	8					270
Ave [†]	36	37	37	46	51	101	156	169	188	140	107	55	1,121

*updated 16/Feb/2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).

⁺Average for 2013/14 to 2016/17.

Table modified from: http://www1.health.nsw.gov.au/IDD/#/ROSS

Table 6. Barmah Forest virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2014 to January 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2014- 2015	10	3	11	11	8	4	12	17	43	43	16	11	189
2015- 2016	6	9	7	9	6	3	4	5	2	3	10	2	66
2016- 2017	4	3	0	0	1	9	9	5	8	6	24	24	93
2017- 2018	8	10	6	8	8	6	4	3					53
Ave [†]	7	5	6	7	5	5	8	9	18	17	17	12	116

*updated 16/Feb/2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).


⁺Average for 2014/15 to 2016/17.

Table modified from: <u>http://www1.health.nsw.gov.au/IDD/#/BF</u>



Mosquito Results

Mosquito abundances are best described in relative terms, and in keeping with the terminology from previous NSWASP Annual Reports, mosquito numbers are depicted on the tables below as:



Each location represents the average for all trapping sites at that location



Inland

Location	Meeswite	Oct	-17				Nov	1			Dec	;				Jar	า-18			Feb)			Mar	,		
Location	Mosquito	1	8	15	22	29	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25
Albury	Cx. annul																										
Albuly	Total Mosq.																										
	-		•	-	-	•									•		-			•		•	•	•			
Bourke	Cx. annul																										
Bourke	Total Mosq.																										
	-			•	•	_														_							
Griffith	Cx. annul																										
	Total Mosq.																										
	-			•	•						-									_							_
Leeton	Cx. annul																										
	Total Mosq.																										
	•	I					1		•	1	1	1				1			1		1				1		-
Macquarie																											
Marshes	Total Mosq.																										
	•	I					1		1	1	1	1				1			1		1				1		-
Mathoura	Cx. annul																										
	Total Mosq.																										
	1																										
Wagga	Cx. annul																										
	Total Mosq.																										



Coastal

Location	Magnuita	Nov	(De	С				Jai	n-18			Feb				Ма	ar			Ар	r			
Location	Mosquito	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25	1	8	15	22	29
Ballina	Ae. vigilax																										
Daililla	Total Mosq.																										
	Ae. vigilax																										
Harbour	Total Mosq.																										
			-		-		-									-							-				
Gosford	Ae. vigilax																										
	Total Mosq.																										
			-		-		-									•							-				
Lake	Ae. vigilax																										
Macquarie	Total Mosq.																										
			-		-		-									•							-				
Port	Ae. vigilax																										
Macquarie	Total Mosq.																										
			-		-		-	-															-				
Tweed	Ae. vigilax																										
I WEEU	Total Mosq.																										
		_		-					-					_				-									
Wyong	Ae. vigilax																										
wyong	Total Mosq.																										



Sydney

Location	Meenvite	Νον	/			De	С				Ja	n-18			Feb				Ма	ar			Ар	r			
Location	Mosquito	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25	1	8	15	22	29
Banks-	Ae. vigilax																										
town	Total Mosq.																										
Blacktown	Ae. vigilax Total Mosq																										
DIACKLOWII	Total Mosq.																										
		-	-		-		-	-	-	-		-			_					-		-	-				
Georges	Ae. vigilax																										
River	Total Mosq.																										
Hawkes-	Ae. vigilax																										
bury	Total Mosq.																										
Hills Shire	Ae. vigilax																										
	Total Mosq.																										
Penrith	Ae. vigilax																										
Femini	Total Mosq.																										
			-		-																						
Sydney	Ae. vigilax																										
Olympic Park	Total Mosq.																										
Dude	Ae. vigilax																										
Ryde	Total Mosq.																										



Sentinel Chicken Flocks

Location	Oct	-17				Νον				Dec					Jan-18				Feb				Mar			
	1	8	15	22	29	5	12	19	26	3	10	17	21	28	7	14	21	28	4	11	18	25	4	11	18	25
Bourke																										
Deniliquin						15N	14N	15N	14N	14N	15N															
Dubbo						15N	15N	15N	15N	15N	14N	14N	14N	14N	14N	14N	14N	14N	14N							
Forbes						15N		15N	15N	15N	14N							15N								
Griffith					15N		15N	15N	15N	15N	15N															
Hay					15N			15N	15N	15N	15N	15N														
Leeton						15N	15N		15N	14N	14N	14N	14N													
Macquarie Marshes							15N	15N	15N	15N	15N		15N	15N		15N	15N	15N								
Menindee										15N	15N	15N	15N	15N	15N	15N	15N	15N	14N							
Moama																										
Moree										15N	15N		15N	15N	15N	15N	15N	15N	15N							
Wee Waa																										

N= Negative for MVEV & KUNV

Prepared by: Stephen Doggett, Manager, Department of Medical Entomology, NSW Health Pathology (ICPMR), Westmead Hospital NSW 2145. Email: <u>Stephen.Doggett@health.nsw.gov.au</u>

Please note that these results remain the property of the NSW Ministry of Health and may not be used or disseminated to unauthorized persons or organizations without permission.

