NSW Arbovirus Surveillance & Mosquito Monitoring Program, 2017-2018

Weekly Update: 13/Apr/2018





creating better health & justice systems www.pathology.health.nsw.gov.au

Contents

Summary	2
Comment	2
Environmental Conditions	3
Rainfall	3
Three Month Rainfall & Temperature Forecast	3
Tidal	4
MVEV Climatic Models	5
Forbes' Hypothesis	5
Nichol's Hypothesis	5
Arboviral Isolates	6
Exotic Detections	6
Human Notifications	7
Monthly RRV notifications	9
Monthly BFV notifications	9
Mosquito Results	10
Inland	11
Coastal	12
Sydney	13
Sentinel Chicken Flocks	14



Summary

- **Climate**: over the last week, only the far north coast received any rainfall and this just a light drizzle. For March, most of the state had below average rainfall with only a pocket of the mid-north coast had above average precipitation.
- Three Month Forecast: rainfall predictions for NSW are for average precipitation for most of the state, with a slight chance of exceeding the average for the far south coast. Maximum and minimum temperatures are expected to be around average, although minimum temperatures are predicted to be above average along the Murray. According to the BOM as 10/Apr/2018, the El Niño–Southern Oscillation remains neutral.
- **Tidal**: the next series of tides that may trigger *Aedes vigilax* hatching are due over 16-22/Apr with tides of up to 1.86m predicted.
- **MVEV models**: the data relevant to both the Forbes' and Nichols' hypotheses have been updated to the end of Mar 2018. Neither model are suggestive of an MVEV epidemic.
- Mosquito Numbers Inland: all inland surveillance activities have ceased for this season.
- **Mosquito Numbers Coast**: collections were 'high' from almost all trapping sites, with few *Aedes vigilax* collected this week.
- **Mosquito Numbers Sydney**: mosquito numbers were down this week, although Homebush Bay continues to produce 'high' numbers.
- **Arboviral Isolates**: there were no further arboviral detections this week.
- Chicken Sentinel Flocks: all inland surveillance activities have ceased for this season.
- Human Notifications: for the current fiscal year, there have been 368 RRV and 74 BFV notifications, this is below the average compared with past years (the prior 18 season averages to the end of March are 406RRV and 240BFV).

Comment: Over the last week temperatures have been above average by up to 10 degrees for large parts of the state and this can only prolong the mosquito season. In fact, mosquito numbers remain 'high' at many sites along the coast. This coincides with the school holidays next week, when people are more likely to be outdoors and exposed to mosquitoes. For the coast, this time of the year is typically when peak virus transmission occurs and we can expect human notifications to go up, with the elevated mosquito numbers.

Holiday-makers enjoying outdoor activities, particularly camping and fishing, should cover-up and use insect repellent containing Diethyl Toluamide (DEET) or Picaridin. The NSW Health factsheet on avoiding mosquito bites is available at:


http://www.health.nsw.gov.au/environment/factsheets/Pages/mosquito-biteavoidance.aspx



Environmental Conditions

Rainfall

Rainfall across Australia for the week ending 12/Apr/2018 is depicted on the left and monthly rainfall deciles for March 2018 are on the right. Over the last week, there has been no rainfall for the majority of the state, with only a drizzle in the far north coast. For March, rainfall was below to very much below average across southern inland regions and around average for the remainder of the state. Only a pocket of the mid-north coast had above average precipitation. Maximum and minimum temperatures for March were 1-2 degrees above average.



Three Month Rainfall & Temperature Forecast

For May to July 2018, rainfall predictions for NSW are for average precipitation for most of the state, with a slight chance of exceeding the average for the far south coast. Maximum and minimum temperatures are expected to be around average, although minimum temperatures are predicted to be above average along the Murray. The following pages contain graphics of the seasonal outlook:

<u>www.bom.gov.au/climate/outlooks/#/rainfall/median</u> (Rainfall outlook). <u>www.bom.gov.au/climate/outlooks/#/temperature/summary</u> (Max & min temperature outlook).


According to the BOM as of 10/Apr/2018, the El Niño–Southern Oscillation remains neutral. The Indian Ocean Dipole (IOD) remains neutral. This all suggests that rainfall patterns are likely to be average over the upcoming months.

For more information: <u>www.bom.gov.au/climate/enso/</u> and, <u>http://www.bom.gov.au/climate/iod/</u>



Tidal

Tidal information is relevant for the prediction of the activity of the salt marsh mosquito, *Aedes vigilax*. Typically for NSW, tides of over 1.8m, as measured at Sydney, can induce hatching of *Aedes vigilax* larvae and the graph below of predicted tide heights can provide some indication of when this is likely to occur. Note this height various between regions, thus at Batemans Bay, a tide height over 0.8m can trigger egg hatching.



The next series of tides that may trigger *Aedes vigilax* hatching are due over 16-22/Apr with tides of up to 1.86m predicted.

Note that actual tide heights can vary by 0.3m (or more in unusual circumstances) due to variations in atmospheric pressure, rainfall, wind and other climatic phenomena. Climate change will also result in much higher tide heights. Thus predicted tide height should be used as a gauge only for potential *Aedes vigilax* activity. The larvae of the saltmarsh mosquito relies on a inundation/drying cycle for the mudflats in which it lives; continual wet weather prevents the drying cycles thereby reducing larval production.



MVEV Climatic Models

Three predictive environmental based models for MVEV activity have been developed; the Forbes (which relies on rainfall in the river catchment basins of Eastern Australia), Nichols (based on the Southern Oscillation), and the Bennett theory (based on the Indian Ocean Dipole). The latter theory is poorly developed (and unreliable), and is not considered below. Note that all the predictive models have been developed on a limited data set and do not always forecast activity. There can also be unusual environmental conditions that may lead to the introduction of the virus to southeastern Australia, such as the movement of low pressure cells from the north to the south of the country during 2008 and 2011. Vertical transmission of the virus (from adult to the egg in *Aedes* species) can result in restricted activity following localized heavy precipitation (as per 2003 at Menindee).

i. Forbes' Hypothesis

Rainfall was not above Decile 7 in all of the river catchment basins in eastern Australia for the last quarter of 2016, the first quarter of 2017, the last quarter of 2017 or the first quarter of 2018 (Table 1).

Table 1. Rainfall indices for the main catchment basins of eastern Australia as per Forbes' hypothesis, relevant to the 2017-2018 season. Note that a value of 1 equals Decile 7 rainfall.

Catchment Basin	Oct-Dec	Jan-Mar	Oct-Dec	Jan-Mar
Catchinent Basin	2016	2017	2017	2018
Darling River	0.58	0.81	0.93	0.52
Lachlan/Murrumbidgee/Murray	0.92	1.01	1.15	0.70
Rivers			_	
Northern Rivers	0.98	1.03	0.81	1.07
North Lake Eyre system	1.09	0.73	0.75	0.69

ii. Nichol's Hypothesis

Table 2. The seasonal atmospheric pressures (in mm) according to Nichol's hypothesis, relevant to the 2017-2018 season.

	Autumn 2017	Winter 2017	Spring 2017
2017 Value	1009.60	1013.23	1009.70
Pre past MVEV seasons	<1009.74	<1012.99	<1009.99

Only the Winter period pertaining to the Nichol's hypothesis is <u>not</u> in line with past MVEV active years.



Arboviral Isolates

LOCATION - Site	Date Trapped	Mosquito Species	Virus
GRIFFITH – Lake Wyangan	3/Jan/2018	Culex annulirostris	Ross River
GEORGES RIVER - Deepwater	30/Jan/2018	*	Stratford
GRIFFITH – Lake Wyangan	31/Jan/2018	Culex annulirostris	Ross River
GRIFFITH – Hanwood	5/Feb/2018	Culex annulirostris	Ross River
GEORGES RIVER – Alfords Point	7/Feb/2018	*	Stratford
GEORGES RIVER - Deepwater	12/Feb/2018	Aedes vigilax	Stratford
CENTRAL COAST – Empire Bay	27/Feb/2018	*	Barmah Forest
CENTRAL COAST – Halekulani	14/Mar/2018	*	Barmah Forest
GEORGES RIVER – Alfords Point	21/Mar/2018	*	Barmah Forest
GEORGES RIVER – Deepwater	21/Mar/2018	*	Barmah Forest

*Detection via PCR on pooled samples; the mosquito species cannot be determined.



Human Notifications

Weekly notifications of human mosquito-borne diseases infections are available from the NSW Ministry of Health, Communicable Disease Weekly Report and summarized in the Table below* (<u>www.health.nsw.gov.au/Infectious/reports/Pages/CDWR.aspx</u>).

Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
1-Jul-17	14	6	3	2	0	0	25
8-Jul-17	6	4	0	4	1	0	15
15-Jul-17	8	0	2	1	0	0	11
22-Jul-17	10	3	7	2	0	0	22
29-Jul-17	6	0	2	2	0	0	10
5-Aug-17	8	0	4	0	0	0	12
12-Aug-17	11	1	3	2	5	0	22
19-Aug-17	5	2	1	2	2	0	12
26-Aug-17	6	3	3	2	0	1	15
2-Sep-17	6	0	1	0	1	0	8
9-Sep-17	14	0	1	2	1	0	18
16-Sep-17	9	1	5	0	0	0	15
23-Sep-17	9	1	3	1	0	0	14
30-Sep-17	7	0	1	1	1	0	10
7-Oct-17	7	0	3	2	0	0	12
14-Oct-17	10	1	2	1	0	0	14
21-Oct-17	11	2	8	2	1	0	24
28-Oct-17	16	1	6	1	1	0	25
4-Nov-17	14	3	7	3	1	0	28
11-Nov-17	5	2	7	0	0	0	14
18-Nov-17	3	2	10	0	0	0	15
25-Nov-17	9	2	6	1	1	0	19
2-Dec-17	14	1	8	0	0	0	23
9-Dec-17	9	0	3	1	0	0	13
16-Dec-17	9	4	2	1	2	0	18
23-Dec-17	7	0	6	0	0	0	13
30-Dec-17	5	0	1	0	0	0	6
Total	238	39	105	33	17	1	433

Table 4. Notifications of Mosquito-Borne Disease in NSW, 2017-2018*

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland. *The data in this table is updated once available from the NSW Ministry of Health.

Comment: It should also be noted that notifications are for NSW residents and that infection may have been acquired elsewhere and that winter notifications of RRV are likely to be false positives.



Week Ending	RRV	BFV	DENV [†]	Malaria [†]	CHIKV [†]	ZIKV [†]	Total
6-Jan-18	5	0	4	2	1	0	12
13-Jan-18	2	2	13	1	0	0	18
20-Jan-18	6	0	9	0	1	0	16
27-Jan-18	3	0	10	1	0	0	14
3-Feb-18	9	3	8	1	0	0	21
10-Feb-18	8	2	6	0	0	0	16
17-Feb-18	4	2	3	0	0	0	9
24-Feb-18	15	1	4	1	1	1	23
3-Mar-18	9	2	3	6	0	0	20
10-Mar-18	19	3	6	0	1	0	29
17-Mar-18	10	1	3	0	0	0	14
24-Mar-18	10	4	1	2	0	0	17
31-Mar-18	8	1	6	0	0	0	16
7-Apr-18	14	2	2	0	0	0	18
Total	360	62	182	47	21	2	672

Table 4 cont. Notifications of Mosquito-Borne Disease in NSW, 2017-2018*

[†]All of these viruses are acquired overseas, although some DENV cases may be from North Queensland. *The data in this table is updated once available from the NSW Ministry of Health.



Table 5. Ross River virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2013 to February 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2013- 2014	36	23	27	36	30	30	33	35	44	72	86	57	509
2014- 2015	38	50	46	67	59	90	117	305	431	264	102	50	1,619
2015- 2016	54	61	53	61	70	54	42	60	78	79	52	16	680
2016- 2017	12	11	20	17	38	216	429	274	200	142	174	89	1,622
2017- 2018	29	37	52	56	37	31	30	40	45	11			368
Ave [†]	27	26	25	30	35	42	65	71	86	78	68	37	589

*updated 13/Apr/2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).

⁺Average for 2001/02 to 2016/17.

Table modified from: <u>http://www1.health.nsw.gov.au/IDD/#/ROSS</u>

Table 6. Barmah Forest virus infection notifications in NSW residents, by month of disease onset per fiscal year, July 2014 to February 2018*.

Year	Jul	Aug	Sep	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	Jun	Total
2014- 2015	10	3	11	11	8	4	12	17	43	43	16	11	189
2015- 2016	6	9	7	9	6	3	4	5	2	3	10	2	66
2016- 2017	4	3	0	0	1	9	9	5	8	6	24	24	93
2017- 2018	8	10	6	8	8	6	5	12	7	4			74
Ave [†]	21	19	18	22	25	21	32	35	48	51	49	28	367

*updated 13/Apr/2018 (this table is updated at different times to Table 4 above, hence there maybe differences in the numbers).


⁺Average for 2001/02 to 2016/17.

Table modified from: <u>http://www1.health.nsw.gov.au/IDD/#/BF</u>



Mosquito Results

Mosquito abundances are best described in relative terms, and in keeping with the terminology from previous NSWASP Annual Reports, mosquito numbers are depicted on the tables below as:



Each location represents the average for all trapping sites at that location



Inland

Location	Meessuite	Oct	-17				Nov	1			Dec	c				Jar	า-18			Feb)			Mar			
Location	Mosquito	1	8	15	22	29	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25
Albury	Cx. annul																										
Albuly	Total Mosq.																										
	-	-			-	•				•				-	-					•		•		-			
Bourke	Cx. annul																										
Bourke	Total Mosq.																										
	-	-			-	•								-	-		-		-	_		-	-	-			
Griffith	Cx. annul																										
ommun	Total Mosq.																										
		-		-	_	-				-						_			-	_			-				
Leeton	Cx. annul																										
Lecton	Total Mosq.																										
		-		-	_	-			_	-			-		-	-			-	-			-				
Macquarie	Cx. annul																										
Marshes	Total Mosq.																										
			_	-		-								-					_	-			_				
Mathoura	Cx. annul																										
	Total Mosq.																										
Wagga	Cx. annul																										
wayya	Total Mosq.																										



Coastal

Location	D.I. a model	Nov	(De	C				Jai	n-18			Feb				Ма	ar			Ар	r			
Location	Mosquito	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25	1	8	15	22	29
Ballina	Ae. vigilax																										
Dallilla	Total Mosq.																										
Coffs	Ae. vigilax																										
Harbour	Total Mosq.																										l
			-		-	-				_	-	-	-	_	_	_		_		-			-		_		
Gosford	Ae. vigilax																										
0031010	Total Mosq.																										
Lake	Ae. vigilax																										
Macquarie	Total Mosq.																										
Port	Ae. vigilax																										
Macquarie	Total Mosq.																										
Tweed	Ae. vigilax																										
	Total Mosq.																										
Wyong	Ae. vigilax																										
vvyong	Total Mosq.																										



Sydney

	Magguita	Nov	/			De	С				Jai	n-18			Feb				Ма	ar			Ар	r			
Location	Mosquito	5	12	19	26	3	10	17	24	31	7	14	21	28	4	11	18	25	4	11	18	25	1	8	15	22	29
Banks-	Ae. vigilax																										
town	Total Mosq.																										
						-				-	-	-	-						-	-	-	-					
Blacktown	Ae. vigilax																										
Diacktown	Total Mosq.																										
	-			-	-		-			-	•	•			-			-		-	_						
Georges	Ae. vigilax																										
River	Total Mosq.																										
Hawkes-	Cx. annul																										
bury	Total Mosq.																										
Hills Shire	Ae. vigilax																										
	Total Mosq.																										
Penrith	Ae. vigilax																										
Ferriti	Total Mosq.																										
Sydney Olympic	Ae. vigilax																										
	Total Mosq.																										



Sentinel Chicken Flocks

Location	Oct	-17				No	V			Dec	;				Jan	-18			Feb)			Mai	r		
Location	1	8	15	22	29	5	12	19	26	3	10	17	21	28	7	14	21	28	4	11	18	25	4	11	18	25
Deniliquin						15N	14N	15N	14N	14N	15N	15N	15N	15N	15N	14N	15N	14N	11N							
Dubbo						15N	15N	15N	15N	15N	14N			14N												
Forbes						15N		15N	15N	15N	14N							15N	15N							
Griffith					15N		15N	14N	14N	14N																
Hay					15N			15N	15N	15N	15N	15N		15N	15N	15N	15N		15N							
Leeton						15N	15N		15N	14N	13N	14N														
Macquarie Marshes							15N	15N	15N	15N	15N		15N	15N		15N	15N	15N		15N	15N	15N	15N		15N	14N
Menindee										15N	14N	15N	16N	15N	15N	15N	15N	15N								
Moree										15N	15N		15N													

N= Negative for MVEV & KUNV

Prepared by: Stephen Doggett, Manager, Department of Medical Entomology, NSW Health Pathology (ICPMR), Westmead Hospital NSW 2145. Email: <u>Stephen.Doggett@health.nsw.gov.au</u>

Please note that these results remain the property of the NSW Ministry of Health and may not be used or disseminated to unauthorized persons or organizations without permission.

SHPN: (EH)170602

